arXiv:1402.0388v1 [math.PR] 3 Feb 2014

CONVERGENCE OF CLOCK PROCESSES AND AGING IN METROPOLIS DYNAMICS OF A TRUNCATED REM

VÉRONIQUE GAYRARD

ABSTRACT. We study the aging behavior of a truncated version of the Random Energy Model evolving under Metropolis dynamics. We prove that the natural time-time correlation function defined through the overlap function converges to an arcsine law distribution function, almost surely in the random environment and in the full range of time scales and temperatures for which such a result can be expected to hold. This establishes that the dynamics ages in the same way as Bouchaud's REM-like trap model, thus extending the universality class of the latter model. The proof relies on a clock process convergence result of a new type where the number of summands is itself a clock process. This reflects the fact that the exploration process of Metropolis dynamics is itself an aging process, governed by its own clock. Both clock processes are shown to converge to stable subordinators below certain critical lines in their time-scale and temperature domains, almost surely in the random environment.

1. INTRODUCTION AND MAIN RESULTS

As evidenced by an extensive body of experiments, glassy systems are never in equilibrium on laboratory time scales [12], [33]; instead, their dynamics become increasingly slower as time elapses. Termed *aging*, this pattern of behavior was most successfully accounted for, at a theoretical level, by Bouchaud's phenomenological trap models [11], [13]. These are effective dynamics that, reviving ideas of Goldstein *et al.* [28], model the long time behavior of spin glass dynamics in terms of thermally activated barrier crossing in a state space reduced to the configurations of lowest energy (see [12] for a review). Main examples of microscopic systems that trap models aim to describe are Glauber dynamics on state spaces $\{-1,1\}^n$ reversible with respect to the Gibbs measures associated to random Hamiltonians of mean-field spin glasses, such as the Random Energy Model (REM) and p-spin SK models [19], [20]. The link between such dynamics and their associated trap models is, however, simply postulated.

When trying to establish this link rigorously, a main question that arises is what Glauber dynamics to choose. While classical choices are Metropolis [31] or Heath-Bath dynamics [27], most of the focus so far was on the so-called *Random Hopping* dynamics whose transition rates do not depend on the variation of energy along a given transition but only on the energy of its starting point [4], [5], [6], [2], [25], [14], [15], [8], [23]. Although physically unrealistic, the relative simplicity of this choice allowed important insights to be gained: a rigorous justification of the connection between the REM dynamics and trap models was given, first on times scales close to equilibrium [3, 4, 5], later also on shorter

Date: February 4, 2014.

2000 *Mathematics Subject Classification.* 82C44,60K35,60G70.

Key words and phrases. random dynamics, random environments, clock process, Lévy processes, spin glasses, ageing, Metropolis dynamics .

I would like to thank the Hausdorff Research Institute for Mathematics and the Institut Henri Poincaré where part of this work was carried out.

(but still exponential in n) time scales [6], and these results were partially extended to the p-spin SK models [2] on a sub-domain of times scales, albeit only in law with respect to the random environment and for $p \geq 3$. The SK model itself ($p = 2$) could be dealt with on time scales that are sub-exponential in n and again in law with respect to the random environment [8]. A variant of the so-called Bouchaud's asymmetric dynamics in which the asymmetry parameter tends to zero as $n \uparrow \infty$ is considered in [30] for the REM.

Beyond model-based analysis, a general aging mechanism was isolated that linked aging to the arcsine law for subordinators through the asymptotic behavior of a partial sum process called *clock process*. First implemented in [6] in the setting of Random Hopping dynamics this mechanism was revisited in [26] and [14] where, using a method developed by Durrett and Resnick [22] to prove functional limit theorems for dependent random variables, simple and robust criteria for convergence of clock processes to subordinators were given, suited for dealing with general Glauber dynamics. Applied to the Random Hopping dynamics of the REM [25] and p-spin SK models [14], [15], these criteria allowed to improve all earlier results, turning statements previously obtained in law into almost sure statements in the random environment.

In the present paper the approach of [26] is applied to Metropolis dynamics of the REM for which it was primarily intended, although only for a truncated version of the REM Hamiltonian. While the ultimate goal is of course to deal with the full REM, the truncated model does captures a number of features that are present in the activated dynamics of the full model, and enables us to clarify a number of issues on a problem for which nothing is known at a theoretical level and no computer simulations are available.

1.1. **The setting.** Before entering into the details, let us specify the model. Denote by $\mathcal{V}_n = \{-1, 1\}^n$ the n-dimensional discrete cube and let $(g(x), x \in \mathcal{V}_n)$ be a collection of independent standard Gaussian random variables, defined on a common probability space $(\Omega, \mathcal{F}, \mathbb{P})$. We will refer to these Gaussians as to the *random environment.* The Hamiltonian or energy function of the standard REM simply is the random funtion

$$H_n^{\mathsf{REM}}(x) \equiv \sqrt{n} g(x)\,, \quad x \in \mathcal{V}_n. \tag{1.1}$$

Given a sequence $u_n > 0$ (our *truncation level*) the *truncated* REM Hamiltonian then is

$$H_n(x) \equiv \begin{cases} \sqrt{n} g(x), & \text{if } \ g(x) \leq -u_n, \\ 0, & \text{else;} \end{cases}\,, \quad x \in \mathcal{V}_n. \tag{1.2}$$

Here we follow the physical convention that the configurations of minimal energy are the most stable ones, that is to say, Gibbs measure at inverse temperature $\beta > 0$ is defined as

$$G_{\beta,n}(x) = e^{-\beta H_n(x)} / (\textstyle\sum_{x \in \mathcal{V}_n} e^{-\beta H_n(x)})\,, \quad x \in \mathcal{V}_n\,. \tag{1.3}$$

We are interested in the single spin-flip continuous time Metropolis dynamics for this model. This is a Markov jump process $(X_n(t), t > 0)$ on $\mathcal{V}_n$ that is usually defined through its jump rates, given by

$$\lambda_n(x, y) = \begin{cases} \frac{1}{n} e^{-\beta [H_n(y) - H_n(x)]^+}, & \text{if } (x, x') \in \mathcal{E}_n, \\ 0, & \text{else;} \end{cases} \tag{1.4}$$

where $a+ = \max\{a, 0\}$, $\mathcal{E}_n = \{(x, y) \in \mathcal{V}_n \times \mathcal{V}_n \,:\, \mathrm{dist}(x, y) = 1\}$ is the set of edges of $\mathcal{V}_n$, and $\mathrm{dist}(x, x') \equiv \frac{1}{2} \sum_{i=1}^n |x_i - x'_i|$ is the graph distance on $\mathcal{V}_n$.

Equivalently, X_n can be defined as a time change of its *jump chain*, namely, the discrete time chain, J_n, that describes the trajectories of X_n, through the relation

$$X(t) = J(\widetilde{S}_n^{\leftarrow}(t)), \quad t \geq 0, \tag{1.5}$$

where $\widetilde{S}_n^{\leftarrow}$ denotes the generalized right continuous inverse of $\widetilde{S}_n$, and $\widetilde{S}_n$, the so-called *clock process*, is the partial sum process that records the total time spent by X_n along the trajectories of J_n. Spelling out these objects explicitly, the jump chain is the Markov chain $(J_n(i), i \in \mathbb{N})$ on $\mathcal{V}_n$ with one-step transition probabilities

$$p_n(x,y) = \frac{e^{-\beta[H_n(y)-H_n(x)]^+}}{\sum_{y:(x,y)\in\mathcal{E}_n} e^{-\beta[H_n(y)-H_n(x)]^+}}, \quad \text{if } (x,y) \in \mathcal{E}_n, \tag{1.6}$$

and $p_n(x,y) = 0$ otherwise, and the clock process is given by

$$\widetilde{S}_n(k) = \sum_{i=0}^{k-1} \lambda_n^{-1}(J_n(i)) e_{n,i}, \quad k \geq 1, \tag{1.7}$$

where $(e_{n,i}, n \in \mathbb{N}, i \in \mathbb{N})$ is a collection of independent mean one exponential random variables, independent of J_n, and the $\lambda_n(\cdot)$'s are the classical holding time parameters

$$\lambda_n(x) \equiv \frac{1}{n} \sum_{y:(x,y)\in\mathcal{E}_n} e^{-\beta[H_n(y)-H_n(x)]^+}, \quad \forall x \in \mathcal{V}_n. \tag{1.8}$$

In the clock process-based aging mechanism, one aims to infer knowledge of the aging behavior of X_n as $n \uparrow \infty$ from the asymptotic behavior of the properly rescaled clock process, using relation (1.5). To formulate this more precisely let $K_n(t)$ be a nondecreasing right continuous function with range $\{0, 1, 2, \dots\}$ and let c_n be a nondecreasing sequence. Both $K_n(t)$ and c_n are time scales. Consider the re-scaled clock process

$$S_n(t) = c_n^{-1} \widetilde{S}_n(K_n(t)), \quad t \geq 0. \tag{1.9}$$

This is a doubly stochastic object: one the one hand, for each fixed realization of the random environment (that is, of the random Hamiltonian H_n), S_n is a partial sum process with increasing paths that increase only by jumps and whose increments depend on the $J_n(i)$'s and the $e_{n,i}$'s; on the other hand, both the $\lambda_n(\cdot)$'s and the law of J_n depend on the random environment. One then asks whether there exist time scales $K_n(t)$ and c_n that make S_n converge weakly, as $n \uparrow \infty$, as a sequence of random elements in Skorokhod's space $D((0,\infty])$, $\mathbb{P}$-almost surely in the random environment. Such a result will be useful for deriving aging information if it enables one to control the behavior of the two-time correlation functions that are used in theoretical physics to quantify this phenomenon, the natural choice in mean-field models being the two-time overlap function

$$\mathcal{C}_n(t,s) = n^{-1}\big(X_n(c_n t), X_n(c_n(t+s)\big) \tag{1.10}$$

where $(\cdot,\cdot)$ denotes the inner product in $\mathbb{R}^n$. Clearly, how successful this can be strongly depends on the topology on the space $D((0,\infty])$ in which weak convergence of S_n is obtained. *Normal* aging is then said to occur if, for some convergence mode,

$$\lim_{n\to\infty} \mathcal{C}_n(t,s) = \mathcal{C}_\infty(t/s) \tag{1.11}$$

for some non trivial function $\mathcal{C}_\infty$ (see [26] for more general aging behaviors). The key idea put forward in [6] is that if S_n converges to an α-stable subordinator with $\alpha \in (0,1)$ then (1.11) is nothing but a manifestation of the self-similarity of such subordinators, as captured by the Dynkin-Lamperti arcsine law Theorem.

For future reference, we denote $\mathcal{F}^J$ and $\mathcal{F}^X$, respectively, the σ-algebra generated by the variables J_n and X_n. We write P for the law of the process J_n, conditional on the σ-algebra $\mathcal{F}$, i.e. for fixed realizations of the random environment. Likewise we call $\mathcal{P}$ the law of X_n conditional on $\mathcal{F}$. If the initial distribution, say μ_n, has to be specified we write $\mathcal{P}_{\mu_n}$ and P_{μ_n}. Expectation with respect to $\mathbb{P}$, P, and $\mathcal{P}$ are denoted by $\mathbb{E}$, E, and $\mathcal{E}$, respectively.

1.2. **Results.** We must now specify the truncation level in (1.2). Given $c_\star > 0$, we let $u_n \equiv u_n(c_\star)$ be the sequence defined through

$$\mathbb{P}(g(x) \leq -u_n(c_\star)) = n^{-c_\star}. \tag{1.12}$$

Viewing the vertices of $\mathcal{V}_n$ as independently occupied with probability (1.12), one sees that this probability increases from 0 to 1 as $c_\star$ decreases from $+\infty$ to 0, and so, the set of occupied vertices evolves from the empty set to the entire $\mathcal{V}_n$. Set

$$\mathcal{V}_n^\star \equiv \{x \in \mathcal{V}_n \mid x \text{ is occupied}\} \setminus I_n^\star, \tag{1.13}$$

where $I_n^\star$ is the set of isolated occupied vertices, namely, $x \in I_n^\star$ if it is occupied but none of its n neighbors is. Our results are closely tied to the graph properties of this set. Let us only mention here that $c_\star$ must be chosen such that $c_\star > c^{\mathsf{crit}} \geq 2$. This precludes the emergence of a giant connected component and guarantees that, $\mathbb{P}$- almost surely, the graph of $\mathcal{V}_n^\star$ is made of an exponentially large number ($\approx \mathcal{O}(2^n/n^{2c_\star - 1})$) of small, disjoint connected components of size smaller than n. In explicit form, the sequence u_n obeys

$$u_n(c_\star) = \sqrt{2c_\star \log n} - \left(\tfrac{\log\log n + \log 4\pi}{2\sqrt{2c_\star \log n}} + \mathcal{O}\Big(\tfrac{1}{\sqrt{\log n}}\Big)\right). \tag{1.14}$$

Hence, the truncation only prunes energies such that $-H_n^{\mathsf{REM}}(x) \lesssim \sqrt{2c_\star n \log n}$, while activated aging typically involves energies of size $-H_n^{\mathsf{REM}}(x) \geq \gamma n$, $\gamma > 0$, that is to say, of the order of $\max_{x \in \mathcal{V}_n}(-H_n^{\mathsf{REM}}(x))$.

We are concerned with finding sequences c_n and K_n for which the rescaled clock process (1.9) converges for some (ideally, the smallest possible) $c_\star > c^{\mathsf{crit}}$. Note that in physical terms, c_n is the time scale on which the continuous time process X_n is observed, while $K_n(t)$ is the total number of steps the discrete time chain J_n takes during the period of observation. In all previously mentioned works on mean-field spin glasses (that is, the REM and p-spin SK models with $p \geq 3$) where convergence of (1.9) could be proved, this was on time scales of the form $c_n \sim \exp(\beta\gamma n)$, $\gamma > 0$. Furthermore, K_n invariably had to be chosen of the form $K_n(t) = \lfloor a_n t \rfloor$, where a_n is defined through $a_n \mathbb{P}(w_n(x) \geq c_n) \sim 1$, and where $w_n(x)$ denotes the Boltzmann weight of the considered model; in the standard REM, this is

$$w_n(x) \equiv \exp\{-\beta H_n^{\mathsf{REM}}(x))\}, \quad x \in \mathcal{V}_n. \tag{1.15}$$

Finally, a common α-stable subordinator emerged as the limit of the clock processes.

As might reasonably be expected, the physical time scale, c_n, on which activated aging occurs in Metropolis dynamics is the same as in the Random Hopping dynamics. What does differ, however, is the choice of K_n. Given a sequence a_n, we now set

$$K_n(t) \equiv \min\left\{ k \geq 1 \,\middle|\, \textstyle\sum_{i=0}^{k-1} \mathbb{1}_{\{J_n(i) \in \mathcal{V}_n \setminus \mathcal{V}_n^\star\}} = \lfloor a_n t \rfloor \right\}, \quad t \geq 0. \tag{1.16}$$

This is the number of steps J_n must take in order to take $\lfloor a_n t \rfloor$ steps outside $\mathcal{V}_n^\star$. Our first theorem states that the resulting rescaled clock process (1.9) converges to the same

limiting subordinator as in the Random Hopping dynamics, for the very same sequences a_n and c_n, and in the same β range. For $0 \le \varepsilon \le 1$ and $0 < \beta < \infty$, set

$$\beta_c(\varepsilon) = \sqrt{\varepsilon 2 \log 2}, \tag{1.17}$$

$$\alpha(\varepsilon) = \beta_c(\varepsilon)/\beta. \tag{1.18}$$

Throughout this paper the initial distribution is the uniform distribution on $\mathcal{V}_n \setminus \mathcal{V}_n^\star$.

Theorem 1.1. *Assume that $c_\star > 3$. Given $0 < \varepsilon < 1$ let a_n and c_n be defined through*

$$\lim_{n\to\infty} \frac{\log a_n}{n \log 2} = \varepsilon, \quad a_n \mathbb{P}(w_n(x) \ge c_n) \sim 1. \tag{1.19}$$

Then, for all $0 < \varepsilon < 1$ and all $\beta > \beta_c(\varepsilon)$, $\mathbb{P}$-almost surely,

$$S_n \Rightarrow_{J_1} S_\infty \tag{1.20}$$

where S_∞ is a stable subordinator with zero drift and Lévy measure ν defined through

$$\nu(u, \infty) = u^{-\alpha(\varepsilon)} \alpha(\varepsilon) \Gamma(\alpha(\varepsilon)), \quad u > 0, \tag{1.21}$$

and where $\Rightarrow_{J_1}$ denotes weak convergence in the space $D([0,\infty))$ of càdlàg functions equipped with the Skorokhod J_1-topology.

In the rest of the paper the symbol $\Rightarrow_{J_1}$ (sometimes only $\Rightarrow$) has the same meaning as in Theorem 1.1.

Let us now elucidate the meaning of K_n. There is a clear parallel between the definitions (1.16) and (1.7) of K_n and $\widetilde{S}_n$. Like $\widetilde{S}_n$, K_n is similar to a time, each step of the chain J_n lasting one time unit. Just like $\widetilde{S}_n$ also, it is a function of an underlying 'faster chain', namely, the chain J_n observed only at its visits to $\mathcal{V}_n \setminus \mathcal{V}_n^\star$. Thus K_n can be viewed as the total time spent by the chain J_n along the first $\lfloor a_n t \rfloor$ steps of that fast chain – in other words, as a clock process for J_n. One may probe this parallel further by asking if there exist sequences b_n for which the rescaled process $b_n^{-1} K_n$ converges. As the next theorem shows, the nature of the limit undergoes a transition at the critical value $\beta = 2\beta_c(\varepsilon/2)$.

Theorem 1.2. *Assume that $c_\star > 3$ and, given $0 < \varepsilon < 1$, let a_n be as in Theorem 1.1.*
(i) If $\beta > 2\beta_c(\varepsilon/2)$, let b_n be defined through $\sqrt{n a_n}\mathbb{P}(w_n(x) \ge (n-1)b_n) \sim 1$. Then, for all $0 < \varepsilon < 1$ and all $\beta > 2\beta_c(\varepsilon/2)$, $\mathbb{P}$-almost surely,

$$b_n^{-1} K_n \Rightarrow_{J_1} S_\infty^\dagger, \tag{1.22}$$

where $S_\infty^\dagger$ is a stable subordinator with zero drift and Lévy measure $\nu^\dagger$ defined through

$$\nu^\dagger(u, \infty) = u^{-2\alpha(\varepsilon/2)} 2\alpha(\varepsilon/2) \Gamma(2\alpha(\varepsilon/2)), \quad u > 0. \tag{1.23}$$

(ii) If $0 < \beta < 2\beta_c(\varepsilon/2)$, set $b_n = a_n \exp(n(\beta/2)^2)/(\beta\sqrt{\pi n})$. Then, for all $0 < \varepsilon < 1$ and all $\beta < 2\beta_c(\varepsilon/2)$, $\mathbb{P}$-almost surely,

$$(b_n^{-1} K_n(t), t \ge 0) \overset{\mathcal{P}-a.s.}{\underset{n\to\infty}{\longrightarrow}} (t, t \ge 0), \tag{1.24}$$

where convergence holds in the space $C([0,\infty))$ of continuous functions equipped with the topology of the uniform convergence on compact sets.

Remark. A transition similar to that of Theorem 1.2 is present in S_n at the critical value $\beta = \beta_c(\varepsilon)$. Since in the region $\beta < \beta_c(\varepsilon)$ activated aging is interrupted (and in order not to make the paper longer) we leave out the explicit statement.

The occurence of stable subordinators as limits of both S_n and $b_n^{-1}K_n$ above the critical lines $\beta = \beta_c(\varepsilon)$ and $\beta = 2\beta_c(\varepsilon/2)$, $0 < \varepsilon < 1$, respectively, can be explained through a single, universal mechanism which is best described as an exploration mechanism of a set of *extreme accessible states* whose effective waiting times are *heavy tailed*. What gives rise to this mechanism, however, is very different depending on whether one considers S_n or $b_n^{-1}K_n$. Let us briefly explain this.

When dealing with S_n, the processes at work are analogous to those already present in the Random Hopping dynamic of the REM: the set of extreme accessible states identifies with the vertices such that $w_n(x) \sim c_n$, and most such vertices belong to the set $\mathcal{I}_n^\star$ of isolated occupied vertices of (1.13), but J_n typically does not revisit the elements of $\mathcal{I}_n^\star$ twice so that the associated effective waiting times typically coincide with the exponential holding times $\lambda_n^{-1}(x)e_{n,i} = w_n(x)e_{n,i}$ (see (1.7)) and these, scaled down by c_n, are asymptotically heavy tailed with parameter $\alpha(\varepsilon)$.

This is in sharp contrast with the mechanisms that govern the behavior of $b_n^{-1}K_n$. Viewing the set $\mathcal{V}_n^\star \cup \mathcal{I}_n^\star$ as the level set of the REM's landscape, and its disjoint components as separated valleys, K_n can be interpreted as the sum of the sojourn times in the valleys of size ≥ 2 that J_n visits along its path. Thus holding times now arise dynamically from metastable trapping times in local valleys. The analysis of these times reveals that the set of extreme accessible states is the set of pairs $(x, y) \in \mathcal{E}_n$ such that $\min(w_n(x), w_n(y)) \sim b_n$, that their effective waiting times have exponential tails of mean value $\min(w_n(x), w_n(y))$, and that, scaled down by b_n, these waiting times are asymptotically heavy tailed with parameter $2\alpha(\varepsilon/2)$.

Below the critical line $\beta = 2\beta_c(\varepsilon/2)$, $0 < \varepsilon < 1$, this picture breaks down. The leading contributions to $b_n^{-1}K_n$ no longer come from extreme events but from *typical* events that consist of visits to valleys whose effective mean waiting times have finite mean values. Note that even here, the jump chain does not resemble the symmetric random walk. In fact, our results show that on the time scales of activated aging, Metropolis dynamics never can be reduced to the Random Hopping dynamics, just as the latter cannot be reduced to Bouchaud's phenomenological trap model. Despite this Bouchaud's trap model does correctly predict the aging behavior of both dynamics:

Theorem 1.3 (Correlation function)**.** *Let $\mathcal{C}_n(t, s)$ be defined in (1.10). Under the hypothesis of Theorem 1.1, for all $\rho \in (0, 1)$, $t > 0$ and $s > 0$, $\mathbb{P}$-almost surely,*

$$\lim_{n\to\infty} \mathcal{P}\big(\mathcal{C}_n(t, s) \geq 1 - \rho\big) = \frac{\sin \alpha\pi}{\pi} \int_0^{t/(t+s)} u^{\alpha(\varepsilon)-1}(1 - u)^{-\alpha(\varepsilon)}\, du. \tag{1.25}$$

Remark. The convergence statement of Theorem 1.2, (i), is of course is a manifestation of the fact that above the critical line the jump chain is itself an aging process. This can be quantified using e.g. the function $\mathcal{C}'_n(t, s) = n^{-1}\big(J_n(\lfloor b_n t\rfloor), J_n(\lfloor b_n(t + s)\rfloor\big)$ for which a statement similar to (1.25) can be proved with $2\alpha(\varepsilon/2)$ subsituted for $\alpha(\varepsilon)$.

Let us highlight the content of the next two sections. What we need to know about the random graph induced by the truncation (1.12) is collected in Section 2. In Section 3 we isolate two processes, called the *front end* and *back end clock processes* (hereafter FECP and BECP), that are central to the proofs of Theorem 1.1 and Theorem 1.2. We show that the processes S_n, respectively K_n, can be written as the sum of FECP, respectively BECP, and remainders. Based on this we decompose the proofs of Theorem 1.1 and Theorem 1.2 into proving on the one hand that FECP and BECP converge, and showing on the other hand that the remainders are asymptotically negligeable. This strategy strongly

relies on two abstract theorems (Theorem 8.2 in Section 8 and Theorem 9.1 in Section 9) that give sufficient conditions for FECP and BECP to converge to Lévy subordinators. The organisation of the rest of the paper is detailed at the end of Section 3.

2. Random Graph properties of the REM's landscape

Given $V \subseteq \mathcal{V}_n$ we denote by $G \equiv G(V)$ the undirected graph which has vertex set V and edge set consisting of pairs of vertices $\{x, y\}$ in V with $\mathrm{dist}(x, y) = 1$. This short section is concerned with the graph properties of the level sets

$$V_n(\rho) = \{x \in \mathcal{V}_n \mid w_n(x) \geq r_n(\rho)\}, \tag{2.1}$$

where, given $\rho > 0$, the truncation level $r_n(\rho)$ is the sequence defined through

$$2^{\rho n}\mathbb{P}(w_n(x) \geq r_n(\rho)) = 1. \tag{2.2}$$

This is a convenient reparametrization of (1.12), that is, (1.12) follows from (2.2) by taking

$$\rho = \rho_n^\star \equiv \frac{c_\star \log n}{n \log 2}, \quad r_n(\rho_n^\star) \equiv \exp(\beta\sqrt{n}u_n(c_\star)). \tag{2.3}$$

Viewing the vertices of $\mathcal{V}_n$ as independently occupied with probability $2^{-\rho n}$, questions on $G(V_n(\rho))$ reduce to questions on random subgraphs of the hypercube graph $\mathcal{Q}_n \equiv G(\mathcal{V}_n)$.

2.1. **Component structure of $V_n(\rho)$.** The set $V_n(\rho)$ of occupied vertices can be decomposed into components that we classify according to their connectedness and size. We call $C \subseteq V_n(\rho)$ a connected component of size $|C|$ if the subgraph $G(C) \subseteq G(V_n(\rho))$ is connected. All connected components have size ≥ 2. We call isolated occupied vertices of $V_n(\rho)$ components of size 1. Given $V_n(\rho)$, $\mathcal{V}_n$ can uniquely be decomposed into

$$\mathcal{V}_n = N_n(\rho) \cup I_n(\rho) \cup \left(\cup_{l=1}^{L} C_{n,l}(\rho)\right), \quad L \equiv L_n(\rho), \tag{2.4}$$

where $N_n(\rho)$ is the set of all non occupied vertices, $I_n(\rho)$ is the set of all isolated occupied ones, and $C_{n,l}(\rho)$, $1 \leq l \leq L$, is a collection of disjoint connected components satisfying

$$G(V_n(\rho)) = \cup_{l=1}^{L} G(C_{n,l}(\rho)), \quad C_{n,l}(\rho) \cap C_{n,k}(\rho) \ \ \forall l \neq k. \tag{2.5}$$

As ρ decreases, the set $V_n(\rho)$ grows and the graph $G(V_n(\rho))$ potentially acquires new edges. Little is known about such graphs compared to those obtained by selecting edges independently. It is chiefly known [10] that the size of the largest $C_{n,l}(\rho)$ undergoes a transion near the value $\rho \approx \frac{\log n}{n \log 2}$, with a unique "giant" componant of size $\mathcal{O}(n^{-1}2^n)$ emerging slightly below this value. We are interested here in choosing ρ in such a way as to garantee that the size of the largest $C_{n,l}(\rho)$ remains small compared to n.

Lemma 2.1. *Let $0 < \eta_n \downarrow 0$ be such that $\eta_n \log n \uparrow \infty$ as $n \uparrow \infty$. Set*

$$\rho_n^{\mathsf{crit}} \equiv c^{\mathsf{crit}} \tfrac{\log n}{n \log 2}, \quad c^{\mathsf{crit}} \equiv c_n^{\mathsf{crit}} = 2\left[1 + \tfrac{\log 2}{2\eta_n \log n}\left(1 + \tfrac{3 \log n}{n \log 2}\right)\right]. \tag{2.6}$$

There exists $\Omega^{\mathsf{CRIT}} \subset \Omega$ with $\mathbb{P}(\Omega^{\mathsf{CRIT}}) = 1$ such that on Ω^{CRIT}, for all but a finite number of indices n, for all $\rho \geq \rho_n^{\mathsf{crit}}$, the connected components in (2.4) satisfy

$$2 \leq \max_{1 \leq l \leq L} |C_{n,l}(\rho)| \leq n\eta_n \ll n. \tag{2.7}$$

Remark. Although c^{crit} in (2.6) is very likely not optimal, we clearly must have $c^{\mathsf{crit}} > 1$.

Proof of Lemma 2.1. The proof relies on a lemma that we first state and prove. Define

$$\widetilde{\Omega}_n(m) = \left\{\omega \in \Omega \,\middle|\, \max_{1\le l\le L} |C_{n,l}(\rho)| < m\right\}, \quad m = 2, 3, \dots \tag{2.8}$$

In what follows $\rho \equiv \rho_n > 0$ and $m \equiv m_n > 1$ are, respectively, positive and integer valued sequences. To keep the notation simple we do not make this dependence explicit.

Lemma 2.2. *If* $\rho \ge \rho_n^+(m) \equiv \frac{1}{m}\big(1 + \frac{(m+2)\log n + \log m!}{n \log 2}\big)$ *then* $\mathbb{P}\big(\liminf_{n\to\infty} \widetilde{\Omega}_n(m)\big) = 1$.

Proof of Lemma 2.2. Call $(\chi_n(x), x \in \mathcal{V}_n)$, $\chi_n(x) \equiv \mathbb{1}_{\{w_n(x)\ge r_n(\rho)\}}$, the occupancy variables. These are i.i.d. Bernoulli r.v.'s with $\mathbb{P}(\chi_n(x) = 1) = 1 - \mathbb{P}(\chi_n(x) = 0) = 2^{-\rho n}$. Set $\mathbb{P}\big(\widetilde{\Omega}_n^c(m)\big) = 1 - \mathbb{P}\big(\widetilde{\Omega}_n(m)\big) = \mathbb{P}\big(\exists_{\mathcal{C}_n\subseteq V_n(\rho):|\mathcal{C}_n|=m} G(\mathcal{C}_n) \text{ is connected}\big)$. The number of connected components of size m is at most $m!n^m 2^n$, and by independence, if $|\mathcal{C}_n| = m$ then $\mathbb{P}\big(G(\mathcal{C}_n) \text{ is connected}\big) = \mathbb{P}\big(\prod_{x\in\mathcal{C}_n} \chi_n(x) = 1\big) = (2^{-\rho n})^m$. Thus, for $\rho \ge \rho_n^+(m)$, $\mathbb{P}\big(\widetilde{\Omega}_n^c(m)\big) \le m!n^m 2^{(1-m\rho)n} \le m!n^m 2^{(1-m\rho_n^+(m))n} \le n^{-2}$, so that $\sum_{n\ge1} \mathbb{P}\big(\widetilde{\Omega}_n^c(m)\big) < \infty$. The lemma now follows from the first Borel-Cantelli Lemma. □

Lemma 2.1 follows from Lemma 2.2 by taking $m = n\eta_n$, observing that for this choice $\rho_n^{\mathsf{crit}} > \rho_n^+(m)$, and setting $\Omega^{\mathsf{CRIT}} \equiv \liminf_{n\to\infty} \widetilde{\Omega}_n(m)$. □

2.2. **Truncation and related quantities.** Throughout the rest of the paper we assume that $c_\star > 2$ in (1.12) so that $c_\star \ge c_n^{\mathsf{crit}}$ for large enough n. According to (2.4)-(2.5), for $\rho = \rho_n^\star$ as in (2.3), $\mathcal{V}_n$ be decomposed in a unique way into

$$\mathcal{V}_n = N_n^\star \cup I_n^\star \cup \big(\cup_{l=1}^{L^\star} C_{n,l}^\star\big), \quad L^\star \equiv L(\rho_n^\star), \tag{2.9}$$

where $N_n^\star \equiv N_n(\rho_n^\star)$, $I_n^\star \equiv I_n(\rho_n^\star)$, and $C_{n,l}^\star \equiv C_{n,l}(\rho_n^\star)$, $1 \le l \le L^\star$. By construction $H_n(x) = 0$ if and only if $x \in N_n^\star$ (see (1.2) and (1.12)). Furthermore $\mathcal{V}_n^\star$ in (1.13) becomes

$$\mathcal{V}_n^\star = \cup_{l=1}^{L^\star} C_{n,l}^\star. \tag{2.10}$$

Lemma 2.3. *Assume that* $c_\star > 2$. *There exists* $\Omega^\star \subset \Omega$ *with* $\mathbb{P}(\Omega^\star) = 1$ *such that on* $\Omega^\star$, *for all but a finite number of indices* n *the following holds:*

$$2 \le \left|C_{n,l}^\star\right| \le \{\rho_n^\star[1 - 2c_\star^{-1}(1 + \mathcal{O}(\log n/n))]\}^{-1}, \quad 1 \le l \le L^\star. \tag{2.11}$$

Furthermore,

$$|I_n^\star| = 2^n n^{-c_\star}(1 - n^{-(c_\star-1)})(1 + \mathcal{O}(n^{-2(c_\star-1)}) + o(n^{-c_\star})), \tag{2.12}$$

$$|V_n(\rho_n^\star)| = \left|\mathcal{V}_n \setminus N_n^\star\right| = 2^n n^{-c_\star}(1 - n^{-c_\star})(1 + o(n^{-c_\star})), \tag{2.13}$$

$$\textstyle\sum_{l=1}^{L^\star} \left|C_{n,l}^\star\right| = |V_n(\rho_n^\star) \setminus I_n^\star| = 2^n n^{-2c_\star+1}(1 + \mathcal{O}(n^{-(c_\star-1)})), \tag{2.14}$$

and, denoting by $\partial A \equiv \{y \in \mathcal{V}_n \setminus A : \mathrm{dist}(y, A) = 1\}$ *the boundary of the set* $A \subset \mathcal{V}_n$,

$$n\left|C_{n,l}^\star\right|(1 - \mathcal{O}(\tfrac{1}{\log n})) \le \left|\partial C_{n,l}^\star\right| \le n\left|C_{n,l}^\star\right|, \tag{2.15}$$

$$|\partial C_{n,l}^\star \cap \partial x| \ge n(1 - \mathcal{O}(\tfrac{1}{\log n})) \ \textit{for all} \ x \in C_{n,l}^\star, \tag{2.16}$$

$$n\left|C_{n,l}^\star\right|(1 - \mathcal{O}(\tfrac{1}{\log n})) \le \textstyle\sum_{x\in C_{n,l}^\star} \sum_{y\in\partial C_{n,l}^\star:\{x,y\}\in\mathcal{E}_n} 1 \le n\left|C_{n,l}^\star\right|. \tag{2.17}$$

Proof. The claim of (2.11) follows from Lemma 2.2. Next,

$$|V_n(\rho_n^\star)| = \textstyle\sum_{x\in\mathcal{V}_n} \chi_n(x), \ \ |I_n^\star| = \sum_{x\in\mathcal{V}_n} \chi_n(x) \prod_{y\in\mathcal{V}_n:(x,y)\in\mathcal{E}_n} (1 - \chi_n(y)), \tag{2.18}$$

and $\sum_{l=1}^{L^\star} \left|C_{n,l}^\star\right| = \sum_{x\in\mathcal{V}_n} \chi_n(x)[1 - \prod_{y\in\mathcal{V}_n:(x,y)\in\mathcal{E}_n}(1 - \chi_n(y))]$, where, as in the proof of Lemma 2.2, $\chi_n(x) \equiv \mathbb{1}_{\{w_n(x)\ge r_n(\rho_n^\star)\}}$ are i.i.d. Bernoulli r.v. with $\mathbb{P}(\chi_n(x) = 1) = n^{-c_\star}$. From these expressions (2.13), (2.12), and (2.14) are easily obtained. Turning to (2.17)

note that the sum therein can be written as $\sum_{x\in C^\star_{n,l}}(n-d_n(x))$ where $d_n(x)$ denotes the connectivity of the vertex x in the graph $G(V_n(\rho^\star_n))$. This, the bound $1\le d_n(x)\le \big|C^\star_{n,l}\big|$, and (2.11) yield the desired result. To prove the lower bound of (2.15) reason that each vertex x in $C^\star_{n,l}$ has at least $n-d_n(x)$ nearest neighbors vertices in $\partial C^\star_{n,l}$, and that no two vertices in $C^\star_{n,l}$ can have more than one common nearest neighbor vertex in $\partial C^\star_{n,l}$. Hence $\big|\partial C^\star_{n,l}\big|\ge\sum_{x\in C^\star_{n,l}}(n-d_n(x)-(\big|C^\star_{n,l}\big|-1))\ge\sum_{x\in C^\star_{n,l}}(n-2\big|C^\star_{n,l}\big|)$ and the lower bound in (2.15) follows from (2.11). Eq. (2.16) is proved in the same way since $|\partial C^\star_{n,l}\cap\partial x|=n-d(x)$ for $x\in C^\star_{n,l}$. Finally, the upper bound of (2.15) is immediate. □

We conclude this section with two elementary lemmata that are repeatedly needed. The first expresses the function $r_n(\rho)$ defined through (2.2).

Lemma 2.4. *For all $\rho>0$, possibly depending on n, such that $\rho n\uparrow\infty$ as $n\uparrow\infty$,*

$$r_n(\rho)=\exp\big\{n\beta\beta_c(\rho)-(\beta/2\beta_c(\rho))\big[\log(\beta_c^2(\rho)n/2)+\log 4\pi\big]+o(\beta/\beta_c(\rho))\big\}\ . \tag{2.19}$$

In particular, for $\rho^\star_n$ as in (2.3) and $c_\star>2$,

$$r_n\big(\rho^\star_n\big)=\exp\Big\{\beta\Big(\sqrt{2c_\star n\log n}-\sqrt{\tfrac{n}{\log n}}\big(\tfrac{\log\log n}{2\sqrt{2c_\star}}+\mathcal{O}(1)\big)\Big)\Big\}. \tag{2.20}$$

Proof. Denote by Φ and ϕ the standard Gaussian distribution function and density, respectively. Setting $b_n=2^{\rho n}$ and $\overline{B}_n=\log r_n(\rho)/\beta\sqrt{n}$, (2.2) becomes $b_n\big(1-\Phi(\overline{B}_n)\big)=1$. It is shown in [25] (see paragraph below (2.20)) that $(\overline{B}_n-B_n)B_n=o(1)$ where B_n is defined through $b_n\frac{\phi(B_n)}{B_n}=1$. Eq. (2.19) then readily follows from the well known fact that (see [18], p. 374) $B_n=(2\log b_n)^{\frac12}-\frac12(\log\log b_n+\log 4\pi)/(2\log b_n)^{\frac12}+\mathcal{O}(1/\log b_n)$. □

Lemma 2.5. *There exists a subset $\Omega_0\subseteq\Omega$ with $\mathbb{P}\big(\Omega_0\big)=1$ such that on Ω_0, for all but a finite number of indices n the following holds: for all $1\le l\le L^\star$*

$$e^{-\beta\min\{\max(H_n(y),H_n(x))\,|\,\{x,y\}\in G(C^\star_{n,l})\}}\le e^{\beta n\sqrt{\log 2}(1+2\log n/n\log 2)}, \tag{2.21}$$

$$e^{-\beta\min\{H_n(x)\,|\,x\in C^\star_{n,l}\}}\le e^{\beta n\sqrt{2\log 2}(1+2\log n/n)}. \tag{2.22}$$

Proof. Set $\rho(1)\equiv 1+2\log n/n\log 2$, $\Omega_n(1)\equiv\{\omega\in\Omega\mid\max_{x\in\mathcal{V}_n}w_n(x)\le r_n(\rho(1))\}$, and $\Omega_\infty(1)\equiv\liminf_{n\to\infty}\Omega_n(1)$. Further set $\rho(2)\equiv\frac12(1+3\log n/n\log 2)$, $\Omega_n(2)\equiv\{\omega\in\Omega\mid\max_{(x,y)\in\mathcal{E}_n}\min(w_n(x),w_n(y))\le r_n(\rho(2))\}$, and $\Omega_\infty(2)\equiv\liminf_{n\to\infty}\Omega_n(2)$. By independence and (2.2), $\mathbb{P}(\Omega_n^c(1))=2^n2^{-n}n^{-2}=n^{-2}$ which is summable, hence $\mathbb{P}(\Omega_\infty(1))=1$. Next, $\mathbb{P}(\Omega_n^c(2))\le n2^{n-1}2^{-n}(r_n(\rho(2)))^2\le n^{-2}$ which is also summable, and so $\mathbb{P}(\Omega_\infty(2))=1$. Taking $\Omega_0\equiv\Omega_\infty(1)\cap\Omega_\infty(2)$ and using (2.19) to bound $r_n(\rho(1))$ and $r_n(\rho(2))$ yields the claim of the lemma. □

3. Front end and back end clock processes, and proofs of the theorems of Section 1.

In this section we formally define the *front end* and *back end* clock processes, and show how they relate to the clock processes S_n and K_n. These relations are then used to decompose the proofs of Theorem 1.1 and Theorem 1.2 into five main steps. Let $C^\star_{n,l}$, $1\le l\le L^\star$, be the collection of connected components defined through (2.9) and set

$$\mathcal{V}_n^\circ\equiv\mathcal{V}_n\setminus\big(\cup_{1\le l\le L^\star}C^\star_{n,l}\big)\,. \tag{3.1}$$

3.1. **Front end clock process.** We call *front end clock process* the process defined through

$$\widetilde{S}_n^\circ(k^\circ) = \sum_{i=0}^{k^\circ-1} \lambda_n^{-1}(J_n^\circ(i)) e_{n,i}^\circ, \quad k^\circ \in \mathbb{N}, \tag{3.2}$$

where $(e_{n,i}^\circ\,, n \in \mathbb{N}, i \in \mathbb{N})$ are independent mean one exponential random variables and where, introducing the times of consecutive visits of J_n to $\mathcal{V}_n^\circ$,

$$\begin{aligned} T_{n,0}^\circ &= \inf\{i \geq 0 \mid J_n(i) \in \mathcal{V}_n^\circ\}\,, & (3.3)\\ T_{n,j+1}^\circ &= \inf\{i > T_{n,j}^\circ \mid J_n(i) \in \mathcal{V}_n^\circ\},\ j = 0, 1, 2, \ldots, & (3.4) \end{aligned}$$

$(J_n^\circ(i), i \in \mathbb{N})$ is the reversible Markov chain on $\mathcal{V}_n^\circ$ obtained by setting $J_n^\circ(i) \equiv J_n(T_{n,i}^\circ)$. Note that J_n° has transition matrix elements

$$p_n^\circ(x,y) = P_x\left(J_n(T_{n,1}^\circ) = y\right),\ x, y \in \mathcal{V}_n^\circ, \tag{3.5}$$

and invariant measure

$$\pi_n^\circ(x) = \frac{\pi_n(x)}{\sum_{x' \in \mathcal{V}_n^\circ} \pi_n(x')}, \quad x \in \mathcal{V}_n^\circ, \tag{3.6}$$

where π_n denotes the invariant measure of J_n (see (6.7) for its expression). We call J_n° the front chain and denote by $(\Omega^{J^\circ}, \mathcal{F}^{J^\circ}, P^\circ)$ its probability space. The associated graph, $G^\circ(\mathcal{V}_n^\circ)$, is described in (6.10).

3.2. **Back end clock process.** The description of this process involves three time sequences. The first two are the intertwined sequences of consecutive hitting times of $\mathcal{V}_n \setminus \mathcal{V}_n^\circ$ and their ensuing exit times. Namely, set

$$\overline{T}_{n,0} = 0, \quad \overline{T}'_{n,0} = \begin{cases} \inf\{i > 0 \mid J_n(i) \notin \mathcal{V}_n^\circ\}, & \text{if } J_n(0) \in \mathcal{V}_n^\circ, \\ 0, & \text{if } J_n(0) \notin \mathcal{V}_n^\circ, \end{cases} \tag{3.7}$$

and, for $j = 0, 1, 2, \ldots,$

$$\begin{aligned} \overline{T}_{n,j+1} &= \inf\{i > \overline{T}'_{n,j} \mid J_n(i) \in \mathcal{V}_n^\circ\}\,, & (3.8)\\ \overline{T}'_{n,j+1} &= \inf\{i > \overline{T}_{n,j+1} \mid J_n(i) \notin \mathcal{V}_n^\circ\}. & (3.9) \end{aligned}$$

Clearly, $0 = \overline{T}_{n,0} \leq \overline{T}'_{n,0} < \overline{T}_{n,1} \leq \overline{T}'_{n,1} < \cdots < \overline{T}_{n,j} \leq \overline{T}'_{n,j} < \ldots$. Clearly also, to each j there corresponds an i such that $T_{n,i-1}^\circ < \overline{T}'_{n,j} = T_{n,i-1}^\circ + 1 < T_{n,i}^\circ$. Merging $\left(T_{n,i}^\circ\right)_{i\geq 0}$ and $\left(\overline{T}'_{n,j}\right)_{j\geq 0}$ into a single sequence, $\left(T_{n,j}^\dagger\right)_{j\geq 0}$, and arranging its elements in increasing order of magnitude,

$$0 \leq T_{n,0}^\dagger < T_{n,1}^\dagger < \cdots < T_{n,j}^\dagger < \ldots. \tag{3.10}$$

we define the *back end clock process* through

$$\widetilde{S}_n^\dagger(k^\dagger) = \sum_{i=0}^{k^\dagger-1} \Lambda_n^\dagger(J_n^\dagger(i)), \quad k^\dagger \in \mathbb{N}, \tag{3.11}$$

where $(J_n^\dagger(i), i \in \mathbb{N})$ is the chain on $\mathcal{V}_n$ obtained by setting $J_n^\dagger(i) \equiv J_n(T_{n,i}^\dagger)$, and where

$$\Lambda_n^\dagger(J_n^\dagger(i)) = \begin{cases} \overline{T}_{n,j+1} - \overline{T}'_{n,j}, & \text{if } J_n^\dagger(i) \notin \mathcal{V}_n^\circ \text{ and } \sum_{k=0}^{i} \mathbb{1}_{\{J_n^\dagger(k) \notin \mathcal{V}_n^\circ\}} = j, \\ 0, & \text{if } J_n^\dagger(i) \in \mathcal{V}_n^\circ. \end{cases} \tag{3.12}$$

Clearly, $J_n^\dagger$ is Markovian with one-step transitions probabilities, $p_n^\dagger(x,y)$, as follows: when it is at $x \in \mathcal{V}_n^\circ$, $J_n^\dagger$ chooses its next step according to the transition probabilities of J_n,

$$p_n^\dagger(x,y) = p_n(x,y), \quad x \in \mathcal{V}_n^\circ, y \in \mathcal{V}_n, \tag{3.13}$$

and when it enters $\cup_{1\le l\le L^\star} C_{n,l}^\star$, say at a vertex of $C_{n,l}^\star$, it exits in just one step through one of the boundary points $\partial C_{n,l}^\star$; that is, for all $x \in C_{n,l}^\star$, $y \in \partial C_{n,l}^\star$, and $1 \le l \le L^\star$,

$$p_n^\dagger(x,y) = P_x(J(T_{n,l}^\star) = y), \tag{3.14}$$

where $T_{n,l}^\star = \inf\{i > 0 \mid J_n(i) \in \partial C_{n,l}^\star\}$. Clearly also, the increments $\Lambda_n^\dagger(J_n^\dagger(i))$ of the clock at the times of the visits of $J_n^\dagger(i)$ to $\cup_{1\le l\le L^\star} C_{n,l}^\star$ are the sojourn times of J_n in the sets $C_{n,l}^\star$ being visited. In other words, $\Lambda_n^\dagger(J_n^\dagger(i))$ is equal in distribution to some $T_{n,l}^\star$.

Summarizing our definitions, FECP (3.2) records the total time spent by the proces X_n in $\mathcal{V}_n^\circ$ along the first k° steps of J_n° whereas BECP (3.11) records the total time spent by the chain J_n in $\cup_{1\le l\le L^\star} C_{n,l}^\star$ along the first $k^\dagger$ steps of $J_n^\dagger$. The chains $J_n^\dagger$ and J_n° differ in that $J_n^\dagger$ does visit the sets $C_{n,l}^\star$, and steps out of these sets right after stepping in, while J_n° straddles over the $C_{n,l}^\star$'s, never entering them. Technically, this makes the two chains very different objects. In particular, J_n° is reversible but $J_n^\dagger$ isn't.

3.3. **Rewriting the clock process.** Our aim is to express the processes K_n and S_n defined in (1.16) and (1.9), respectively, using FECP and BECP. We first deal with K_n. For a_n as in (1.16) let $k_n^\dagger(t)$ be defined through

$$k_n^\dagger(t) = \min\left\{k \ge 1 \,\middle|\, \textstyle\sum_{i=0}^{k-1} \mathbb{1}_{\{J_n^\dagger(i)\in\mathcal{V}_n^\circ\}} = \lfloor a_n t\rfloor\right\}, \quad t \ge 0, \tag{3.15}$$

and, taking $k^\dagger = k_n^\dagger(t)$ in (3.11), set

$$S_n^\dagger(t) = b_n^{-1}\widetilde{S}_n^\dagger(k_n^\dagger(t)), \quad t \ge 0. \tag{3.16}$$

Then $K_n(t)$ can be writen as

$$K_n(t) = \lfloor a_n t\rfloor + b_n S_n^\dagger(t), \quad t \ge 0. \tag{3.17}$$

To see this write $K_n(t) = \sum_{i=0}^{K_n(t)-1} \mathbb{1}_{\{J_n(i)\in\mathcal{V}_n^\circ\}} + \sum_{i=0}^{K_n(t)-1} \mathbb{1}_{\{J_n(i)\notin\mathcal{V}_n^\circ\}}$ and note that

$$\textstyle\sum_{i=0}^{K_n(t)-1} \mathbb{1}_{\{J_n(i)\notin\mathcal{V}_n^\circ\}} = \sum_{i=0}^{k_n^\dagger(t)-1} \Lambda_n^\dagger(J_n^\dagger(i)) = b_n S_n^\dagger(t), \tag{3.18}$$

$$\textstyle\sum_{i=0}^{K_n(t)-1} \mathbb{1}_{\{J_n(i)\in\mathcal{V}_n^\circ\}} = \sum_{i=0}^{k_n^\dagger(t)-1} \mathbb{1}_{\{J_n^\dagger(i)\in\mathcal{V}_n^\circ\}} = \lfloor a_n t\rfloor \equiv k_n^\circ(t), \tag{3.19}$$

where we introduced the notation $k_n^\circ(t)$ for later convenience. In words, when J_n takes $K_n(t)$ steps, $J_n^\dagger$ takes $k_n^\dagger(t)$ steps, of which $k_n^\circ(t)$ are visits of $J_n^\dagger$ to $\mathcal{V}_n^\circ$.

To deal with the clock process S_n we likewise split the sum in (1.9) in two terms according to whether $J_n(i) \in \mathcal{V}_n^\circ$ or $J_n(i) \notin \mathcal{V}_n^\circ$. From the above definitions and those of $J_n^\dagger$ and J_n° we have that on the one hand, writing $\stackrel{d}{=}$ for equality in distribution,

$$\begin{aligned}
\textstyle\sum_{i=0}^{K_n(t)-1} \lambda_n^{-1}(J_n(i)) e_{n,i} \mathbb{1}_{\{J_n(i)\in\mathcal{V}_n^\circ\}} &\stackrel{d}{=} \textstyle\sum_{j=0}^{k_n^\dagger(t)-1} \lambda_n^{-1}(J_n^\dagger(j)) e_{n,j}^\dagger \mathbb{1}_{\{J_n^\dagger(i)\in\mathcal{V}_n^\circ\}} &(3.20)\\
&\stackrel{d}{=} \textstyle\sum_{j=0}^{k_n^\circ(t)-1} \lambda_n^{-1}(J_n^\circ(j)) e_{n,j}^\circ \mathbb{1}_{\{J_n^\circ(i)\in\mathcal{V}_n^\circ\}} &(3.21)\\
&= \widetilde{S}_n^\circ(\lfloor a_n t\rfloor), &(3.22)
\end{aligned}$$

where $(e^{\dagger}_{n,j})$ and $(e^{\circ}_{n,j})$ are families of independent mean one exponential random variables, and $\widetilde{S}^{\circ}_n$ is the front end clock process (3.2). On the other hand,

$$\sum_{i=0}^{K_n(t)} \lambda_n^{-1}(J_n(i))e_{n,i}\mathbb{1}_{\{J_n(i)\notin\mathcal{V}_n^{\circ}\}} \tag{3.23}$$

$$= \sum_{j=0}^{k_n^{\dagger}(t)-1}\Big(\sum_{i=0}^{\Lambda_n^{\dagger}(J_n^{\dagger}(j))-1}\lambda_n^{-1}(J_n(\overline{T}'_{n,j}+i))e_{n,i}\Big)\,\mathbb{1}_{\{J_n^{\dagger}(j)\notin\mathcal{V}_n^{\circ}\}} \tag{3.24}$$

$$\equiv \sum_{j=0}^{k_n^{\dagger}(t)-1}\widehat{\Lambda}_n^{\dagger}(J_n^{\dagger}(j)) \tag{3.25}$$

where the last line defines $\widehat{\Lambda}_n^{\dagger}(J_n^{\dagger}(j))$. If we now set, for $t\geq 0$,

$$S_n^{\circ}(t) \equiv c_n^{-1}\widetilde{S}_n^{\circ}(\lfloor a_n t\rfloor), \tag{3.26}$$

$$\widehat{S}_n(t) \equiv c_n^{-1}\sum_{j=0}^{k_n^{\dagger}(t)-1}\widehat{\Lambda}_n^{\dagger}(J_n^{\dagger}(j)), \tag{3.27}$$

the rescaled clock process (1.9) can be rewritten as

$$S_n(t)\stackrel{d}{=}S_n^{\circ}(t)+\widehat{S}_n(t). \tag{3.28}$$

Here the rescaled front end clock process, $S_n^{\circ}(t)$, records the time spent *by the process* X_n during its visits to the set $\mathcal{V}_n^{\circ}$, while the remainder term, $\widehat{S}_n(t)$, records the time spent in its complement. The back end clock process $b_nS_n^{\dagger}(t)$ is the time needed to actually be able to observe a transition *of the chain* J_n from one vertex of $\mathcal{V}_n^{\circ}$ to the next.

3.4. **Proofs of the theorems of Section 1.** The proofs of the theorems of Section 1.2 rely on five theorems stated below. Each of them controls one of the processes $k_n^{\dagger}(t)$, $S_n^{\circ}(t)$, $\widehat{S}_n(t)$, and $S_n^{\dagger}(t)$ above, respectively below, the critical line $\beta=2\beta_c(\varepsilon/2)$, $0<\varepsilon<1$. As in Section 1.2 the initial distribution of J_n is the uniform distribution on $\mathcal{V}_n^{\circ}$. By (6.5), this is nothing but the invariant measure, π_n°, of J_n°. Hence J_n° and $J_n^{\dagger}$ also start in π_n°.

The first theorem shows that $k_n^{\dagger}(t)$ behaves like $k_n^{\circ}(t)=\lfloor a_nt\rfloor$ for large n.

Theorem 3.1. *Assume that $c_\star>2$. For all $0<t<\infty$, any constant $c_\circ>0$, and any sequence $a_n>0$ we have that on $\Omega^\star$, for all but a finite number of indices n,*

$$P_{\pi_n^{\circ}}\big(1\leq k_n^{\dagger}(t)/k_n^{\circ}(t)\leq 1+n^{-c_\circ}\big)\geq 1-n^{-2(c_\star-1)+c_\circ}(1+\mathcal{O}(n^{-(c_\star-1)})). \tag{3.29}$$

The next two theorems are the building blocks of the proof of Theorem 1.1. The first establishes convergence of the front end clock process, S_n°. The second implies, in particular, that the contribution of $\widehat{S}_n$ to (3.28) vanishes as n diverges.

Theorem 3.2 (Front end clock process)**.** *Assume that $c_\star>3$. Let the sequences a_n and c_n be as in Theorem 1.1. Then, for all $0<\varepsilon<1$ and $\beta>\beta_c(\varepsilon)$, $\mathbb{P}$-almost surely,*

$$S_n^{\circ}\Rightarrow_{J_1}S_\infty^{\circ}, \tag{3.30}$$

where S_∞° is a subordinator with zero drift and Lévy measure $\nu^{\circ}=\nu$ defined in (1.21).

Theorem 3.3 (Remainder)**.** *Assume that $c_\star>2$. Let the sequences a_n and c_n be as in Theorem 1.1. Then, for all $0<\varepsilon<1$ and $\beta>\beta_c(\varepsilon)$, $\mathbb{P}$-almost surely,*

$$\limsup_{n\to\infty}\mathcal{P}_{\pi_n^{\circ}}\big(\rho_\infty\big(S_n(\cdot),S_n^{\circ}(\cdot)\big)>\epsilon\big)=0,\quad\forall\epsilon>0, \tag{3.31}$$

where ρ_∞ is Skorohod metric on $D([0,\infty))$.

We now turn to the back end clock process. The next result parallels Theorem 3.2.

Theorem 3.4 (Back end clock process above the critical line)**.** *Assume that $c_\star > 3$. Let the sequence a_n and b_n be as in Theorem 1.1 and Theorem 1.2, (i), respectively. Then, for all $0 < \varepsilon < 1$ and $\beta > 2\beta_c(\varepsilon/2)$, $\mathbb{P}$-almost surely,*

$$S_n^\dagger \Rightarrow_{J_1} S_\infty^\dagger, \tag{3.32}$$

where $S_\infty^\dagger$ is a stable subordinator with zero drift and Lévy measure $\nu^\dagger$ defined in (1.23).

Finally, the last result covers the high temperature domain.

Theorem 3.5 (Back end clock process below the critical line)**.** *Assume that $c_\star > 3$. Let the sequence a_n and b_n be as in Theorem 1.1 and Theorem 1.2, (ii), respectively. Then, for all $0 < \varepsilon < 1$, all $0 < \beta < 2\beta_c(\varepsilon/2)$, and all $0 < T < \infty$, $\mathbb{P}$-almost surely,*

$$\mathcal{P}_{\pi_n^\circ}\Big(\lim_{n\to\infty} \sup_{t\in[0,T]} \big|b_n^{-1} S_n^\dagger(t) - t\big| = 0\Big) = 1. \tag{3.33}$$

Assuming these theorems we may prove Theorem 1.1 and Theorem 1.2. The proof of Theorem 1.3 also uses properties of the chain J_n° that are taken from Section 6.

Proof of Theorem 1.1. In view of (3.28) Theorem 1.1 is an immediate consequence of Theorem 3.2 and Theorem 3.3 □

Proof of Theorem 1.2. Recall the expression (3.17) of K_n and notice that $a_n/b_n \downarrow 0$ both under the assumptions on a_n and b_n of Theorem 3.5 and Theorem 3.4 (use (2.19) to check this in the latter case). Thus the first assertion of Theorem 1.2 is a an immediate consequence of Theorem 3.4 while the second assertion follows from Theorem 3.5. □

Proof of Theorem 1.3. The proof follows classical arguments. Let $A_n^\rho(t,s)$ be the event $A_n^\rho(t,s) \equiv \{\mathcal{C}_n(t,s) \geq 1-\rho\}$. Denote respectively by $\mathcal{R}_n$ and $\mathcal{R}_n^\circ$ the ranges of the processes $c_n^{-1}\widetilde{S}_n$ and $c_n^{-1}\widetilde{S}_n^\circ$. Clearly, for all $\rho \in (0,1)$, $A_n^\rho(t,s) \supseteq \{\mathcal{R}_n \cap (t,t+s) = \emptyset\}$. Now $\lim_{n\to\infty} \mathcal{P}_{\pi_n^\circ}(\mathcal{R}_n \cap (t,t+s) = \emptyset) = \lim_{n\to\infty} \mathcal{P}_{\pi_n^\circ}(\mathcal{R}_n^\circ \cap (t,t+s) = \emptyset)$ as follows from Theorem 3.3, and by Theorem 3.2, proceeding as in the proof of Theorem 1.6 in [26], $\lim_{n\to\infty} \mathcal{P}_{\pi_n^\circ}(\mathcal{R}_n^\circ \cap (t,t+s) = \emptyset) = \mathcal{P}(\{S_\infty^\circ(u), u > 0\} \cap (t,t+s) = \emptyset)$, but by the arcsine law for stable subordinators (see e.g. Theorem 1.8 of [26]) the last probability is equal to the right hand side of (1.25).

It thus remains to show that $\lim_{n\to\infty} \mathcal{P}_{\pi_n^\circ}\big(A_n^\rho(t,s) \cap \{\mathcal{R}_n \cap (t,t+s) \neq \emptyset\}\big) = 0$. Invoking as before Theorem 3.3, we can substitute $\mathcal{R}_n^\circ$ for $\mathcal{R}_n$ in the probability. Consider the set $\mathcal{T}_n(\epsilon) \equiv \{x \in I_n^\star \mid w_n(x) \geq \epsilon c_n\}$, $\epsilon > 0$. By Theorem 3.2, if $\mathcal{R}_n^\circ \cap (t,t+s) \neq \emptyset$ then with a probability that tends to one as $n \uparrow \infty$ and $\epsilon \downarrow 0$ there exists $u_- \leq u_+$ such that on the one hand $c_n^{-1}\widetilde{S}_n^\circ(\lfloor a_n u_- \rfloor - 1) < t < c_n^{-1}\widetilde{S}_n^\circ(\lfloor a_n u_- \rfloor)$ while $c_n^{-1}\widetilde{S}_n^\circ(\lfloor a_n u_+ \rfloor) < t+s < c_n^{-1}\widetilde{S}_n^\circ(\lfloor a_n u_+ \rfloor + 1)$ on the other, and these two increments correspond to visits to vertices z_- and z_+ in $\mathcal{T}_n(\epsilon)$ (that is to say, with probability one, the points t and $t+s$ lie in constancy intervals of the process, and such intervals are produced, asymptotically, by visits to $\mathcal{T}_n(\epsilon)$). Let us now argue that, firstly, that starting from a given vertex $z_- \in \mathcal{T}_n(\epsilon)$, the chain J_n° quickly moves at a distance greater than $n\rho/2$ from it, and secondly, that it does not visit any vertex in $\{z \in \mathcal{T}_n(\epsilon) \mid \mathrm{dist}(z_-, z) \leq n\rho/2\}$ in the ensuing $\lfloor Ca_n \rfloor$ steps, for any $0 < C < \infty$, $0 < \rho < 1$, and small $\epsilon > 0$. For this we use three results of Section 6. By Proposition 6.1, the chain J_n° started in z_- reaches stationarity in $\ell_n^\circ \sim n^3$ steps, and by Proposition 6.3, $\pi_n^\circ(\{z \in \mathcal{V}_n^\circ \mid \mathrm{dist}(z_-, z) > n\rho/2\}) \geq 1 - \exp\{-n\mathcal{I}(\rho)\}$, where $\mathcal{I}(\rho) > 0$ if $0 < \rho < 1$. This proves the first claim. The second claim is an immediate consequence of Proposition 6.4. The proof of Theorem 1.3 is done. □

The rest of this paper is organized as follows. In Section 4 we focus on the increments of the process $\widehat{S}_n$ and prove an upper bound on their tail distribution. A similar analysis is carried out in Section 5 for the increments of the back end clock process $\widetilde{S}_n^\dagger$; an explicit expression is also obtained for the distribution of the sojourn times of J_n in sets $C_{n,l}^\star$ of size 2. The properties of J_n° (invariant measure, mixing time through spectral gap, mean local times) are studied in Section 6, where it is shown that J_n° has several of the attributes of the symmetric random walk. Using these preparations, the proofs of Theorem 3.1 and Theorem 3.3 are carried out in Section 7. Those of Theorem 3.2, Theorem 3.4, and Theorem 3.5 are carried out in Section 8, Section 9, and Section 10, respectively. Finally, Appendix 11 gathers needed results on the speed of convergence to Perron projector for nonnegative and primitive matrices (Subsection 11.1), and for irreducible and periodic matrices with period 2 (Subsection 11.2).

4. Distribution of the increments of the process $\widehat{S}_n$.

In this section we focus on the increments of the process $\widehat{S}_n$, that is to say, on the quantities defined through (3.24)-(3.25) by

$$\widehat{\Lambda}_n^\dagger(J_n^\dagger(j)) \equiv \sum_{i=0}^{\Lambda_n^\dagger(J_n^\dagger(j))-1} \lambda_n^{-1}(J_n(\overline{T}'_{n,j}+i))e_{n,i} \tag{4.1}$$

if $J_n^\dagger(j) \in \cup_{1\le l\le L^\star} C_{n,l}^\star$, and $\widehat{\Lambda}_n^\dagger(J_n^\dagger(j)) = 0$ otherwise. These are the sojourn times of the process X_n in the sets $C_{n,l}^\star$ (we may think of them as "effective holding times" in those sets). As expected, these times have exponential tails. For $1 \le l \le L^\star$, set

$$\bar{\varrho}_{n,l}(0) = e^{-\beta \min\left\{H_n(x)\,|\,x\in C_{n,l}^\star\right\}}, \tag{4.2}$$

$$\bar{\varrho}_{n,l}(1) = e^{-\beta \min\left\{H_n(x)\,|\,x\in C_{n,l}^\star\right\}+\beta \max\left\{H_n(y)\,|\,y\in C_{n,l}^\star\right\}}, \tag{4.3}$$

and

$$\bar{\theta}_{n,l}^\star \equiv 3\beta n^2 |C_{n,l}^\star|^3 \bar{\varrho}_{n,l}(1). \tag{4.4}$$

Proposition 4.1. *Let $\Omega^\star$ and Ω_0 be as in Lemma 2.3 and Lemma 2.5, respectively. On $\Omega^\star \cap \Omega_0$, for all but a finite number of indices n, the following holds for all $1 \le l \le L^\star$. For all $t \ge \bar{\theta}_{n,l}^\star$ and all x in $C_{n,l}^\star$,*

$$\mathcal{P}\left(\widehat{\Lambda}_n^\dagger(J_n^\dagger(j)) > t \mid J_n^\dagger(j) = x\right) \le e^{-t/\tau_{n,l}^\star}(1+\bar{\epsilon}_n^\star), \tag{4.5}$$

where $0 \le \bar{\epsilon}_n^\star \le \frac{\bar{\theta}_{n,l}^\star |C_{n,l}^\star|}{\bar{\varrho}_{n,l}(0)}(1+o(1))$ and

$$\tfrac{1}{|C_{n,l}^\star|}\bar{\varrho}_{n,l}(0) \le \tau_{n,l}^\star \le (1+\mathcal{O}(\tfrac{1}{\log n}))\bar{\varrho}_{n,l}(0). \tag{4.6}$$

Moreover $(\tau_{n,l}^\star, 1 \le l \le L^\star)$ are independent random variables on $(\Omega, \mathcal{F}, \mathbb{P})$.

The next corollary is a crucial ingredient of the proof of Theorem 3.3.

Corollary 4.2. *Assume that $a_n \le 2^n$. On $\Omega_0 \cap \Omega^\star$, for all but a finite number of indices n,*

$$\mathcal{P}_{\pi_n^\circ}\left(\exists_{0\le j\le k_n^\dagger(t)-1}\exists_{1\le l\le L^\star}\widehat{\Lambda}_n^\dagger(J_n^\dagger(j))\mathbb{1}_{\{J_n^\dagger(j)\in C_{n,l}^\star\}} > n\bar{\varrho}_{n,l}(0)\right) \le te^{-n}+n^{-2(c_\star-1)+c_\circ} \tag{4.7}$$

where $c_\circ > 0$ is a constant that can be chosen arbitrarily small.

Let us sketch the proof of Proposition 4.1. Suppose that $J_n^\dagger(j) \in C_{n,l}^\star$. Then $\widehat{\Lambda}_n^\dagger(J_n^\dagger(j))$ is equal in distribution to the absorption time of the restriction of the process X_n to $C_{n,l}^\star$, killed on the boundary $\partial C_{n,l}^\star$ and started in $J_n^\dagger(j)$. It is well known that its law is governed by the spectral characteristics of the associated infinitesimal generator (more specifically by the corresponding Dirichlet eigenvalues) and is, on suitable time scales, approximated to very good precision by an exponential law. This section is thus organized as follows. In Subsection 4.1 we introduce the absorbing processes of interest, study their low lying Dirichlet eigenvalues in Subsection 4.2 and, in Subsection 4.3, deduce from this the distribution of the absorption times, using a result from Appendix 11. The proof of Corollary 4.2 is done in Subsection 4.4.

4.1. **The 'starred' absorbing processes.** Let $C_{n,l}^\star$, $1 \le l \le L^\star$, be the collection of connected components defined through (2.9). To each component $C_{n,l}^\star$ we associate an absorbing Markov process $X_{n,l}^\star$ with state space $C_{n,l}^\star \cup \Delta$, where the absorbing point, Δ, represents the boundary $\partial C_{n,l}^\star$; its infinitesimal generator $\overline{\mathcal{L}}_{n,l}^\star = \big(\bar\lambda_{n,l}^\star(x,y)\big)$ has entries $\bar\lambda_{n,l}^\star : \{C_{n,l}^\star \cup \Delta\} \times \{C_{n,l}^\star \cup \Delta\} \to \mathbb{R}$, given by

$$\bar\lambda_{n,l}^\star(x,y) = \begin{cases} \lambda_n(x,y) & \text{if} \quad (x,y) \in G(C_{n,l}^\star), \\ \sum_{y' \notin C_{n,l}^\star} \lambda_n(x,y') & \text{if} \quad x \in C_{n,l}^\star, y = \Delta, \\ -\sum_{y' \in \mathcal{V}_n} \lambda_n(x,y') & \text{if} \quad x = y \in C_{n,l}^\star, \\ 0 & \text{else.} \end{cases} \tag{4.8}$$

Thus $X_{n,l}^\star$ can be viewed as the restriction of X_n to $C_{n,l}^\star$, killed on the boundary $\partial C_{n,l}^\star$. We also call $\mathcal{L}_{n,l}^\star = \big(\lambda_{n,l}^\star(x,y)\big)$ the sub-Markovian restriction of $\overline{\mathcal{L}}_{n,l}^\star$ to $C_{n,l}^\star$, namely $\lambda_{n,l} : C_{n,l}^\star \times C_{n,l}^\star \to \mathbb{R}$,

$$\lambda_{n,l}^\star(x,y) = \begin{cases} \lambda_n(x,y) & \text{if} \quad (x,y) \in G(C_{n,l}^\star) \\ -\sum_{y' \in \mathcal{V}_n} \lambda_n(x,y') & \text{if} \quad x = y \in C_{n,l}^\star. \end{cases} \tag{4.9}$$

With this notation $\widehat{\Lambda}_n^\dagger(J_n^\dagger(j))$ in (4.1) is nothing but the absorption time

$$\Lambda_{n,l}^\star \equiv \inf\{t > 0 \mid X_{n,l}^\star(t) = \Delta\} \tag{4.10}$$

of the process $X_{n,l}^\star$ started in $X_{n,l}^\star(0) = J_n^\dagger(j)$.

4.2. **Spectrum of $-\mathcal{L}_{n,l}^\star$ and other properties.** It follows from the properties of the sets $C_{n,l}^\star$ that $\{\Delta\}$ is the unique recurrence class of $\overline{\mathcal{L}}_{n,l}^\star$ and that $\mathcal{L}_{n,l}^\star$ is irreducible (in fact primitive) and non periodic. Also note that $\mathcal{L}_{n,l}^\star$ is reversible with respect to the restriction of Gibbs measure (see (1.3)) to $C_{n,l}^\star$,

$$\mu_{\beta,n,l}^\star(x) = \frac{G_{\beta,n}(x)}{\sum_{x' \in C_{n,l}^\star} G_{\beta,n}(x')}, \quad x \in C_{n,l}^\star. \tag{4.11}$$

Let I_d denote the identity matrix in $\mathbb{R}^d$ and set $N \equiv N_{n,l} = |C_{n,l}^\star|$. In view of (1.4) the matrix $\overline{R}_{n,l} \equiv \overline{\mathcal{L}}_{n,l}^\star + I_{N+1}$ is stochastic (its entries are in $[0,1]$ and its rows sum up to one) and $\{\Delta\}$ is its unique recurrence class. From this and Perron's theorem we deduce (see (11.1)) that the eigenvalues of $-\mathcal{L}_{n,l}^\star$ satisfy

$$0 < \varsigma_{n,l}(0) < \varsigma_{n,l}(1) \le \varsigma_{n,l}(2) \cdots \le \varsigma_{n,l}(N-1). \tag{4.12}$$

The next two Lemmata state bounds on the eigenvalues $\varsigma_{n,l}(0)$ and $\varsigma_{n,l}(1)$. Let $\bar\varrho_{n,l}(0)$ and $\bar\varrho_{n,l}(1)$ be given by (4.2) and (4.3) respectively.

Lemma 4.3. *On $\Omega^\star$, for all but a finite number of indices n, $\frac{1-\mathcal{O}(1/\log n)}{\bar\varrho_{n,l}(0)} \le \varsigma_{n,l}(0) \le \frac{|C^\star_{n,l}|}{\bar\varrho_{n,l}(0)}$.*

Lemma 4.4. $\varsigma_{n,l}(1) \ge 1/\left(n|C^\star_{n,l}|^2\bar\varrho_{n,l}(1)\right)$.

Proof of Lemma 4.3. Rather than considering the matrix $-\mathcal{L}^\star_{n,l}$, it is convenient to consider

$$R_{n,l} \equiv \mathcal{L}^\star_{n,l} + I_N \tag{4.13}$$

that is to say, the nonnegative and substochastic matrix $R_{n,l} = (r_{n,l}(x,y))$ with entries

$$r^\star_{n,l}(x,y) = \begin{cases} \lambda_n(x,y), & \text{if} \quad (x,y) \in G(C^\star_{n,l}), \\ 1 - \sum_{y' \in \mathcal{V}_n} \lambda_n(x,y'), & \text{if} \quad x = y \in C^\star_{n,l} \end{cases} \tag{4.14}$$

and eigenvalues $1 > \vartheta_{n,l}(0) > \vartheta_{n,l}(1) \ge \cdots \ge \vartheta_{n,l}(N-1) > -1$,

$$\vartheta_{n,l}(i) \equiv 1 - \varsigma_{n,l}(i). \tag{4.15}$$

To bound $\vartheta_{n,l}(0)$ from below note that since $R_{n,l}$ is reversible with respect to $\mu^\star_{\beta,n,l}$, it is similar to the matrix with entries $r_{n,l}(x,y)\sqrt{\mu^\star_{n,l}(x)/\mu^\star_{n,l}(y)}$, and the latter being symmetric, the minimax characterization of its largest eigenvalue (see e.g. [29], p. 176) gives

$$\vartheta_{n,l}(0) = \sup_{u \ne 0} \frac{\sum_{\{x,y\} \in C^\star_{n,l} \times C^\star_{n,l}} u(x)u(y)\mu^\star_{n,l}(x) r_{n,l}(x,y)}{\sum_{x \in C^\star_{n,l}} u^2(x)\mu^\star_{n,l}(x)}, \tag{4.16}$$

where supremum is taken over all non zero functions real valued function u on $C^\star_{n,l}$. Thus $\vartheta_{n,l}(0)$ is bounded below by the ratio in the r.h.s. of (4.16) evaluated at constant functions,

$$\vartheta_{n,l}(0) \ge \textstyle\sum_{\{x,y\} \in C^\star_{n,l} \times C^\star_{n,l}} \mu^\star_{n,l}(x) r_{n,l}(x,y) = 1 - A/B \tag{4.17}$$

where, by (4.11), (4.14), and the definition (1.4) of $\lambda_n(x,y)$,

$$A \equiv \textstyle\sum_{x \in C^\star_{n,l}} \sum_{y \in \partial C^\star_{n,l} : \{x,y\} \in \mathcal{E}_n} e^{-\beta \max(H_n(y), H_n(x))}, \tag{4.18}$$

$$B \equiv n \textstyle\sum_{x \in C^\star_{n,l}} e^{-\beta H_n(x)}. \tag{4.19}$$

Now by construction (see (2.9) in Subsection 2.2) the maximum in (4.18) is equal to zero. Hence $A = \sum_{x \in C^\star_{n,l}} \sum_{y \in \partial C^\star_{n,l} : \{x,y\} \in \mathcal{E}_n} 1 \le n|C^\star_{n,l}|$. This and the bound

$$n \le B e^{\beta \min\left\{H_n(x) \,|\, x \in C^\star_{n,l}\right\}} \le n|C^\star_{n,l}| \tag{4.20}$$

yields $\vartheta_{n,l}(0) = 1 - \varsigma(0) \ge 1 - \frac{|C^\star_{n,l}|}{\bar\varrho_{n,l}(0)}$, which is the desired lower bound.

We now turn to the upper bound. Denote by $R^\star_{n,l}$ the stochastic matrix $R^\star_{n,l} \equiv R_{n,l} + D_{n,l}$ where $D_{n,l} = (d_{n,l}(x,y))$ is the diagonal matrix with entries $d_{n,l} : C^\star_{n,l} \times C^\star_{n,l} \to [0,1]$,

$$d_{n,l}(x,y) = \begin{cases} \sum_{y' \notin C^\star_{n,l}} \lambda_n(x,y'), & \text{if} \quad x = y, x \in C^\star_{n,l}, \\ 0, & \text{else.} \end{cases} \tag{4.21}$$

For $0 \le k \le N-1$ denote by $\vartheta^\star_{n,l}(k)$ and $\vartheta^d_{n,l}(k)$ the eigenvalues of $R^\star_{n,l}$ and $D_{n,l}$, respectively, arranged in decreasing order of magnitude. Then, since $R_{n,l} = R^\star_{n,l} - D_{n,l}$, by Weyl's theorem on eigenvalues (see e.g. [29] p. 181),

$$\vartheta_{n,l}(k) \le \vartheta^\star_{n,l}(k) - \vartheta^d_{n,l}(N-1), \quad 0 \le k \le N-1. \tag{4.22}$$

Here $\vartheta^\star_{n,l}(0) = 1$ and $\vartheta^d_{n,l}(N-1) = \min_{x\in C^\star_{n,l}} \sum_{y'\in\partial C^\star_{n,l}:\{x,y'\}\in\mathcal{E}_n} \lambda_n(x,y')$. Thus,

$$\vartheta_{n,l}(0) \le 1 - \min_{x\in C^\star_{n,l}} \sum_{y'\in\partial C^\star_{n,l}:\{x,y'\}\in\mathcal{E}_n} \lambda_n(x,y') = 1 - \frac{|\partial C^\star_{n,l}\cap \partial x|}{n\bar\varrho_{n,l}(0)}, \tag{4.23}$$

where we used that $0 = H_n(y') \ge H_n(x)$ for all $\{x,y'\}\in\mathcal{E}_n$ with $x\in C^\star_{n,l}$ and $y'\in\partial C^\star_{n,l}$. Inserting the bound (2.16) of Lemma 2.3 we obtain that, on $\Omega^\star$, for all but a finite number of indices n, $\vartheta_{n,l}(0) = 1-\varsigma(0) \le 1-(1-\mathcal{O}(\frac{1}{\log n}))\frac{1}{\bar\varrho_{n,l}(0)}$ for all $x\in C^\star_{n,l}$. The proof of Lemma 4.3 is done. □

Proof of Lemma 4.4. Applying the bound (4.22) with $k=1$ gives $\vartheta_{n,l}(1) \le \vartheta^\star_{n,l}(1)$ since, clearly, $\vartheta^d_{n,l}(N-1)\ge 0$. $R^\star_{n,l}$ being a stochastic matrix, questions on $\vartheta^\star_{n,l}(1)$ then reduce to classical questions on the spectral gap $\tau^{-1}_{n,l} \equiv 1-\vartheta^\star_{n,l}(1)$. It is in particular well known that

$$\tau_{n,l} = \sup\left\{\frac{\mathcal{E}(u,u)}{Var(u)} \,:\, u \text{ is nonconstant}\right\}, \tag{4.24}$$

where, denoting by u a real valued function on $C^\star_{n,l}$,

$$Var(u) = \tfrac12 \sum_{x,y} \left(u(x)-u(y)\right)^2 \mu^\star_{n,l}(x)\mu^\star_{n,l}(y), \tag{4.25}$$

$$\mathcal{E}(u,u) = \tfrac12 \sum_{x,y} \left(u(x)-u(y)\right)^2 \mu^\star_{n,l}(x) r_{n,l}(x,y). \tag{4.26}$$

It is also well known that based on (4.24), one may derive bounds on $\tau_{n,l}$ in terms of "canonical paths" (see e.g. [32]). The bound stated below is taken from [21] (see Proposition 1' p. 38). For each pair of distinct vertices $x,y\in C^\star_{n,l}$, choose a path $\gamma_{x,y}$ going from x to y in the graph $G(C^\star_{n,l})$. Paths may have repeated vertices but a given edge appears at most once in a given path. Let $\Gamma_{n,l}$ denote a collection of paths (one for each pair $\{x,y\}$). Then

$$\tau_{n,l} \le \max_e \rho_n^{-1}(e) \sum_{\gamma_{x,y}\ni e} |\gamma_{x,y}|\, \mu^\star_{n,l}(x)\mu^\star_{n,l}(y), \tag{4.27}$$

where the max is over all edges $e=\{x',y'\}$ of $G(C^\star_{n,l})$, $\rho_n(e) \equiv \mu^\star_{n,l}(x') r_{n,l}(x',y')$, and the summation is over all paths in $\Gamma_{n,l}$ that pass through e. The quality of this bound usually depends on the choice of $\Gamma_{n,l}$. Here however we simply bound the length of the longest path by the total number of edges in the graph, that is $|\gamma_{x,y}| \le |C^\star_{n,l}|$. Eq. (4.27) then immediately gives $\tau_{n,l} \le \left|C^\star_{n,l}\right| \max_e \rho_n^{-1}(e)$, and so

$$\vartheta_{n,l}(1) \le 1 - \left|C^\star_{n,l}\right|^{-1} \min_{\{x,y\}\in G(C^\star_{n,l})} \mu^\star_{n,l}(x) r_{n,l}(x,y). \tag{4.28}$$

Proceeding as in (4.17) to evaluate the r.h.s. above, we obtain

$$\vartheta_{n,l}(1) \le 1 - \left|C^\star_{n,l}\right|^{-1} \min_{\{x,y\}\in G(C^\star_{n,l})} B^{-1} e^{-\beta\max(H_n(y),H_n(x))}, \tag{4.29}$$

where B is defined in (4.19). Finally, plugging in the upper bound of (4.20) we arrive at $\vartheta_{n,l}(1) = 1-\varsigma_{n,l}(1) \le 1-(n|C^\star_{n,l}|^2\bar\varrho_{n,l}(1))^{-1}$. The proof of Lemma 4.4 is complete. □

We close this subsection with bounds that will be needed in several places.

Lemma 4.5. *For $r_n(\rho^\star_n)$ as in (2.20),*

$$e^{\beta n\sqrt{2\log 2}(1+2\log n/n)} \ge \bar\varrho_{n,l}(0) \ge \varrho_{n,l}(1)\frac{\bar\varrho_{n,l}(0)}{\bar\varrho_{n,l}(1)} = e^{-\beta\max\left\{H_n(x)\,|\,x\in C^\star_{n,l}\right\}} \ge r_n\left(\rho^\star_n\right),$$

where the first inequality holds on Ω_0, for all but a finite number of indices n.

Proof. The first inequality is (2.22). The second is immediate. The third follows by construction from the truncation (see (1.2), (1.12), and (2.9)) and relation 2.3. □

4.3. **Absorption times: proof of Proposition 4.1.**

Proof of Proposition 4.1. Assume that (4.5) holds for integer values of t in $[\bar\theta^\star_{n,l}-1,\infty)$. Then, writing $t-1<\lfloor t\rfloor\le t$ where $\lfloor t\rfloor$ is the largest integer smaller than or equal to t, we have, for all $t\ge\bar\theta^\star_{n,l}$,

$$\mathcal{P}_x\left(\Lambda^\star_{n,l}>t\right)\le\mathcal{P}_x\left(\Lambda^\star_{n,l}>\lfloor t\rfloor\right)\le e^{-\lfloor t\rfloor/\tau^\star_{n,l}}(1+\bar\epsilon^\star_n)\le e^{-t/\tau^\star_{n,l}}e^{1/\tau^\star_{n,l}}(1+\bar\epsilon^\star_n),\quad(4.30)$$

where by (4.6), $e^{1/\tau^\star_{n,l}}(1+\bar\epsilon^\star_n)\le\frac{\bar\theta^\star_{n,l}|C^\star_{n,l}|}{\bar\varrho_{n,l}(0)}(1+o(1))$.

It thus suffices to prove (4.5) for integer t's in $[\bar\theta^\star_{n,l}-1,\infty)$. Assume from now on that $t=m>1$ is an integer and set $A\equiv e^{\mathcal{L}^\star_{n,l}}$. Writing $A^m_{n,l}=(a^{(m)}_{n,l}(x,y))$, it follows from (4.8)-(4.10) and the semigroup property that

$$\mathcal{P}_x\left(\Lambda^\star_{n,l}>m\right)=\textstyle\sum_{y\in C^\star_{n,l}}(\delta_x,e^{m\mathcal{L}^\star_{n,l}}\delta_y)=\sum_{y\in C^\star_{n,l}}a^{(m)}_{n,l}(x,y),\quad(4.31)$$

where $(\cdot,\cdot)$ denotes the inner product in $\mathbb{R}^N$, $N\equiv|C^\star_{n,l}|$, and δ_x is the vector with components $\delta_x(x')=1$ if $x'=x$ and zero otherwise. Clearly, $A_{n,l}$ is a nonnegative and primitive matrix, and is reversible with respect to $\mu^\star_{\beta,n,l}$. We can therefore use Lemma 11.1 to evaluate the right hand side of (4.31). For this note that $A_{n,l}$ has eigenvalues $\exp\{-\varsigma_{n,l}(i)\}$, $0\le i\le N-1$, where the $\varsigma_{n,l}(i)$'s obey (4.12), and denote by $u^\star_{n,l}$ and $v^\star_{n,l}$ the left and right Perron eigenvectors of $A_{n,l}$ (see (11.1)-(11.2)),

$$u^\star_{n,l}A_{n,l}=e^{-\varsigma_{n,l}(0)}u^\star_{n,l},\quad u^\star_{n,l}>0,\quad(4.32)$$

$$A_{n,l}v^\star_{n,l}=e^{-\varsigma_{n,l}(0)}v^\star_{n,l},\quad v^\star_{n,l}>0,\quad(4.33)$$

normalized to make

$$\textstyle\sum_{x\in C^\star_{n,l}}u^\star_{n,l}(x)=1,\quad\sum_{x\in C^\star_{n,l}}u^\star_{n,l}(x)v^\star_{n,l}(x)=1.\quad(4.34)$$

Then, by (11.5) of Lemma 11.1,

$$\mathcal{P}_x\left(\Lambda^\star_{n,l}>m\right)=v^\star_{n,l}(x)e^{-m\varsigma_{n,l}(0)}\left\{1+\mathcal{R}_ne^{-m(\varsigma_{n,l}(1)-\varsigma_{n,l}(0))}\right\},\quad(4.35)$$

where

$$|\mathcal{R}_n|\le\left(\min_{x\in C^\star_{n,l}}\left\{v^\star_{n,l}(x)(\mu^\star_{n,l}(x))^{1/2}\right\}\right)^{-1}.\quad(4.36)$$

In order to control $|\mathcal{R}_n|$ we need a lower bound on $v^\star_{n,l}(x)$.

Lemma 4.6. *For all $x\in C^\star_{n,l}$, $v^\star_{n,l}(x)\ge(1/\bar\varrho_{n,l}(0))^{|C^\star_{n,l}|}$.*

Proof of Lemma 4.6. Write $A_{n,l}=e^{\mathcal{L}^\star_{n,l}}=e^{-I_N+(\mathcal{L}^\star_{n,l}+I_N)}=e^{-1}e^{R_{n,l}}$ and recall from (4.13) that $R_{n,l}\equiv\mathcal{L}^\star_{n,l}+I_N$ is a nonnegative and primitive matrix. By the spectral decomposition the left and right Perron eigenvectors of $A_{n,l}$ and $R_{n,l}$ coincide, and thus

$$u^\star_{n,l}R_{n,l}=(1-\varsigma_{n,l}(0))u^\star_{n,l},\quad u^\star_{n,l}>0,\quad(4.37)$$

$$R_{n,l}v^\star_{n,l}=(1-\varsigma_{n,l}(0))v^\star_{n,l},\quad v^\star_{n,l}>0,\quad(4.38)$$

where $u^\star_{n,l}$ and $v^\star_{n,l}$ obey (4.34). This implies in particular that

$$\textstyle\sum_{x\in C^\star_{n,l}}v^\star_{n,l}(x)\ge1.\quad(4.39)$$

Indeed, by (4.34), $u^\star_{n,l}(x) \le 1$ for all $x \in C^\star_{n,l}$ and so $1 = \sum_{x\in C^\star_{n,l}} u^\star_{n,l}(x)v^\star_{n,l}(x) \le \sum_{x\in C^\star_{n,l}} v^\star_{n,l}(x)$. Equipped with (4.38) and (4.39) we prove the lemma by contradiction. Assume that $v^\star_{n,l}(x) < \varepsilon_0 \equiv (1/\bar\varrho_{n,l}(0))^{|C^\star_{n,l}|}$ for some $x \in C^\star_{n,l}$. Then, by (4.38),

$$\sum_{x\in C^\star_{n,l}} r_{n,l}(y,x)v^\star_{n,l}(x) = (1-\varsigma_{n,l}(0))v^\star_{n,l}(y) \le \varepsilon_0, \tag{4.40}$$

and since $\min\{r_{n,l}(y,x) \,|\, x,y \in C^\star_{n,l}\} \ge \frac{r_n(\rho^\star_n)}{n\bar\varrho_{n,l}(0)}$,

$$v^\star_{n,l}(x) \le \varepsilon_1 \equiv \varepsilon_0 \tfrac{n\bar\varrho_{n,l}(0)}{r_n(\rho^\star_n)} \text{ for each } x \in C^\star_{n,l} \cap \partial y. \tag{4.41}$$

Repeating this reasoning for each $x \in C^\star_{n,l} \cap \partial y$, then for each $x \in C^\star_{n,l} \cap \partial_2 y$, and so on and so forth, we arrive at

$$\max\{v^\star_{n,l}(x) \,|\, x \in C^\star_{n,l}\} \le \varepsilon_0 \big(\tfrac{n\bar\varrho_{n,l}(0)}{r_n(\rho^\star_n)}\big)^{|C^\star_{n,l}|} < \tfrac{n}{r_n(\rho^\star_n)}. \tag{4.42}$$

In view of (2.20) this implies that $\sum_{x\in C^\star_{n,l}} v^\star_{n,l}(x) \ll 1$, contradicting (4.39). Therefore $v^\star_{n,l}(x) \ge \varepsilon_0$ for all $x \in C^\star_{n,l}$. □

Clearly, $\mu^\star_{n,l}(x) \ge 1/(\bar\varrho_{n,l}(1)|C^\star_{n,l}|)$ for all $x \in C^\star_{n,l}$. Using this bound and Lemma 4.6 in (4.36), it follows from (2.11) and the first and last inequality of Lemma 4.5 that on $\Omega_0 \cap \Omega^\star$, for all but a finite number of indices n,

$$|\mathcal{R}_n| \le \sqrt{n/r_n(\rho^\star_n)}\, e^{\beta n(|C^\star_{n,l}|+1/2)\sqrt{2\log 2}(1+2\log n/n)}. \tag{4.43}$$

Next, by Lemma 4.3, Lemma 4.4, and Lemma 4.5 we have that on $\Omega^\star$, for all but a finite number of indices n,

$$\varsigma_{n,l}(1) - \varsigma_{n,l}(0) \ge (1-o(1))/n|C^\star_{n,l}|^2\bar\varrho_{n,l}(1) \equiv 3\beta n|C^\star_{n,l}|(1-o(1))/\bar\theta^\star_{n,l}, \tag{4.44}$$

where $\bar\theta^\star_{n,l}$ is defined in (4.4). Inserting (4.43) and (4.44) in (4.35) we obtain that on $\Omega_0 \cap \Omega^\star$, for all $m \ge \bar\theta^\star_{n,l} - 1$ and all but a finite number of indices n.

$$\mathcal{P}_x\left(\Lambda^\star_{n,l} > m\right) = v^\star_{n,l}(x)e^{-m\varsigma_{n,l}(0)}(1+\bar\epsilon'_n), \tag{4.45}$$

where $\bar\epsilon'_n = \mathcal{O}(\exp(-\beta n))$. $\mathcal{P}_x\left(\Lambda^\star_{n,l} > \bar\theta^\star_{n,l}\right) \le 1$ yields $(1+\bar\epsilon'_n)v^\star_{n,l}(x) \le e^{\bar\theta^\star_{n,l}\varsigma_{n,l}(0)}$. Plugging in the upper bound on $\varsigma_{n,l}(0)$ of Lemma 4.3 and inserting the result in (4.45) we obtain that

$$\mathcal{P}_x\left(\Lambda^\star_{n,l} > m\right) = e^{-m\varsigma_{n,l}(0)}(1+\bar\epsilon^\star_n), \tag{4.46}$$

where $0 \le \bar\epsilon^\star_n \le \frac{\bar\theta^\star_{n,l}|C^\star_{n,l}|}{\bar\varrho_{n,l}(0)}(1+o(1))$. Taking $\tau^\star_{n,l} = 1/\varsigma_{n,l}(0)$ in (4.45) now yields the statement of (4.5) for all integer values of t in $[\bar\theta^\star_{n,l}-1,\infty)$. The claimed independance of the $\tau^\star_{n,l}$'s is a direct consequence of the definition of the $\varsigma_{n,l}(0)$'s. The proof of Proposition 4.1 is complete. □

4.4. **Proof of Corollary 4.2.** First note that the choice $t = n\bar\varrho_{n,l}(0)$ in Proposition 4.1 is admissible since by (4.2), (4.3), (4.4), (2.11), Lemma 4.5, and (2.20) of Lemma 2.4, on $\Omega^\star$, for all n large enough,

$$\tfrac{n\bar\varrho_{n,l}(0)}{\theta^\star_{n,l}} = [3\beta n|C^\star_{n,l}|^3]^{-1}\tfrac{\bar\varrho_{n,l}(0)}{\bar\varrho_{n,l}(1)} \ge [3\beta n|C^\star_{n,l}|^3]^{-1}\varrho_{n,l}(1) \ge n^{-4}r_n(\rho^\star_n) \gg 1. \tag{4.47}$$

Moreover by (4.6), for this choice, $t/\tau^\star_{n,l} = n\bar\varrho_{n,l}(0)/\tau^\star_{n,l} \ge n|C^\star_{n,l}| \ge 2n$. Hence Proposition 4.1 applies so that on $\Omega_0 \cap \Omega^\star$, for all but a finite number of indices n,

$$\mathcal{P}\big(\widehat\Lambda^\dagger_n(J^\dagger_n(j)) > n\bar\varrho_{n,l}(0) \mid J^\dagger_n(j) = x\big) \le e^{-2n}(1+o(1)) \tag{4.48}$$

for all $1 \le l \le L^\star$ and all $x \in C^\star_{n,l}$. Let $\mathcal{A}$ be the event in the left hand side of (4.7). By Theorem 3.1, $\mathcal{P}_{\pi^\circ_n}(\mathcal{A}) \le \mathcal{P}_{\pi^\circ_n}(\mathcal{A}, \{k^\dagger_n(t) \le \lfloor a_n t \rfloor (1+n^{-c_\circ})\}) + 2n^{-2(c_\star-1)+c_\circ}$, and by (4.48), on $\Omega_0 \cap \Omega^\star$, $\mathcal{P}_{\pi^\circ_n}(\mathcal{A}, \{k^\dagger_n(t) \le \lfloor a_n t \rfloor (1+n^{-c_\circ})\}) \le 2\lfloor a_n t \rfloor (1+n^{-c_\circ}) e^{-2n}$ for all but a finite number of indices n. Since $a_n < 2^n$, (4.7) follows.

5. Distribution of the increments of the back end clock process $\widetilde{S}^\dagger_n$.

This section parallels Section 4, focusing this time on the increments, $\Lambda^\dagger_n$, of the process $\widetilde{S}^\dagger_n$. Just as the $\widehat{\Lambda}^\dagger_n$'s are the sojourn times of the process X_n in the sets $C^\star_{n,l}$, the $\Lambda^\dagger_n$'s are the sojourn times of the chain J_n in those sets. Indeed, by (3.12),

$$\Lambda^\dagger_n(J^\dagger_n(j)) \equiv \inf\{i > j \mid J_n(i) \in \partial C^\star_{n,l}\} - j \tag{5.1}$$

if $J^\dagger_n(j) \in C^\star_{n,l}$ for some $1 \le l \le L^\star$, and $\widehat{\Lambda}^\dagger_n(J^\dagger_n(j)) = 0$ otherwise. For $1 \le l \le L^\star$, set

$$\varrho_{n,l}(0) = e^{-\beta \min\left\{\max(H_n(y), H_n(x)) \,|\, \{x,y\} \in G(C^\star_{n,l})\right\}}, \tag{5.2}$$

$$\varrho_{n,l}(1) = e^{-\beta \max\left\{H_n(x) \,|\, x \in C^\star_{n,l}\right\}}, \tag{5.3}$$

and

$$\theta^\star_{n,l} = 2\beta n |C^\star_{n,l}|^5 \left(\varrho_{n,l}(0)/\varrho_{n,l}(1)\right). \tag{5.4}$$

Proposition 5.1. *(i) For each $1 \le l \le L^\star$ such that $|C^\star_{n,l}| = 2$ we have, for all $i > 0$ and all x in $C^\star_{n,l}$,*

$$\mathcal{P}\left(\Lambda^\dagger_n(J^\dagger_n(j)) > i \mid J^\dagger_n(j) = x\right) = \left(1 - \tfrac{1}{1+\varrho_{n,l}(0)/(n-1)}\right)^i. \tag{5.5}$$

(ii) Furthermore, on $\Omega^\star \cap \Omega_0$ (for $\Omega^\star$ and Ω_0 as in Lemma 2.3 and Lemma 2.5, respectively), for all but a finite number of indices n, the following holds for all $1 \le l \le L^\star$: for all $i \ge \theta^\star_{n,l}$ and all x in $C^\star_{n,l}$,

$$\mathcal{P}\left(\Lambda^\dagger_n(J^\dagger_n(j)) > i \mid J^\dagger_n(j) = x\right) \le e^{-i(n/\varrho_{n,l}(0)|C^\star_{n,l}|)(1-o(1))}(1+\hat{\epsilon}_n), \tag{5.6}$$

where $0 \le \hat{\epsilon}_n \le \frac{\theta^\star_{n,l} n}{\varrho_{n,l}(0)|C^\star_{n,l}|}(1+o(1))$.

This section is organized as Section 4. In Subsection 5.1 we introduce an absorbing Markov chain $J^\star_{n,l}$, defined as the restriction of J_n to $C^\star_{n,l}$, killed on the boundary $\partial C^\star_{n,l}$. In Subsection 5.2 we establish properties of the spectrum of an associated sub-Markovian transition matrix, $Q_{n,l}$, needed in Subsection 5.3 to prove bounds on the time to absorption.

5.1. The 'starred' absorbing chains.

Let $C^\star_{n,l}$, $1 \le l \le L^\star$, be the collection of connected components defined through (2.9). To each component $C^\star_{n,l}$ we associate an absorbing Markov chain $J^\star_{n,l}$ with state space $C^\star_{n,l} \cup \Delta$, where the absorbing point Δ represents the boundary $\partial C^\star_{n,l}$; its transition matrix $P^\star_{n,l} = \left(p^\star_{n,l}(x,y)\right)$ has entries $p^\star_{n,l} : \{C^\star_{n,l} \cup \Delta\} \times \{C^\star_{n,l} \cup \Delta\} \to [0,1]$,

$$p^\star_{n,l}(x,y) = \begin{cases} p_n(x,y) & \text{if} \quad (x,y) \in G(C^\star_{n,l}), \\ 1 - \sum_{y' \in C^\star_{n,l}} p_n(x,y'), & \text{if} \quad x \in C^\star_{n,l}, y = \Delta, \\ 1, & \text{if} \quad x = y = \Delta, \\ 0, & \text{else.} \end{cases} \tag{5.7}$$

Thus $J^\star_{n,l}$ can be viewed as the restriction of J_n to $C^\star_{n,l}$, killed on the boundary $\partial C^\star_{n,l}$. We also call $Q_{n,l} = (q_{n,l}(x,y))$ the sub-Markovian restriction of $P^\star_{n,l}$ to $C^\star_{n,l}$, namely $q_{n,l} : C^\star_{n,l} \times C^\star_{n,l} \to [0,1]$,

$$q_{n,l}(x,y) = \begin{cases} p_n(x,y) & \text{if} \quad (x,y) \in G(C^\star_{n,l}), \\ 0, & \text{else.} \end{cases} \tag{5.8}$$

Then, $\Lambda^\dagger_n(J^\dagger_n(j))$ in (5.1) is equal in distribution to the absorption time

$$T^\star_{n,l} = \inf\{i \in \mathbb{N} \mid J^\star_{n,l}(i) = \Delta\} \tag{5.9}$$

of the process $J^\star_{n,l}$ started in $J^\star_{n,l}(0) = J^\dagger_n(j)$.

5.2. **Spectrum of $Q_{n,l}$.** Clearly, $Q_{n,l}$ is reversible with respect to the measure $\pi^\star_{n,l}$,

$$\pi^\star_{n,l}(x) = \frac{\pi_n(x)}{\sum_{x' \in C^\star_{n,l}} \pi_n(x')}, \quad x \in C^\star_{n,l}, \tag{5.10}$$

where π_n denotes the invariant measure of J_n (see (6.7)). From the properties of the sets $C^\star_{n,l}$ it follows that $\{\Delta\}$ is the unique recurrence class of $P^\star_{n,l}$ and that $Q_{n,l}$ is irreducible and periodic with period 2 (indeed $G(C^\star_{n,l}) \subseteq \mathcal{Q}_n$ is a connected sub-graph of the bipartite graph $\mathcal{Q}_n$). Therefore, by (11.10) and (11.11), the eigenvalues of $Q_{n,l}$ satisfy

$$1 > \vartheta_{n,l}(0) > \vartheta_{n,l}(1) \geq \vartheta_{n,l}(2) \cdots \geq \vartheta_{n,l}(N-1) > -1, \tag{5.11}$$

where $N \equiv N_{n,l} = |C^\star_{n,l}|$. The next two lemmata give bounds on the first and second eigenvalues, respectively, in terms of the quantities (5.2) and (5.3).

Lemma 5.2. *If $|C^\star_{n,l}| = 2$ then*

$$\vartheta_{n,l}(0) = 1 - [1 + \varrho_{n,l}(0)/(n-1)]^{-1}. \tag{5.12}$$

Furthermore on $\Omega^\star$, for all but a finite number of indices n, for all $1 \leq l \leq L^\star$,

$$1 - \frac{1 + \mathcal{O}(1/\log n)}{1 + 2\varrho_{n,l}(0)/(n|C^\star_{n,l}|)} \leq \vartheta_{n,l}(0) \leq 1 - \frac{1 - \mathcal{O}(1/\log n)}{1 + \varrho_{n,l}(0)|C^\star_{n,l}|/n}. \tag{5.13}$$

Remark. Note that the bounds of Lemma 5.2 are rather sharp, i.e. the prefactors of $\varrho_{n,l}(0)$ only differ through the terms $|C^\star_{n,l}|^{\pm 1}$, and $2 \leq |C^\star_{n,l}| \leq cnst.n/\log n$.

Lemma 5.3. *On $\Omega^\star$, for all but a finite number of indices n, for all $1 \leq l \leq L^\star$,*

$$\vartheta_{n,l}(1) \leq 1 - \varrho_{n,l}(1) \left[n|C^\star_{n,l}|^3(1 + \varrho_{n,l}(0)|C^\star_{n,l}|/n)\right]^{-1}. \tag{5.14}$$

Proof of Lemma 5.2 and Lemma 5.3. Bounds on the first two eigenvalues of $Q_{n,l}$ are established in exactly the same way as in the case of the matrix $R_{n,l}$ defined in (4.13) (see the proof of Lemma 4.3 and Lemma 4.4). The case $|C^\star_{n,l}| = 2$ is easily worked out by hand. We skip the details. □

Corollary 5.4. *On $\Omega^\star$, for all but a finite number of indices n,*

$$\vartheta_{n,l}(1)/\vartheta_{n,l}(0) \leq \exp\left\{-|C^\star_{n,l}|^{-4}(\varrho_{n,l}(1)/\varrho_{n,l}(0))(1 + o(1))\right\}. \tag{5.15}$$

Proof. By Lemmata 5.2 and 5.3, $\vartheta_{n,l}(1)/\vartheta_{n,l}(0) \leq |C^\star_{n,l}|^{-4}(\varrho_{n,l}(1)/\varrho_{n,l}(0))(1+o(1))$, and since $1 - x \leq e^{-x}$ for $0 < x < 1$, (5.15) follows. □

Finally, the lemma below collects useful bounds.

Lemma 5.5. *For each $1 \le l \le L^\star$ and for $r_n\big(\rho_n^\star\big)$ as in (2.20),*

$$e^{\beta n\sqrt{\log 2}(1+o(1))} \ge \varrho_{n,l}(0) \ge \varrho_{n,l}(1) \ge r_n\left(\rho_n^\star\right), \tag{5.16}$$

where the first inequality holds on Ω_0, for all but a finite number of indices n.

Proof. The first inequality is (2.21). The remaining ones are immediate. □

5.3. **Absorption time: proof of Proposition 5.1.** This section and the next draw on Appendix 11 where needed results on irreducible and periodic matrices are collected. We begin with a little notation. Recall that **1** denotes the vertex of $\mathcal{V}_n$ whose coordinates are identically 1. Write $\mathcal{V}_n \equiv \mathcal{V}_n^- \cup \mathcal{V}_n^+$ where $\mathcal{V}_n^-$ and $\mathcal{V}_n^+$ are, respectively, the subsets of vertices that are at odd and even distance of the vertex **1**. Every edge of the hypercube graph $\mathcal{Q}_n$ connects a vertex in $\mathcal{V}_n^-$ to one in $\mathcal{V}_n^+$ and so, every edge of the graph $G(C_{n,l}^\star)$ connects a vertex in $C_{n,l}^\star \cap \mathcal{V}_n^-$ to one in $C_{n,l}^\star \cap \mathcal{V}_n^+$. The matrices $Q_{n,l}$ can thus be written in the canonical form (11.9). Denote by $u_{n,l}^\star$ and $v_{n,l}^\star$ the left and right Perron eigenvectors of $Q_{n,l}$ (see (11.12)-(11.15)),

$$u_{n,l}^\star Q_{n,l} = \vartheta_{n,l}(0) u_{n,l}^\star, \quad u_{n,l}^\star > 0, \tag{5.17}$$

$$Q_{n,l} v_{n,l}^\star = \vartheta_{n,l}(0) v_{n,l}^\star, \quad v_{n,l}^\star > 0, \tag{5.18}$$

normalized to make $\sum_{x\in C_{n,l}^\pm} 2u_{n,l}^\star(x) = 1$ and $\sum_{x\in C_{n,l}^\pm} 2u_{n,l}^\star(x)v_{n,l}^\star(x) = 1$, where $C_{n,l}^\pm \equiv C_{n,l}^\star \cap \mathcal{V}_n^\pm$. The next lemma is the key ingredient to the proof Proposition 5.1

Lemma 5.6. *On $\Omega_0 \cap \Omega^\star$, for all but a finite number of indices n we have, for each x in $C_{n,l}^\star$ and all $i \ge \theta_{n,l}^\star$,*

$$P_x\left(T_{n,l}^\star > i\right) = v_{n,l}^\star(x)\vartheta_{n,l}^i(0)(1+\epsilon_n^\star), \tag{5.19}$$

where $|\epsilon_n^\star| \le \exp(-\beta n/4)$.

The proof of this result relies on the following a priori lower bound on $v_{n,l}^\star$.

Lemma 5.7. *For all $x \in C_{n,l}^\star$, $v_{n,l}^\star(x) \ge (1/\varrho_{n,l}(0))^{|C_{n,l}^\star|}$.*

Proof. The proof is a simple adaptation of the proof of Lemma 4.6. □

Proof of Lemma 5.6. By (11.19) and (11.20) of Lemma 11.2 applied to $Q_{n,l}$ we have, for all $x \in C_{n,l}^\star$ and all $i \in \mathbb{N}$,

$$P_x\left(T_{n,l}^\star > i\right) = v_{n,l}^\star(x)\vartheta_{n,l}^i(0)\Big\{1 + \Big(\tfrac{\vartheta_{n,l}(1)}{\vartheta_{n,l}(0)}\Big)^i \widehat{R}_n\Big\}, \tag{5.20}$$

where $|\widehat{R}_n| \le \Big(\min_{x\in C_{n,l}^\star}\big\{v_{n,l}^\star(x)(\pi_{n,l}^\star(x))^{1/2}\big\}\Big)^{-1}$. The lemma will be proved if we can show that on $\Omega_0 \cap \Omega^\star$, for all but a finite number of indices n,

$$\Big(\tfrac{\vartheta_{n,l}(1)}{\vartheta_{n,l}(0)}\Big)^{\theta_{n,l}^\star} |\widehat{R}_n| \le \exp(-\beta n). \tag{5.21}$$

To this end note that, by (5.10), (6.7), and Lemma 5.5, for all $x \in C_{n,l}^\star$, $\pi_{n,l}^\star(x) \ge r_n\left(\rho_n^\star\right)/n|C_{n,l}^\star|\varrho_{n,l}(0)$. Inserting this bound and that of Lemma 5.7 in our bound on $|\widehat{R}_n|$, and using (2.11), Lemma 5.5, and (2.20) we get that on $\Omega_0 \cap \Omega^\star$, for all but a finite number of indices n, $|\widehat{R}_n| \le e^{\beta n(|C_{n,l}^\star|+1/2)\sqrt{\log 2}(1+o(1))}$. Using (5.15) of Corollary 5.4 to bound the ratio $\vartheta_{n,l}(1)/\vartheta_{n,l}(0)$ and choosing $\theta_{n,l}^\star$ as in (5.4) then yield (5.21). The proof of the lemma is done. □

Proof of Proposition 5.1. When $|C^\star_{n,l}| = 2$, $Q_{n,l} = \vartheta_{n,l}(0) I_2$ where I_2 denotes the identity matrix in $\mathbb{R}^2$. Thus $P_x\left(T^\star_{n,l} > i\right) = \vartheta^i_{n,l}(0)$ which by (5.12) leads to (5.5). To prove (5.6) use that by (5.19) with $i = \theta^\star_{n,l}$,

$$v^\star_{n,l}(x) = P_x\left(T^\star_{n,l} > \theta^\star_{n,l}\right) \vartheta^{-\theta^\star_{n,l}}_{n,l}(0)(1+\epsilon^\star_n)^{-1} \le \vartheta^{-\theta^\star_{n,l}}_{n,l}(0)(1+\epsilon^\star_n)^{-1}. \tag{5.22}$$

Hence $P_x\left(T^\star_{n,l} > i\right) \le \vartheta^{-\theta^\star_{n,l}}_{n,l}(0)\vartheta^i_{n,l}(0)$ where, by (5.13), $\vartheta^i_{n,l}(0) \le e^{-i(n/\varrho_{n,l}(0)|C^\star_{n,l}|)(1-o(1))}$ and $\vartheta^{-\theta^\star_{n,l}}_{n,l}(0) \le 1 + \frac{\theta^\star_{n,l} n}{\varrho_{n,l}(0)|C^\star_{n,l}|}(1+o(1))$. □

6. Properties of the effective jump chain J°_n

Thi section gathers needed results on the chain J°_n. The first proposition, which is central to the strategy of Sections 8, 9, and 10, states that J°_n is fast mixing. Define

$$\ell^\circ_n = \left\lceil n^3 \right\rceil. \tag{6.1}$$

Proposition 6.1. *Assume that $c_\star > 1 + \log 4$. For all $\beta > 0$, there exists a subset $\Omega_1 \subset \Omega$ with $\mathbb{P}(\Omega_1) = 1$ such that, on Ω_1, for all but a finite number of indices n, for all pairs $x \in \mathcal{V}^\circ_n, y \in \mathcal{V}^\circ_n$, and all $i \ge 0$,*

$$\left|P^\circ_{\pi^\circ_n}\left(J^\circ_n(i+\ell^\circ_n) = y, J^\circ_n(i) = x\right) - \pi^\circ_n(x)\pi^\circ_n(y)\right| \le \delta_n \pi^\circ_n(x)\pi^\circ_n(y), \tag{6.2}$$

where $0 \le \delta_n \le 2^{-n}$.

Thus the random variables $J^\circ_n(\ell^\circ_n i)$, $i \in \mathbb{N}$, are close to independent and distributed according to the invariant distribution π°_n. The next proposition provides bounds on certain mean local times that are needed to control stretches of trajectories of length ℓ°_n. Recall that $I^\star_n$ is the set of isolated vertices in the partition (2.9).

Proposition 6.2. *Assume that $c_\star > 3$. Then, there exists a subset $\Omega^{\textsc{srw}} \subset \Omega$ with $\mathbb{P}(\Omega^{\textsc{srw}}) = 1$ such that, on $\Omega^{\textsc{srw}} \cap \Omega^\star$, for all but a finite number of indices n, the following holds: there exist constants $0 < C_\circ, C'_\circ < \infty$ such that,*
(i) for all $z \in I^\star_n$,

$$\sum_{l=1}^{\ell^\circ_n - 1} P^\circ\left(J^\circ_n(l+2) = z \mid J^\circ_n(0) = z\right) \le \frac{C_\circ}{n \log n}, \tag{6.3}$$

(ii) for all $1 \le l \le L^\star$ and all $z, z' \in \partial C^\star_{n,l}$,

$$\sum_{l=1}^{\ell^\circ_n - 1} P^\circ\left(J^\circ_n(l) = z \mid J^\circ_n(0) = z'\right) \le \frac{C'_\circ}{n}. \tag{6.4}$$

SRW in $\Omega^{\textsc{srw}}$ above stands for Symmetric Random Walk. The reason for this will become clear from the proof (see Lemma 6.12). One may however already observe that the behavior of J°_n in Proposition 6.1 and Proposition 6.2 is reminiscent of SRW (see e.g. Section 3 of [25]) and as the next proposition shows, so is that of its invariant measure.

Proposition 6.3. *Assume that $c_\star > 2$. For all $\beta > 0$,*

$$\pi^\circ_n(x) = 1/|\mathcal{V}^\circ_n|, \quad x \in \mathcal{V}^\circ_n, \tag{6.5}$$

where, on $\Omega^\star$, for all but a finite number of indices n,

$$|\mathcal{V}^\circ_n| = 2^n\left[1 - n^{-2c_\star+1}(1 + \mathcal{O}(n^{-(c_\star-1)}))\right]. \tag{6.6}$$

Let us immediately give the short proof of Proposition 6.3.

Proof. Because the process X_n has a unique reversible invariant measure, $G_{\beta,n}$, the jump chain also has unique reversible invariant measure, which is the measure defined on $\mathcal{V}_n$ by

$$\pi_n(x) = \frac{\lambda_n(x)G_{\beta,n}(x)}{\sum_{x\in\mathcal{V}_n}\lambda_n(x)G_{\beta,n}(x)} = \frac{\sum_{y:(x,y)\in\mathcal{E}_n} e^{-\beta\max(H_n(y),H_n(x))}}{\sum_{x\in\mathcal{V}_n}\sum_{y:(x,y)\in\mathcal{E}_n} e^{-\beta\max(H_n(y),H_n(x))}}. \tag{6.7}$$

By this and (3.6) $\pi_n^\circ(x) = (nW_{\beta,n}^\circ)^{-1}\sum_{y:(x,y)\in\mathcal{E}_n} e^{-\beta\max(H_n(y),H_n(x))}$, $x\in\mathcal{V}_n^\circ$, where $W_{\beta,n}^\circ = n^{-1}\sum_{x\in\mathcal{V}_n^\circ}\sum_{y:(x,y)\in\mathcal{E}_n} e^{-\beta\max(H_n(y),H_n(x))}$. But by (2.9) and the definition (3.1) of $\mathcal{V}_n^\circ$, $\max(H_n(y),H_n(x)) = 0$ whenever one of the two vertices $\{x,y\}$ lies in $\mathcal{V}_n^\circ$. Hence $W_{\beta,n}^\circ = |\mathcal{V}_n^\circ|$, yielding (6.5). Since $|\mathcal{V}_n^\circ| = 2^n - \sum_{l=1}^{L^\star}|C_{n,l}^\star|$, (6.6) follows from (2.14). □

Our last proposition contains a rough lower bound on hitting times at stationarity that is needed in the proof of Theorem 1.3. Write

$$T^\circ(A) \equiv \inf\{i\in\mathbb{N} \mid J_n^\circ(i)\in A\}, \quad A\subseteq\mathcal{V}_n^\circ. \tag{6.8}$$

Proposition 6.4. *Assume that $c_\star > 1+\log 4$. On $\Omega_1\cap\Omega^\star$, for all but a finite number of indices n, we have that for all $A\subseteq\mathcal{V}_n^\circ$ and for $I_n^\star$ as in (2.9),*

$$P_{\pi_n^\circ}^\circ\left(T^\circ(A\cap I_n^\star) > t\right) \geq (1+o(1))\exp\left(-2t|A\cap I_n^\star|/|\mathcal{V}_n^\circ|\right) - \mathcal{O}(\tfrac{1}{\log n}), \quad t>0. \tag{6.9}$$

The rest of this section is organized as follows. The proof of Proposition 6.1, given in Subsection 6.3, relies on a bound on the spectral gap of J_n° established in Subsection 6.2, which itself relies on estimates on the transition probabilities of J_n° that are established in Subsection 6.1. The proof of Proposition 6.2 and Proposition 6.4 are carried out in Subsection 6.4 and Subsection 6.5, respectively

6.1. **Estimates on the transition probabilities.** We now examine the transition probabilities (3.5) of J_n°. In what follows we denote by $G^\star(A)$ the complete graph on A. Let $G^\circ(\mathcal{V}_n^\circ)$ be the graph with vertex set $\mathcal{V}_n^\circ$ such that (x,y) is an egde of the graph if and only if $p_n^\circ(x,y)>0$. In view of (3.4)-(3.5),

$$G^\circ(\mathcal{V}_n^\circ) = G(\mathcal{V}_n^\circ)\bigcup_{1\leq l\leq L^\star} G^\star(\partial C_{n,l}^\star). \tag{6.10}$$

Proposition 6.5. *For all $(x,y)\in G(\mathcal{V}_n^\circ)$,*

$$p_n^\circ(x,y) = 1/n, \tag{6.11}$$

and, for all $1\leq l\leq L^\star$ and all (x,y) in $G^\star(\partial C_{n,l}^\star)$,

$$p_n^\circ(x,y) = \frac{m_{n,l}^\star(x)m_{n,l}^\star(y)}{\sum_{z\in\partial C_{n,l}^\star} m_{n,l}^\star(z)}(1+o(1)), \tag{6.12}$$

where $nm_{n,l}^\star(x)$ is the number of vertices of $C_{n,l}^\star$ that are are distance one from x,

$$m_{n,l}^\star(x) \equiv n^{-1}|\left\{y\in C_{n,l}^\star \mid \mathrm{dist}(y,x)=1\right\}|, \;\; x\in\partial C_{n,l}^\star. \tag{6.13}$$

Proof. Clearly, if $(x,y)\in G(\mathcal{V}_n^\circ)$, $p_n^\circ(x,y) = p_n(x,y) = 1/n$, yielding (6.11). We now turn to (6.12). Let us first state two useful a piori relations

$$p_n^\circ(x,y) = p_n^\circ(y,x) \quad \forall (x,y)\in G^\circ(\mathcal{V}_n^\circ), \tag{6.14}$$

$$m_{n,l}^\star(y) = \sum_{x\in\partial C_{n,l}^\star} p_n^\circ(x,y) \quad \forall y\in\partial C_{n,l}^\star. \tag{6.15}$$

Eq. (6.14) is reversibility. Eq. (6.15) follows from the relation $\sum_y p_n^\circ(x,y) = 1$, (6.11), (6.14), and the definition (6.13).

Given $A \subseteq \mathcal{V}_n$ write $T(A) \equiv \inf\{i \in \mathbb{N} \mid J_n(i) \in A\}$. Also recall that for $1 \le l \le L^\star$, $T^\star_{n,l} \equiv \inf\{i \in \mathbb{N} \mid J_n(i) \in \partial C^\star_{n,l}\}$. Then, for all $(x,y) \in G^\star(\partial C^\star_{n,l})$,

$$p^\circ_n(x,y) = \textstyle\sum_{z \in C^\star_{n,l}} p_n(x,z) P_z\left(J_n(T^\star_{n,l}) = y\right). \tag{6.16}$$

The next lemma establishes that the exit distribution from $C^\star_{n,l}$ is independent from the entrance point, provided that the exit probability is not too small.

Lemma 6.6. *For any two distinct vertices z and $\bar z$ in $C^\star_{n,l}$ and any $y \in \partial C^\star_{n,l}$,*

$$P_z\left(J_n(T^\star_{n,l}) = y\right) = (1-\tilde\epsilon_n) P_{\bar z}\left(J_n(T^\star_{n,l}) = y\right) + \tilde\epsilon_n, \tag{6.17}$$

where $\tilde\epsilon_n \le |\partial C^\star_{n,l}|/\varrho_{n,l}(1)$.

Proof of Lemma 6.6. Note that for any two vertices z and $\bar z$ in $C^\star_{n,l}$,

$$\begin{aligned} P_z\left(T^\star_{n,l} \le T(\bar z)\right) &= \textstyle\sum_{y \in \partial C^\star_{n,l}} P_z\left(T(y) \le T(\bar z \cup (\partial C^\star_{n,l}))\right) && (6.18)\\ &= \textstyle\sum_{y \in \partial C^\star_{n,l}} \frac{\pi_n(y)}{\pi_n(z)} P_y\left(T(z) \le T(\bar z \cup (\partial C^\star_{n,l}))\right) && (6.19)\\ &\le |\partial C^\star_{n,l}| \varrho^{-1}_{n,l}(1), && (6.20)\end{aligned}$$

where the second equality is reversibility. Next decompose the event $\{J_n(T^\star_{n,l}) = y\}$ according to whether $\{T(\bar z) \ge T^\star_{n,l}\}$ or $\{T(\bar z) < T^\star_{n,l}\}$: by the strong Markov property,

$$P_z\left(T(\bar z) < T^\star_{n,l}, J_n(T^\star_{n,l}) = y\right) = P_z\left(T(\bar z) < T^\star_{n,l}\right) P_{\bar z}\left(J_n(T^\star_{n,l}) = y\right), \tag{6.21}$$

whereas $P_z\left(T^\star_{n,l} \le T(\bar z), J_n(T^\star_{n,l}) = y\right) \le P_z\left(T^\star_{n,l} \le T(\bar z)\right)$. Eq. (6.17) now follows. □

Now pick an arbitrary vertex $z^\star_{n,l} \in C^\star_{n,l}$ and denote by $\mathcal{L}^\star_{n,l}$ the exit distribution

$$\mathcal{L}^\star_{n,l}(y) = P_{z^\star_{n,l}}\left(J_n(T^\star_{n,l}) = y\right), \quad y \in \partial C^\star_{n,l}. \tag{6.22}$$

Lemma 6.7. *For all $z \in C^\star_{n,l}$ and $y \in \partial C^\star_{n,l}$*

$$P_z\left(J_n(T^\star_{n,l}) = y\right) = (1+o(1))\mathcal{L}^\star_{n,l}(y). \tag{6.23}$$

Proof of Lemma 6.7. We readily deduce from Lemma 6.6 that if $y \in \partial C^\star_{n,l}$ is such that $\mathcal{L}^\star_{n,l}(y) \ge n\tilde\epsilon_n$, then $P_z\left(J_n(T^\star_{n,l}) = y\right) = (1+o(1))\mathcal{L}^\star_{n,l}(y)$, otherwise

$$P_z\left(J_n(T^\star_{n,l}) = y\right) < (n+1)\tilde\epsilon_n. \tag{6.24}$$

Let us prove by contradiction that $\mathcal{L}^\star_{n,l}(y) \ge n\tilde\epsilon_n$ for all $y \in \partial C^\star_{n,l}$. Assume that there exists $y \in \partial C^\star_{n,l}$ such that $\mathcal{L}^\star_{n,l}(y) < n\tilde\epsilon_n$. Then, by (6.24) and (6.16),

$$p^\circ_n(x,y) \le (n+1)\tilde\epsilon_n \textstyle\sum_{z \in C^\star_{n,l}} p_n(x,z) = (n+1)\tilde\epsilon_n m^\star_{n,l}(x). \tag{6.25}$$

Summing both sides over $x \in \partial C^\star_{n,l}$,

$$\textstyle\sum_{x \in \partial C^\star_{n,l}} p^\circ_n(x,y) \le (n+1)\tilde\epsilon_n \sum_{x \in \partial C^\star_{n,l}} m^\star_{n,l}(x) \le n^5 \varrho^{-1}_{n,l}(1) \ll n^{-1}. \tag{6.26}$$

However, by (6.15), $\sum_{x \in \partial C^\star_{n,l}} p^\circ_n(x,y) = m^\star_{n,l}(x) \ge n^{-1}$, which is a contradiction. □

We are now ready to conclude the proof of 6.6. By (6.16) and (6.23),

$$p^\circ_n(x,y) = m^\star_{n,l}(x)\mathcal{L}^\star_{n,l}(y)(1+o(1)). \tag{6.27}$$

Inserting this in (6.14) and summing both sides over $x \in \partial C^\star_{n,l}$ we get

$$\mathcal{L}^\star_{n,l}(y) = \tfrac{m^\star_{n,l}(y)}{\sum_{x \in \partial C^\star_{n,l}} m^\star_{n,l}(x)}(1+o(1)), \tag{6.28}$$

and inserting this in turn in (6.27) yields (6.12). The proof of the proposition is done. □

6.2. **Bound on the spectral gap of J_n°.** Let

$$1 = \vartheta_n^\circ(0) > \vartheta_n^\circ(1) \geq \vartheta_n^\circ \cdots \geq \vartheta_{n,l}(|\mathcal{V}_n^\circ| - 1) > -1 \tag{6.29}$$

denote the eigenvalues of the matrix with entries (3.5) and set $\tau_n^\circ = 1/(1 - \vartheta_n^\circ(1))$. The proof of Proposition 6.1 relies on the following lower bound on the spectral gap of J_n°.

Proposition 6.8. *Assume that $c_\star > 1 + \log 4$. For all $\beta > 0$, there exists a subset $\Omega_2 \subset \Omega$ with $\mathbb{P}(\Omega_2) = 1$ such that, on Ω_2, for all but a finite number of indices n,*

$$\tau_n^\circ \leq \tfrac{1}{2} n^2 (1 + o(1)). \tag{6.30}$$

Proof. The proof of Proposition 6.8 is based on a bound on spectral gaps already stated in Subsection 4.1 which now reads as follows: if $\Gamma_n^\circ = \{\gamma_{x,y}^\circ\}$ is a set of paths in the graph $G^\circ(\mathcal{V}_n^\circ)$ such as described in the paragraph above (4.27), then

$$\tau_n^\circ \leq \max_e \rho_n^{-1}(e) \sum_{\gamma_{x,y}^\circ \ni e} \left|\gamma_{x,y}^\circ\right| \pi_n^\circ(x) \pi_n^\circ(y), \tag{6.31}$$

where the max is over all edges $e = \{x', y'\}$ of $G^\circ(\mathcal{V}_n^\circ)$, $\rho_n(e) \equiv \pi_{n,l}^\circ(x') p_n^\circ(x', y')$, and the summation is over all paths $\gamma_{x,y}^\circ$ in Γ_n° that pass through e. The structure (6.10) of the graph $G^\circ(\mathcal{V}_n^\circ)$ naturally prompts us to write

$$\Gamma_n^\circ = \Gamma_n \cup \left(\cup_{1 \leq l \leq L^\star} \Gamma_{n,l}^\star\right), \tag{6.32}$$

where Γ_n is a set of paths in $G(\mathcal{V}_n^\circ)$ and, for each $1 \leq l \leq L^\star$,

$$\Gamma_{n,l}^\star = \left\{\gamma_{x,y} \equiv (x, y),\, x, y \in \partial C_{n,l}^\star\right\}. \tag{6.33}$$

That is, the paths of $\Gamma_{n,l}^\star$ simply are the edges of $G^\star(\partial C_{n,l}^\star)$. Eq. (6.31) then becomes

$$\tau_n^\circ \leq \max\left\{\tau_n, \max_{1 \leq l \leq L^\star} \tau_{n,l}^\star\right\}, \tag{6.34}$$

where $\tau_{n,l}^\star \equiv \max_{e=(x,y), x,y \in \partial C_{n,l}^\star} \pi_n^\circ(y)/p_n^\circ(x,y)$, and

$$\tau_n \equiv \max_e \rho_n^{-1}(e) \sum_{\gamma_{x,y} \ni e} |\gamma_{x,y}|\, \pi_n^\circ(x) \pi_n^\circ(y), \tag{6.35}$$

the max now being over all edges e of $G(\mathcal{V}_n^\circ)$, and the summation over all paths $\gamma_{x,y}$ in Γ_n that pass through e. Using Proposition 6.3 and the fact that $\max(H_n(z), H_n(z')) = 0$ whenever $(z, z') \in \mathcal{E}_n$ is an edge with at least one endpoint in $\mathcal{V}_n^\circ$, (6.35) reduces to

$$\tau_n = (n/|\mathcal{V}_n^\circ|) \max_{e=(x,x') \in G(\mathcal{V}_n^\circ)} \sum_{\gamma_{y,y'} \ni e} |\gamma_{y,y'}|. \tag{6.36}$$

The quality of the bound (6.36) now depends on making a judicious choice of the set of paths Γ_n. We will adopt a very clever choice made in [24].

• ***A choice of*** **Γ_n.** The set Γ_n is defined as

$$\Gamma_n = \left\{\gamma_{x,y} \in \Gamma_n',\, x, y \in \mathcal{V}_n^\circ\right\}, \tag{6.37}$$

where Γ_n' is a subset of paths in $G(\mathcal{V}_n)$ constructed as follows. Given $i \in \{1, \dots n\}$, and given two vertices x and $x' \in \mathcal{V}_n$ such that $x_i \neq x_i'$, let $\gamma_{x,x'}^i$ be the path obtained by going left to right cyclically from x to x', successively flipping the disagreeing coordinates, starting from the i-th coordinate. Set $\Gamma_n^i = \left\{\gamma_{x,x'}^i, x, x' \in \mathcal{V}_n\right\}$, $1 \leq i \leq n$. These paths are ordered in an obvious way. Given x, x' and $\gamma_{x,x'}$, let $\overline{\gamma}_{x,x'}$ be the set of vertices visited by the path $\gamma_{x,x'}$, and let $\gamma_{x,x'}^{int} = \overline{\gamma}_{x,x'} \setminus \{x, x'\}$ be the subset of "interior" vertices. We next split the set of vertices $\mathcal{V}_n$ into *good* ones and *bad* ones. Recalling (2.9), we say that a vertex is good if it belongs to $N_n^\star$; otherwise it is bad. We say that a path γ is good if all its interior points γ^{int} are good, and that a set of paths is good if all its elements are good.

The (random) set of path Γ_n' is then constructed as follows:

(i) Consider pairs x and x' such that $\mathrm{dist}(x,x') \geq n/\log n$. If $\{\gamma^i_{x,x'}, 1 \leq i \leq n\}$ contains a good path, choose the first such for Γ'_n; otherwise choose $\gamma^1_{x,x'}$.
(ii) Consider pairs x and x' such that $\mathrm{dist}(x,x') < n/\log n$. If there is a good vertex $x'' \in \mathcal{V}_n$ such that $\mathrm{dist}(x,x'') \geq n/\log n$ and $\mathrm{dist}(x'',x') \geq n/\log n$, and if there are good paths, one in $\{\gamma^i_{x,x''}, 1 \leq i \leq n\}$ and one in $\{\gamma^i_{x'',x'}, 1 \leq i \leq n\}$, such that the union of these two good paths is a self avoiding path of length less than n, select this union as the path connecting x to x' in Γ'_n (notice that this is a good path); otherwise choose $\gamma^1_{x,x'}$.

The key point of this construction is that Γ'_n is almost surely good. More precisely, set $\Omega^{\text{GOOD}}_n = \{\omega \in \Omega \,\big|\, \Gamma'_n \equiv \Gamma'_n(\omega) \text{ is good}\}$, $n \geq 1$, and $\Omega^{\text{GOOD}} = \liminf_{n\to\infty} \Omega^{\text{GOOD}}_n$.

Proposition 6.9 (Proposition 4.1 of [24]). *If $c_\star > 1 + \log 4$ then $\mathbb{P}\big(\Omega^{\text{GOOD}}\big) = 1$.*

Going back to (6.37), we see that the set Γ_n is obtained from Γ'_n by removing the paths whose endpoints lie in $\cup_{1\leq l\leq L^\star} C^\star_{n,l}$. Thus on Ω^{GOOD} the paths of Γ_n only visit vertices in $\mathcal{V}^\circ_n$. This finishes our construction. Note that the paths constructed in this way have length smaller than n. Thus (6.36) yields

$$\tau_n \leq (n^2/|\mathcal{V}^\circ_n|)\max_{e\in G(\mathcal{V}^\circ_n)} |\{\gamma \in \Gamma \mid e \in \gamma\}| \,. \tag{6.38}$$

• ***Bound on*** τ_n**.** From now on we assume that $\omega \in \Omega^{\text{GOOD}}$ so that, for all large enough n, $\Gamma_n \equiv \Gamma_n(\omega)$ is good. In that case a bad vertex can appear only at the ends of any path. Let us write

$$\tau_n = (n^2/|\mathcal{V}^\circ_n|)(\tau^1_n + \tau^2_n) \,, \tag{6.39}$$

where τ^1_n, repectively τ^2_n, is obtained by restricting the sum in (6.38) to paths connecting vertices at distance $n/\log n$ or more apart, repectively, less than $n/\log n$ apart.

On the one hand it is well known that (see e.g. Example 2.2, p. 45 in [21])

$$\tau^1_n \leq 2^{n-1}. \tag{6.40}$$

On the other hand, arguing as in [24] (see Subsection 4.2.2, page 934) that the sum in τ^2_n is over a set of paths that connect vertices in a hypercube of dimension at most $n/\log n$ around e, we have

$$\tau^2_n \leq 2^{2n/\log n}. \tag{6.41}$$

Plugging (6.40) and (6.41) in (6.39), and using (6.6) of Proposition 6.3 to bound $|\mathcal{V}^\circ_n|$, we we get that on $\Omega^{\text{GOOD}} \cap \Omega^\star$, for large enough n,

$$\tau_n \leq n^2 2^{-n} \left[1 - n^{-2c_\star+1}(1+o(1))\right]^{-1} \left(2^{n-1} + 2^{2n/\log n}\right) \leq (n^2/2)(1+o(1)). \tag{6.42}$$

• ***Bound on*** $\tau^\star_{n,l}, 1 \leq l \leq L^\star$**.** Consider the terms $\tau^\star_{n,l}$ from (6.34). By Proposition 6.3 and (6.12) of Proposition 6.5, on $\Omega_0 \cap \Omega^\star$, for large enough n we have, for all $e = (x,y), x, y \in \partial C^\star_{n,l}$ and all $1 \leq l \leq L^\star$,

$$\frac{\pi^\circ_n(y)}{p^\circ_n(x,y)} \leq \frac{1}{|\mathcal{V}^\circ_n|} \frac{n\sum_{z\in\partial C^\star_{n,l}} m^\star_{n,l}(z)}{m^\star_{n,l}(x) m^\star_{n,l}(y)}(1+o(1)) \leq \frac{n^2|C^\star_{n,l}|^2}{|\mathcal{V}^\circ_n|} \leq 2n^4 2^{-n}. \tag{6.43}$$

Thus, on $\Omega_0 \cap \Omega^\star$, for large enough n,

$$\max_{1\leq l\leq L^\star} \tau^\star_{n,l} \leq 2n^4 2^{-n}. \tag{6.44}$$

Setting $\Omega_2 = \Omega^{\text{GOOD}} \cap \Omega_0 \cap \Omega^\star$ and plugging (6.44) and (6.42) in (6.34) finally yields the upper bound (6.30) on τ°_n. The proof of Proposition 6.8 is done. □

6.3. **Proof of Proposition 6.1.** Consider the continuous time Markov chain $(J_n^*(t), t > 0)$ with jump chain $(J_n^\circ(k), k \in \mathbb{N})$ and rate one exponential waiting times. That is, given a family $(e_{n,i}^*, i \in \mathbb{N})$ of independent mean one exponential r.v.'s, independent of J_n°,

$$J_n^*(t) = J_n^\circ(i) \text{ if } s_n(i) \le t < s_n(i+1) \text{ for some } i, \tag{6.45}$$

where $s_n(k) \equiv \sum_{i=0}^{k-1} e_{n,i}^*$, $k \in \mathbb{N}$. Write P_x^* for the law of J_n^* started in x. We know that:

Lemma 6.10 (Proposition 3 of [21])**.** *For all $x \in \mathcal{V}_n^\circ$ and all $t \in (0, \infty)$,*

$$\left\| P_x^* \left(J_n^*(t) = \cdot \right) - \pi_n^\circ(\cdot) \right\|_{TV} \le \sqrt{(1 - \pi_n^\circ(x))/(4\pi_n^\circ(x))} e^{-t/\tau_n^\circ}. \tag{6.46}$$

We now aim to compare the discrete and continuous time transition kernels. Set

$$\mathcal{I}(\zeta) = \zeta - \log(1+\zeta), \quad \zeta > 0. \tag{6.47}$$

Lemma 6.11. *Let $0 < m_n \uparrow \infty$ be an integer valued sequence. For all $\zeta > 0$ we have*

$$|P_x^\circ \left(J_n^\circ(m_n) = y \right) - \pi_n^\circ(y)| \le (4\pi_n^\circ(x))^{-1/2} e^{-m_n(1-\zeta)/\tau_n^\circ} + 2e^{-m_n \mathcal{I}(\zeta)}, \quad \forall x, y \in \mathcal{V}_n^\circ.$$

Proof. By (6.45), $P_x^\circ \left(J_n^\circ(m_n) = y \right) = P_x^* \left(J_n^*(s_n(m_n)) = y \right)$. Now consider the event $A_{\zeta,n} = \{|s_n(m_n) - m_n| \ge \zeta m_n\}$, $\zeta > 0$. A classical large deviation estimates yields

$$P_x^* \left(A_{\zeta,n} \right) \le 2e^{-m_n \mathcal{I}(\zeta)}. \tag{6.48}$$

Hence

$$0 \le P_x^* \left(J_n^*(s_n(m_n)) = y, A_{\zeta,n} \right) \le 2e^{-m_n \mathcal{I}(\zeta)}. \tag{6.49}$$

Writing $A_{\zeta,n}^c = \{|s_n(m_n) - m_n| < \zeta m_n\}$,

$$\begin{aligned} P_x^* \left(J_n^*(s_n(m_n)) = y, A_{\zeta,n}^c \right) &\le \sup_{|t-m_n|<\zeta m_n} P_x^* \left(J_n^*(t) = y, A_{\zeta,n}^c \right) \\ &\le \sup_{|t-m_n|<\zeta m_n} P_x^* \left(J_n^*(t) = y \right) \\ &\le \pi_n^\circ(y) + (4\pi_n^\circ(x))^{-1/2} e^{-m_n(1-\zeta)/\tau_n^\circ}, \end{aligned} \tag{6.50}$$

where the last inequality follows from Lemma 6.10. Similarly,

$$P_x^* \left(J_n^*(s_n(m_n)) = y, A_{\zeta,n}^c \right) \ge \inf_{|t-m_n|<\zeta m_n} P_x^* \left(J_n^*(t) = y, A_{\zeta,n}^c \right). \tag{6.51}$$

But $P_x^* \left(J_n^*(t) = y, A_{\zeta,n}^c \right) \ge P_x^* \left(J_n^*(t) = y \right) - 2e^{-m_n \mathcal{I}(\zeta m_n)}$ so that, by Lemma 6.10 again,

$$P_x^* \left(J_n^*(s_n(m_n)) = y, A_{\zeta,n}^c \right) \ge \pi_n^\circ(y) - (4\pi_n^\circ(x))^{-1/2} e^{-m_n(1+\zeta)/\tau_n^\circ} - 2e^{-m_n \mathcal{I}(\zeta m_n)}. \tag{6.52}$$

Combining (6.49), (6.50), and (6.52) proves the claim of the lemma. □

We can now proceed with the proof of Proposition 6.1. Since π_n° is the invariant measure of J_n°, we have $P_{\pi_n^\circ}^\circ \left(J_n^\circ(i + \ell_n^\circ) = y, J_n^\circ(0) = x \right) = P_x^\circ \left(J_n^\circ(i + \ell_n^\circ) = y \right) \pi_n^\circ(x)$. It thus suffices to show that there exists a subset $\Omega_1 \subset \Omega$ with $\mathbb{P}(\Omega_1) = 1$ such that, on Ω_1, for all but a finite number of indices n, for all pairs $x \in \mathcal{V}_n^\circ, y \in \mathcal{V}_n^\circ$, and all $i \ge 0$,

$$|P_x^\circ \left(J_n^\circ(i + \ell_n^\circ) = y \right) - \pi_n^\circ(y)| \le \delta_n \pi_n^\circ(y), \tag{6.53}$$

for some $0 \le \delta_n \le 2^{-n}$. To do this choose $\zeta = 1/2$ and $m_n = \ell_n^\circ$ in Lemma 6.11. By (6.1) and Proposition 6.8, on Ω_2, $\ell_n^\circ / \tau_n^\circ \ge 2n(1 - o(1))$ for all n large enough. Futhermore, by Proposition 6.3, on $\Omega^\star$, for all n large enough, $(\pi_n^\circ(y)\sqrt{4\pi_n^\circ(x)})^{-1} \le 2^{3n/2}$. Thus, on $\Omega_1 \equiv \Omega^\star \cap \Omega_2$, for all n large enough,

$$\delta_n \equiv (\pi_n^\circ(y)\sqrt{4\pi_n^\circ(x)})^{-1} e^{-\ell_n^\circ(1-\zeta)/\tau_n^\circ} + (\pi_n^\circ(y))^{-1} 2e^{-\ell_n^\circ \mathcal{I}(\zeta m_n)} \le 2^{-n}. \tag{6.54}$$

Inserting (6.54) in (6.46) yields the claim of 6.53. The proof of Proposition 6.1 is done.

6.4. **Mean local times: proof of Proposition 6.2.** Let J_n^{SRW} and P^{SRW} denote, respectively, the symmetric random walk on $\mathcal{V}_n$ (hereafter SRW) and its law. More precisely,

$$p_n^{\mathsf{SRW}}(x,y) \equiv P^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(1)=y \mid J_n^{\mathsf{SRW}}(1)=x\right) = \begin{cases} \frac{1}{n} & \text{if } \operatorname{dist}(x,y)=1, \\ 0, & \text{else.} \end{cases} \tag{6.55}$$

We also write P_x^{SRW} for the law of J_n^{SRW} started in x. The proof of Proposition 6.2 relies on three key properties of J_n^{SRW} that we state below in the form of three lemmata.

Our first lemma is a reformulation of Theorem 1.1 of [16] on the hitting time of so-called percolation clouds, namely, sets of the form $V_n(\rho)$ (see (2.1)). For $A \subset \mathcal{V}_n$ let $T^{\mathsf{SRW}}(A)$ be the hitting time

$$T^{\mathsf{SRW}}(A) = \inf\left\{k \in \mathbb{N} : J_n^{\mathsf{SRW}}(k) \in A\right\}. \tag{6.56}$$

Lemma 6.12. *Let $\rho_n^\star$ be as in (2.3) for some $c_\star$ such that $n^{c_\star} \gg n\log n$. There exists a subset $\Omega^{\mathsf{SRW}} \subset \Omega$ with $\mathbb{P}\left(\Omega^{\mathsf{SRW}}\right)=1$ such that, on Ω^{SRW}, for all but a finite number of indices n the following holds: for all sequences $l_n > 0$ such that $l_n/n^{c_\star} \le C$ for some constant $0<C<\infty$,*

$$\max_{x\in\mathcal{V}_n}\left|P_x^{\mathsf{SRW}}\left(T^{\mathsf{SRW}}(V_n(\rho_n^\star)\setminus x) \ge l_n\right) - e^{-l_n/n^{c_\star}}\right| \le C'\left[\frac{1}{n}+\frac{1}{\log n^{c_\star}}+\frac{n\log n}{n^{c_\star}}\right], \tag{6.57}$$

where $0<C'<\infty$ is a numerical constant.

The next two lemmata bound the mean number of returns to a given vertex, z, respectively the mean local time in z, in a time interval of the form $\{3,\dots,m\}$, $m \le \lceil n^3\rceil$.

Lemma 6.13. *For all $m \le \lceil n^3\rceil$, all $z\in\mathcal{V}_n$, and $a\in\{0,1,2,3\}$,*

$$\sum_{l=1}^{m} P_z^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(l+a)=z\right) \le \frac{c}{n^b}, \quad b=\begin{cases}1, & \text{if } a\in\{0,1\}\\ 2, & \text{if } a\in\{2,3\}\end{cases}, \tag{6.58}$$

where $0<c<\infty$ is a numerical constant.

Proof. The lemma is proved in exactly the same way as Proposition 3.2 of [25]. □

Lemma 6.14. *For all $m \le \lceil n^3\rceil$ and all y,z such that $\operatorname{dist}(y,z)\ge 1$,*

$$\sum_{l=1}^{m} P_y^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(l)=z\right) \le \frac{c'}{n} \tag{6.59}$$

where $0<c'<\infty$ is a numerical constant.

Proof of Lemma 6.14. The proof draws on the results of [7] where a d-dimensional version of the Ehrenfest scheme, called lumping, was introduced and analyzed (hereafter and whenever possible we use the notations of [7]). Without loss of generality we may take $y\equiv 1$ to be the vertex all of whose coordinates take the value 1. Let γ^Λ be the map (1.7) of [7] derived from the partition of $\Lambda\equiv\{1,\dots,n\}$ into $d=2$ classes, $\Lambda=\Lambda_1\cup\Lambda_2$, defined through the relation: $i\in\Lambda_1$ if the i^{th} coordinate of z is 1, and $i\in\Lambda_2$ otherwise. The resulting lumped chain, $X_n^\Lambda \equiv \gamma^\Lambda(J_n^{\mathsf{SRW}})$, has range $\Gamma_{n,2}=\gamma^\Lambda(\mathcal{V}_n)\subset[-1,1]^2$. Note that the vertices y and z of $\mathcal{V}_n$ are mapped, respectively, onto the corners $1\equiv(1,1)$ and $x\equiv(1,-1)$ of $[-1,1]^2$. Denoting by $\mathbb{P}^\Lambda$ the law of X_n^Λ, we have,

$$P_y^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(l)=z\right) = \mathbb{P}^\Lambda(X_n^\Lambda(l)=x \mid X_n^\Lambda(0)=1). \tag{6.60}$$

Write $\tau_x^{x'} = \inf\{k>0 \mid X_n^\Lambda(0)=x', X_n^\Lambda(k)=x\}$. Without loss of generality we may assume that $0\in\Gamma_{n,2}$ (namely, both Λ_1 and Λ_2 have even cardinality). Then, decomposing

(6.60) according to whether, starting from 1, X_n^Λ visits 0 before it visits x or not, we get: $\mathbb{P}^\Lambda(X_n^\Lambda(l) = x \mid X_n^\Lambda(0) = 1) = A + B$,

$$A = \mathbb{P}^\Lambda(X_n^\Lambda(l) = x, \tau_0^1 < \tau_x^1), \tag{6.61}$$

$$B = \mathbb{P}^\Lambda(X_n^\Lambda(l) = x, \tau_0^1 \geq \tau_x^1). \tag{6.62}$$

By Theorem 3.2 of [7], for all y, z such that $\mathrm{dist}(z, y) \geq 5$,

$$B \leq \mathbb{P}^\Lambda(\tau_x^1 \leq \tau_0^1) \leq F_{n,2}(\mathrm{dist}(z, y)) \leq c_1 n^{-4} \tag{6.63}$$

for some constant $0 < c_1 < \infty$. Of course $A = 0$ for all l such that $l < n/2$ since the chain X_n^Λ needs at least $n/2$ steps to travel from the vertex 1 to 0. To bound A when $l \geq n/2$ we condition on the time of the last visit to 0 before time l, and bound the probability of the latter event by 1. This readily yields

$$A \leq l\mathbb{P}^\Lambda(\tau_x^0 < \tau_0^0) = l\frac{\mathbb{Q}_n(x)}{\mathbb{Q}_n(0)}\mathbb{P}^\Lambda(\tau_0^x < \tau_x^x) \leq l\frac{\mathbb{Q}_n(x)}{\mathbb{Q}_n(0)}, \tag{6.64}$$

where the equality in the middle is reversibility, and where $\mathbb{Q}_n$, defined in Lemma 2.2 of [7], denotes the invariant measure of X_n^Λ. We are thus left to estimate the ratio of invariant masses in (6.64). By (2.4) of [7] we get that $\frac{\mathbb{Q}_n(x)}{\mathbb{Q}_n(0)} \leq |\{x' \in \mathcal{V}_n \mid \gamma^\Lambda(x') = 0\}|^{-1} \leq e^{-c_2 n}$ for some constant $0 < c_2 < \infty$. Gathering our bounds we get that for all y, z such that $\mathrm{dist}(y, z) \geq 5$,

$$P_y^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(l) = z\right) = A + B \leq c_1 n^{-4} + le^{-c_2 n} \leq c_3 n^{-4} \tag{6.65}$$

for some constant $0 < c_3 < \infty$, so that for all $m \leq \lceil n^3 \rceil$,

$$\textstyle\sum_{l=1}^m P_y^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(l) = z\right) \leq c_3 n^{-1}. \tag{6.66}$$

It remains to deal with the cases $1 \leq \mathrm{dist}(y, z) \leq 4$. Thus assume from now on that $1 \leq \mathrm{dist}(y, z) \leq 4$. Consider the event $\mathcal{A}_z \equiv \{\forall i \leq l\, \mathrm{dist}(J_n^{\mathsf{SRW}}(i), z) < 5\}$ and denote by $\mathcal{A}_z^c$ its complement. Decomposing $\mathcal{A}_z^c$ on the place and time of the first visit of the chain to the ball of radius 5, we readily get by the Markov property and (6.65) that

$$P_y^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(l) = z, \mathcal{A}_z^c\right) \leq c_3 n^{-4}. \tag{6.67}$$

Next, by reversibility (the invariant measure of J_n^{SRW} being the uniform measure),

$$P_y^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(l) = z, \mathcal{A}_z\right) = P_z^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(l) = y, \mathcal{A}_z\right) \leq P_z^{\mathsf{SRW}}\left(\mathcal{A}_z\right). \tag{6.68}$$

Let us thus estimate the probability $P_z^{\mathsf{SRW}}\left(\mathcal{A}_z\right)$ that starting in z, the chain did not exit a ball of radius 4 centered at z by time l. This means that at every step it takes the chain flips a coordinate of z in such a way that the total number of coordinates of z and $J_n^{\mathsf{SRW}}(i)$ that disagree is at most 4 for each $i \leq l$. If $l \geq 4$, this implies that $(l-4)/2$ of its l steps (respectively, $(l-4+1)/2$ of them) consist in flipping back a coordinate to its initial position if $l - 4$ is even (respectively, if $l - 4$ is odd). Each time such a backward flip occurs the chain chooses one in at most 4 flipped coordinates. Thus, for all $l \geq 4$,

$$P_y^{\mathsf{SRW}}\left(\mathcal{A}_z\right) \leq (4/n)^{\frac{l-4}{2}}\, \mathbb{1}_{l\,\text{even}} + (4/n)^{\frac{l-3}{2}}\, \mathbb{1}_{l\,\text{odd}}. \tag{6.69}$$

Plugging (6.69) in (6.68) yields that for all y, z such that $1 \leq \mathrm{dist}(y, z) \leq 4$,

$$\textstyle\sum_{l=5}^m P_y^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(l) = z, \mathcal{A}_z\right) \leq c_4 n^{-1}, \tag{6.70}$$

for all $m \leq \lceil n^3 \rceil$ and some constant $0 < c_4 < \infty$, while by simple combinatorics,

$$\textstyle\sum_{l=1}^4 P_y^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(l) = z, \mathcal{A}_z\right) \leq \sum_{l=1}^4 P_y^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(l) = z\right) \leq c_5 n^{-1}, \tag{6.71}$$

for some $0 < c_5 < \infty$. Combining (6.66), (6.70) and (6.71) finishes the proof. □

We are now ready to give the proof of Proposition 6.2.

Proof of Proposition 6.2, (i). Given $y \in \mathcal{V}_n$ denote respectively by P°_y, P_y, and P^{SRW}_y the laws of J°_n, J_n, and J^{SRW}_n started in y. The idea behing the proof is to decompose the paths of J_n at visits to the set $\mathcal{V}^\star_n \equiv \cup_{1\le l\le L^\star} C^\star_{n,l}$, and use that, away from this set, J_n reduces to SRW. To this end define $T^{\mathsf{SRW}}(A) \equiv \inf\{i \in \mathbb{N} \mid J^{\mathsf{SRW}}_n(i) \in A\}$, $A \subseteq \mathcal{V}_n$, and set

$$T^{\mathsf{SRW},\star}_n \equiv \inf\left\{k \in \mathbb{N} : J^{\mathsf{SRW}}_n(k) \in \mathcal{V}^\star_n\right\}. \tag{6.72}$$

Similarly set

$$T^\star_n \equiv \inf\left\{i \in \mathbb{N} \mid J_n(i) \in \mathcal{V}^\star_n\right\}. \tag{6.73}$$

Let $z \in I^\star_n$ be fixed. Using that by definition $J^\circ_n(i) \equiv J_n(T^\circ_{n,i})$, we may write

$$\textstyle\sum_{k=1}^{\ell^\circ_n-1} P^\circ_z\left(J^\circ_n(k+2)=z\right) = \sum_{k=1}^{\ell^\circ_n-1} P_z\left(J_n(T^\circ_{n,k+2})=z\right) = I_1 + I_2 \tag{6.74}$$

where

$$I_1 \equiv \textstyle\sum_{k=1}^{\ell^\circ_n-1} P_z\left(J_n(T^\circ_{n,k+2})=z, T^\star_n > k+2\right), \tag{6.75}$$

$$I_2 \equiv \textstyle\sum_{k=1}^{\ell^\circ_n-1} P_z\left(J_n(T^\circ_{n,k+2})=z, T^\star_n \le k+2\right). \tag{6.76}$$

In view of (3.3)-(3.4), $T^\circ_{n,i} = i$ for all $i \in \{0,\dots,T^\star_n - 1\}$. Hence

$$I_1 = \textstyle\sum_{k=1}^{\ell^\circ_n-1} P_z\left(J_n(k+2)=z, T^\star_n > k+2\right), \tag{6.77}$$

and since up to time $T^\star_n$ the transition probabilities of J_n are those of SRW,

$$I_1 \le \textstyle\sum_{k=1}^{\ell^\circ_n-1} P^{\mathsf{SRW}}_z\left(J^{\mathsf{SRW}}_n(k+2)=z\right) \le cn^{-2}, \tag{6.78}$$

where the last inequality is (6.58).

To Bound I_2 note that the event $\{T^\star_n \le k+2\}$ can be written as the disjoint union

$$\{T^\star_n \le k+2\} = \cup_{i\le k+2} \cup_{y\in\mathcal{V}^\star_n} \{T^\star_n = i, J_n(T^\star_n) = y\}. \tag{6.79}$$

Thus

$$I_2 = \textstyle\sum_{k=1}^{\ell^\circ_n-1} \sum_{i=1}^{k+2} \sum_{y\in\mathcal{V}^\star_n} P_z\left(J_n(T^\circ_{n,k+2})=z, T^\star_n = i, J_n(T^\star_n)=y\right). \tag{6.80}$$

As above note that $T^\circ_{n,i} = i$ for all $i \in \{0,\dots,T^\star_n - 1 = i-1\}$, that $T^\star_n = T^\circ_{n,i-1}+1$, and that in the time interval $\{0,\dots,T^\star_n\}$, J_n has the same transition probabilities as SRW. By this and the Markov property, the probability in (6.80) is equal to

$$P^{\mathsf{SRW}}_z\left(T^{\mathsf{SRW},\star}_n = i, J^{\mathsf{SRW}}_n(i) = y\right) P_y\left(J_n(T^\circ_{n,k+2-i})=z\right). \tag{6.81}$$

Consider now the last factor in (6.81). By construction $y \in \mathcal{V}^\star_n$. Hence, by (3.3),

$$\begin{aligned} P_y\left(J_n(T^\circ_{n,k+2-i})=z\right) &= P_y\left(J_n(T^\circ_{n,0})=z\right)\mathbb{1}_{\{k+2-i=0\}} \\ &\quad + \textstyle\sum_x P_y\left(J_n(T^\circ_{n,0})=x\right) P_x\left(J_n(T^\circ_{n,k+2-i})=z\right)\mathbb{1}_{\{k+2-i>0\}} \end{aligned} \tag{6.82}$$

where the sum is over x in $\partial\mathcal{V}^\star_n = \cup_{1\le l\le L^\star}\partial C^\star_{n,l}$. Note that $P_y\left(J_n(T^\circ_{n,0})=x\right)$ is nothing but the exit distribution from the set $C^\star_{n,l}$ containing y, so by Lemma 6.7, for all $y \in C^\star_{n,l}$,

$$\begin{aligned} P_y\left(J_n(T^\circ_{n,k+2-i})=z\right) &= (1+o(1))\big\{\mathcal{L}^\star_{n,l}(z)\mathbb{1}_{\{z\in\partial C^\star_{n,l}\}}\mathbb{1}_{\{k+2-i=0\}} \\ &\quad + \textstyle\sum_x \mathcal{L}^\star_{n,l}(x) P_x\left(J_n(T^\circ_{n,k+2-i})=z\right)\mathbb{1}_{\{k+2-i>0\}}\big\}. \end{aligned} \tag{6.83}$$

Observe that since by assumption z belongs to $I_n^\star$ then, by (2.9), $z \notin \partial\mathcal{V}_n^\star$. Thus the first term in the r.h.s. of (6.83) is zero. Now for indices i, k such that $k+2-i > 0$, let us rewrite the probability in the remaining term as $P_x\left(J_n(T^\circ_{n,k+2-i}) = z\right) = I_3(x) + I_4(x)$,

$$\begin{aligned} I_3(x) &\equiv P_x\left(J_n(T^\circ_{n,k+2-i}) = z, T_n^\star > k+2-i\right), \qquad (6.84)\\ I_4(x) \equiv A_x(k+2-i) &\equiv P_x\left(J_n(T^\circ_{n,k+2-i}) = z, T_n^\star \le k+2-i\right). \qquad (6.85)\end{aligned}$$

Proceeding as we did for I_1 to express $I_3(x)$ yields

$$I_3(x) = P_x^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(k+2-i) = z\right). \qquad (6.86)$$

Thus, up to a prefactor $1+o(1)$, the contribution to I_2 coming from the terms $I_3(x)$ is

$$\begin{aligned} K_1 \equiv \textstyle\sum_{k=1}^{\ell_n^\circ - 1}\sum_{i=1}^{k+1}\sum_{1\le l\le L^\star}\sum_{y\in C^\star_{n,l}}\big\{ & P_z^{\mathsf{SRW}}\left(T_n^{\mathsf{SRW},\star} = i, J_n^{\mathsf{SRW}}(i) = y\right)\\ & \textstyle\sum_{x\in\partial C^\star_{n,l}} \mathcal{L}^\star_{n,l}(x) P_x^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(k+2-i) = z\right)\big\}. \qquad (6.87)\end{aligned}$$

By a change of indices,

$$K_1 = \textstyle\sum_{1\le l\le L^\star}\sum_{j=1}^{\ell_n^\circ}\sum_{x\in\partial C^\star_{n,l}} \mathcal{L}^\star_{n,l}(x) P_x^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(j) = z\right) R_1(j), \qquad (6.88)$$

where

$$\begin{aligned} R_1(j) &\equiv \textstyle\sum_{m=1}^{\ell_n^\circ + 1 - j} P_z^{\mathsf{SRW}}\left(T_n^{\mathsf{SRW},\star} = m, J_n^{\mathsf{SRW}}(m) = y\right)\\ &\le P_z^{\mathsf{SRW}}\left(T_n^{\mathsf{SRW},\star} = T^{\mathsf{SRW}}(C^\star_{n,l}) \le \ell_n^\circ + 1 - j\right)\\ &\le P_z^{\mathsf{SRW}}\left(T_n^{\mathsf{SRW},\star} = T^{\mathsf{SRW}}(C^\star_{n,l}) \le \ell_n^\circ\right). \qquad (6.89)\end{aligned}$$

As observed earlier, $\mathrm{dist}(x,z) \ge 1$. Thus, by Lemma 6.14, and since $\sum_x \mathcal{L}^\star_{n,l}(x) = 1$,

$$\textstyle\sum_{j=1}^{\ell_n^\circ}\sum_{x\in\partial C^\star_{n,l}} \mathcal{L}^\star_{n,l}(x) P_x^{\mathsf{SRW}}\left(J_n^{\mathsf{SRW}}(j) = z\right) \le c'/n. \qquad (6.90)$$

Plugging (6.89) in (6.88) and using (6.90),

$$\begin{aligned} K_1 &\le (c'/n)\textstyle\sum_{1\le l\le L^\star} P_z^{\mathsf{SRW}}\left(T_n^{\mathsf{SRW},\star} = T^{\mathsf{SRW}}(C^\star_{n,l}) \le \ell_n^\circ\right)\\ &\le (c'/n) P_z^{\mathsf{SRW}}\left(T_n^{\mathsf{SRW},\star} \le \ell_n^\circ\right)\\ &\le (c'/n) P_z^{\mathsf{SRW}}\left(T^{\mathsf{SRW}}(V_n(\rho_n^\star)\setminus z) \le \ell_n^\circ + 1\right). \qquad (6.91)\end{aligned}$$

Note that by (6.1) and the assumption that $c_\star > 3$, $n^{c_\star} > \ell_n^\circ \log n$ for all n large enough. It thus follows from Lemma 6.12 that on Ω^{SRW}, for all but a finite number of indices n,

$$K_1 \le \Big(\frac{c'}{n}\Big)\Big(\frac{2 + C'/c_\star}{\log n}\Big). \qquad (6.92)$$

We assume from now on that $\omega \in \Omega^{\mathsf{SRW}}$. It remains to deal with the contribution to I_2 coming from the terms $I_4(x)$: in view of (6.85) this is given, up to a prefactor $1+o(1)$, by

$$\begin{aligned} K_2 \equiv \textstyle\sum_{k=1}^{\ell_n^\circ - 1}\sum_{i=1}^{k+1}\sum_{1\le l\le L^\star}\sum_{y\in C^\star_{n,l}}\big\{ & P_z^{\mathsf{SRW}}\left(T_n^{\mathsf{SRW},\star} = i, J_n^{\mathsf{SRW}}(i) = y\right)\\ & \textstyle\sum_{x\in\partial C^\star_{n,l}} \mathcal{L}^\star_{n,l}(x) A_x(k+2-i)\big\}. \qquad (6.93)\end{aligned}$$

Proceeding as in K_1 to rearrange the indices and using (6.89),

$$\begin{aligned} K_2 \le \textstyle\sum_{1\le l\le L^\star} & P_z^{\mathsf{SRW}}\left(T_n^{\mathsf{SRW},\star} = T^{\mathsf{SRW}}(C^\star_{n,l}) \le \ell_n^\circ + 1\right)\\ & \textstyle\sum_{x\in\partial C^\star_{n,l}} \mathcal{L}^\star_{n,l}(x)\sum_{j=1}^{\ell_n^\circ} A_x(j). \qquad (6.94)\end{aligned}$$

Summing up, we have established that $I_2 \leq (1+o(1))(K_1+K_2)$ with K_1 bounded by (6.92) and K_2 bounded by (6.94). But by (6.76) and (6.85),

$$I_2 = \sum_{k=1}^{\ell_n^\circ -1} A_z(k+2). \tag{6.95}$$

Hence

$$\sum_{k=1}^{\ell_n^\circ -1} A_z(k+2) \leq (c'/n)\alpha_n + \sum_{1\leq l\leq L^\star} \alpha_{n,l} \sum_{x\in\partial C^\star_{n,l}} \mathcal{L}^\star_{n,l}(x) \sum_{j=1}^{\ell_n^\circ} A_x(j), \tag{6.96}$$

where $\alpha_{n,l}\geq 0$, $\sum_{1\leq l\leq L^\star}\alpha_{n,l}=\alpha_n$, and $\alpha_n \leq c''/(\log n)$ for some constant $0<c''<\infty$.

The idea now simply is to iterate this relation. One easily checks, going through the preceeding arguments that, indeed, $\sum_{j=1}^{\ell_n^\circ} A_x(j)$ can be bounded in exactly the same way as I_2. Doing so and iterating κ times gives

$$I_2 \leq (c'/n)\sum_{m=1}^{\kappa}\alpha_n^m + \alpha_n^{\kappa-1}\sum_{1\leq l\leq L^\star}\alpha_{n,l}\sum_{x\in\partial C^\star_{n,l}}\mathcal{L}^\star_{n,l}(x)\sum_{j=1}^{\ell_n^\circ}A_x(j). \tag{6.97}$$

Since $A_x(j)\leq 1$, choosing $\kappa = \lfloor \log n\rfloor$ yields

$$I_2 \leq \frac{c'}{n}\frac{\alpha_n}{1-\alpha_n} + \alpha_n^\kappa \ell_n^\circ \leq \frac{c'}{n}\frac{\alpha_n}{1-\alpha_n} + o\Big(\frac{1}{n^2}\Big). \tag{6.98}$$

Inserting this bound and (6.78) in (6.74) we finally obtain

$$\sum_{k=1}^{\ell_n^\circ -1} P_z^\circ\left(J_n^\circ(k+2)=z\right) \leq cn^{-2}(1+o(1)) + c'n^{-1}\frac{\alpha_n}{1-\alpha_n} \leq \frac{C_\circ}{n\log n}, \tag{6.99}$$

for some constant $0<C_\circ<\infty$, and this is valid on Ω^{SRW} for all but a finite number of indices n. The proof of assertion (i) of Proposition 6.2 is complete. □

Proof of Proposition 6.2, (ii). The proof is a rerun of the proof of assertion (i) with only three minor changes that we now indicate. Let $1\leq l'\leq L^\star$ and $z,z'\in\partial C^\star_{n,l'}$ be given. As in (6.74) write

$$\sum_{k=1}^{\ell_n^\circ -1} P_{z'}^\circ\left(J_n^\circ(k)=z\right) = \sum_{k=1}^{\ell_n^\circ -1} P_{z'}\left(J_n(T^\circ_{n,k})=z\right) = I_1 + I_2 \tag{6.100}$$

where I_1 and I_2 are the analogues of (6.75) and (6.76), respectively. Arguing as in (6.77)-(6.78) to bound I_1, but using (6.58) of Lemma 6.13 if $z=z'$ and (6.59) of Lemma 6.14 if $z\neq z'$, we get that

$$I_1 \leq \sum_{k=1}^{\ell_n^\circ -1} P^{\mathsf{SRW}}_{z'}\left(J_n^{\mathsf{SRW}}(k)=z\right) \leq c''n^{-1} \tag{6.101}$$

for some constant $0<c''<\infty$. This is the most impacting change since, as we shall see, this term now becomes the leading one. The second change is in the treatment of (6.83) where the term $\mathcal{L}^\star_{n,l}(z)\mathbb{1}_{\{z\in\partial C^\star_{n,l}\}}\mathbb{1}_{\{k+2-i=0\}}$ is non zero whenever $l=l'$. In that case we deduce from (6.28) and (6.13) that $\mathcal{L}^\star_{n,l}(z)\leq (1+o(1))|C^\star_{n,l}|/|\partial C^\star_{n,l}|$, so that by (2.15) of Lemma 2.3,

$$\mathcal{L}^\star_{n,l}(z) \leq n^{-1}(1+o(1)) \tag{6.102}$$

on $\Omega^\star$, for all but a finite number of indices n. One then easily checks that the contribution to I_2 coming from this term can be bounded in the same way as the contribution coming from the terms $I_3(x)$, that is to say, as K_1. The third and last minor change is in the treatment of K_1. Notice that the sum over x in (6.87) contains the term $x=z$. For this term we bound $I_3(x)$ (see (6.86)) using (6.58). Clearly, this does not affect (6.90) and thus, the bound (6.92) (valid on Ω^{SRW}, for all but a finite number of indices n) is unchanged.

Poceeding from there on as in the proof of assertion (i) we finally get that on $\Omega^{\mathsf{SRW}}\cap\Omega^\star$, for all but a finite number of indices n,

$$\sum_{k=1}^{\ell_n^\circ -1} P_{z'}^\circ\left(J_n^\circ(k)=z\right) \leq c''n^{-1} + c'n^{-1}\frac{\alpha_n}{1-\alpha_n} + o(n^{-2}) \leq C'_\circ n^{-1} \tag{6.103}$$

for some constant $0 < C'_\circ < \infty$. The proof of assertion (ii) of Proposition 6.2 is done. □

6.5. **Hitting time at stationarity: proof of Proposition 6.4.** Let us first prove, that under the assumptions of Proposition 6.4, (6.9) holds for the continous time Markov chain J^*_n introduced in Subsection 6.3. For this we use results from [1]. Set $B \equiv A \cap I^\star_n$ and write $T^*(B) \equiv \inf\{t > 0 \mid J^*_n(t) \in B\}$. Then, by Theorem 3 and Lemma 2 of [1] we have,

$$P^*_{\pi^\circ_n}\left(T^*(B) > t\right) \geq \left(1 - \tau^\circ_n \tfrac{q(B,B^c)}{1-\pi^\circ_n(B)}\right) \exp\left(-t \tfrac{q(B,B^c)}{1-\pi^\circ_n(B)}\right), \quad t > 0, \tag{6.104}$$

where τ°_n is as in (6.30) and where $q(B, B^c) = \sum_{x\in B} \sum_{y\notin B} \pi^\circ_n(x) p^\circ_n(x, y) = \pi^\circ_n(B)$ as follows from (6.11) and the fact that $B \subseteq I^\star_n$. By Proposition 6.3,

$$\pi^\circ_n(B) = |B|/|\mathcal{V}^\circ_n| \leq |I^\star_n|/|\mathcal{V}^\circ_n| \leq n^{-c_\star}(1 + o(1)), \tag{6.105}$$

where we used (6.6) and (2.12) in the last inequality. From this and (6.30), we get that

$$P^*_{\pi^\circ_n}\left(T^*(B) > t\right) \geq \left(1-n^{-(c_\star-2)}(1+o(1))\right) \exp\left(-t\tfrac{|B|}{|\mathcal{V}^\circ_n|}(1+o(n^{-c_\star}))\right), \quad t > 0. \tag{6.106}$$

The idea then is that for s_n as in (6.45), $T^*(B) - T^\circ(B) = s_n(T^\circ(B)) - T^\circ(B)$, which by (6.48) should be small for $T^\circ(B)$ large. We thus need an a priori lower bound on $T^\circ(B)$. Now note that $B \subset V_n(\rho^\star_n)$ so that by Lemma 6.12 and (6.105), for all $l_n \leq n^{c_\star}/\log n$,

$$P^\circ_{\pi^\circ_n}(T^\circ(B) > l_n) \geq (1 - \pi^\circ_n(B)) \inf_{x\notin B} P^{\mathsf{SRW}}_x\left(T^{\mathsf{SRW}}(V_n(\rho^\star_n) \setminus x) \geq l_n\right) \geq 1 - \mathcal{O}(\tfrac{1}{\log n}).$$

From this bound, (6.48), and (6.106) we easily get that for $\mathcal{I}(\zeta)$ as in (6.47) and any $\zeta > 0$,

$$P^*_{\pi^\circ_n}\left(T^*(B) > t\right) \leq P^\circ_{\pi^\circ_n}\left(T^\circ(B) > t/(1+\zeta)\right) + 2e^{-l_n\mathcal{I}(\zeta)} + \mathcal{O}(\tfrac{1}{\log n}). \tag{6.107}$$

Taking e.g. $\zeta = 1/2$ and $l_n = n^2$ yields (6.9) and finishes the proof of Proposition 6.4.

7. Proof of Theorem 3.3 and of Theorem 3.1

The proofs of Theorem 3.3 and of Theorem 3.1 hinge upon the next two lemmata.

7.1. **Preparatory Lemmata.** Let $0 < \rho < 1$ and, for $V_n(\rho)$ defined in (2.1), set

$$C^\star_{n,l}(\rho) = \begin{cases} C^\star_{n,l} & \text{if } C^\star_{n,l} \cap V_n(\rho) \neq \emptyset, \\ \emptyset & \text{else.} \end{cases} \tag{7.1}$$

Lemma 7.1. *Assume that $c_\star > 2$. There exists a subset $\Omega_3 \subset \Omega$ with $\mathbb{P}(\Omega_3) = 1$ such that on Ω_3, for all but a finite number of indices n, for all $\rho^\star_n \leq \rho \leq 1 - 3\rho^\star_n$,*

$$\left|\cup_{1\leq l\leq L^\star} C^\star_{n,l}(\rho)\right| / |\mathcal{V}^\circ_n| \leq n^{-c_\star+1} 2^{-n\rho}(1 + o(1)), \tag{7.2}$$

and, for $m^\star_{n,l}(x)$ as in (6.13),

$$\sum_{1\leq l\leq L^\star} \sum_{x\in\partial C^\star_{n,l}(\rho)} \pi^\circ_n(x) m^\star_{n,l}(x) \leq n^{-c_\star+1} 2^{-n\rho}(1 + o(1)). \tag{7.3}$$

Lemma 7.2. *Assume that $c_\star > 2$. On $\Omega^\star$, for all but a finite number of indices n,*

$$\pi^\circ_n\left(\partial\left(\cup_{1\leq l\leq L^\star} C^\star_{n,l}\right)\right) \leq n^{-2(c_\star-1)}(1 + \mathcal{O}(n^{-(c_\star-1)})). \tag{7.4}$$

Proof of Lemma 7.2. By (6.5), $\pi^\circ_n\left(\partial(\cup_{1\leq l\leq L^\star} C^\star_{n,l})\right) \leq n|\cup_{1\leq l\leq L^\star} C^\star_{n,l}|/|\mathcal{V}^\circ_n|$. By (2.14) of Lemma 2.3 and (6.6) of Proposition 6.3, on $\Omega^\star$, for all but a finite number of indices n,

$$n|\cup_{1\leq l\leq L^\star} C^\star_{n,l}|/|\mathcal{V}^\circ_n| = n\left|V_n(\rho^\star_n) \setminus I^\star_n\right| / |\mathcal{V}^\circ_n| \leq nn^{-2c_\star+1}(1 + \mathcal{O}(n^{-(c_\star-1)})), \tag{7.5}$$

proving (7.4). □

Proof of Lemma 7.1. Set $k_n^\star \equiv \max_{2\le l\le L^\star} |C_{n,l}^\star(\rho)|$ and let

$$S_n(k) \equiv \sum_{l=2}^{L^\star} |C_{n,l}^\star(\rho)| \mathbb{1}_{\{|C_{n,l}^\star(\rho)|=k\}} \tag{7.6}$$

be the total number of vertices that belong to sets $C_{n,l}^\star(\rho)$ that have cardinality k. Note that by (2.3) and (2.11) of Lemma 2.3, on $\Omega^\star$, for large enough n,

$$k_n^\star \le n/((c_\star - 2)\log n). \tag{7.7}$$

Now, on the one hand,

$$|\cup_{1\le l\le L^\star} C_{n,l}^\star(\rho)|/|\mathcal{V}_n^\circ| = \tfrac{1}{|\mathcal{V}_n^\circ|}\sum_{l=2}^{L^\star} |C_{n,l}^\star(\rho)| = \tfrac{1}{|\mathcal{V}_n^\circ|}\sum_{k=2}^{k_n^\star} S_n(k). \tag{7.8}$$

On the other hand, by Proposition 6.3,

$$\begin{aligned} &\textstyle\sum_{1\le l\le L^\star}\sum_{x\in\partial C_{n,l}^\star(\rho)} \pi_n^\circ(x) m_{n,l}^\star(x) \\ =\ &\textstyle\sum_{k=2}^{k_n^\star}\sum_{l=2}^{L^\star} \mathbb{1}_{\{|C_{n,l}^\star(\rho)|=k\}} \frac{1}{|\mathcal{V}_n^\circ|}\sum_{x\in\partial C_{n,l}^\star(\rho)} m_{n,l}^\star(x) \le \tfrac{1}{|\mathcal{V}_n^\circ|}\sum_{k=2}^{k_n^\star} S_n(k), \end{aligned} \tag{7.9}$$

where we used in the final inequality that by (6.13),

$$\begin{aligned} \textstyle\sum_{x\in\partial C_{n,l}^\star(\rho)} m_{n,l}^\star(x) &= n^{-1}\textstyle\sum_{y\in C_{n,l}^\star(\rho)}\sum_{x\in\partial C_{n,l}^\star(\rho)} \mathbb{1}_{\{\mathrm{dist}(y,x)=1\}} &(7.10)\\ &= n^{-1}\textstyle\sum_{y\in C_{n,l}^\star(\rho)} |\partial C_{n,l}^\star(\rho)\cap\partial y| \le |C_{n,l}^\star(\rho)|, &(7.11)\end{aligned}$$

since $|\partial C_{n,l}^\star(\rho)\cap\partial y|\le n$. Let us now focus on the quantities $S_n(k)$, $2\le k\le k_n^\star$. We claim that if $c_\star > 2$ there exists a subset $\Omega^{\star\star}\subset\Omega$ with $\mathbb{P}(\Omega^{\star\star})=1$ such that, on $\Omega^{\star\star}$, for all but a finite number of indices n, for all $\rho_n^\star \le \rho \le 1-3\rho_n^\star$,

$$S_n(2) \le n^{-c_\star+1}2^{n(1-\rho)}(1+\mathcal{O}(n^{-(c_\star-1)})), \tag{7.12}$$

$$S_n(3) \le n^{-2(c_\star-1)}2^{n(1-\rho)}(1+\mathcal{O}(n^{-(c_\star-1)})), \tag{7.13}$$

and, for all $4\le k\le k_n^\star$,

$$S_n(k) \le n^{-1}n^{-c_\star+1}2^{n(1-\rho)}(1+\mathcal{O}(n^{-(c_\star-1)})). \tag{7.14}$$

We first prove (7.12). For this let us introduce the variables $\chi^\rho(x) \equiv \mathbb{1}_{\{w_n(x)\ge r_n(\rho)\}}$, $\chi_n^{\star,\rho}(x) \equiv \mathbb{1}_{\{r_n(\rho_n^\star)\le w_n(x)<r_n(\rho)\}}$, and $\chi_n(x)\equiv\mathbb{1}_{\{w_n(x)\ge r_n(\rho_n^\star)\}}$. They are Bernoulli r.v.'s with $\mathbb{P}(\chi^\rho(x)=1)=2^{-\rho n}$, $\mathbb{P}(\chi_n(x)=1)=n^{-c_\star}$, and $\mathbb{P}(\chi_n^{\star,\rho}(x)=1)=n^{-c_\star}-2^{-\rho n}$ respectively, that inherit the independence of the variables $(w_n(x), x\in\mathcal{V}_n)$. We then may write $S_n(2)=S_n^0(2)+S_n^1(2)$ where, for $\mathcal{G}_2$ as in (9.70) (see also the paragraph above (9.72)),

$$\begin{aligned} S_n^0(2) &\equiv \textstyle\sum_{\mathcal{C}=\{x,y\}\in\mathcal{G}_2}(Y_n(x,y)+Y_n(y,x)), &(7.15)\\ S_n^1(2) &\equiv \textstyle\sum_{\mathcal{C}=\{x,y\}\in\mathcal{G}_2} Z_n(x,y), &(7.16)\end{aligned}$$

where

$$\begin{aligned} Y_n(x,y) &\equiv \chi_n^\rho(x)\chi_n^{\star,\rho}(y)\textstyle\prod_{z\in(\partial x\cup\partial y)\setminus\{x,y\}}(1-\chi_n(z)), &(7.17)\\ Z_n(x,y) &\equiv \chi_n^\rho(x)\chi_n^\rho(y)\textstyle\prod_{z\in(\partial x\cup\partial y)\setminus\{x,y\}}(1-\chi_n(z)). &(7.18)\end{aligned}$$

To bound the sums (7.15) and (7.16) we proceed as in the proof of (9.105) (see (9.100)-(9.105)). Namely, we decompose $\mathcal{G}_2$ into $\mathcal{G}_2=\cup_{1\le j\le n}\cup_{1\le i\le 4}\mathcal{G}_2^{j,i}$, where the $\mathcal{G}_2^{j,i}$'s are defined in (9.101), and use Bennett's bound (9.103) to estimate the sum over each $\mathcal{G}_2^{j,i}$. Doing this we readily get that $\mathbb{E}S_n^0(2)=n(n^{-c_\star}-2^{-\rho n})2^{n(1-\rho)}(1-n^{-c_\star})^{2(n-1)}$ and

$$\mathbb{P}\left(|S_n^0(2)-\mathbb{E}S_n^0(2)|\ge 2n\sqrt{\mathbb{E}S_n^0(2)}\right)\le 4ne^{-n}. \tag{7.19}$$

Similarly, $\mathbb{E}S_n^1(2) = n2^{n(1-2\rho)}(1-n^{-c_\star})^{2(n-1)}$ and for all $\rho_n^\star \le \rho \le (1-4\rho_n^\star)/2$,

$$\mathbb{P}\left(|S_n^1(2) - \mathbb{E}S_n^1(2)| \ge 2n\sqrt{\mathbb{E}S_n^1(2)}\right) \le 4ne^{-n}. \tag{7.20}$$

For $\rho > (1-4\rho_n^\star)/2$ we simply use that by Tchebychev's first order order inequality,

$$\mathbb{P}\left(S_n^1(2) \ge 2^{-n\rho/2}\mathbb{E}S_n^0(2)\right) \le 2^{-n\rho/2}. \tag{7.21}$$

From the assumptions that $\rho_n^\star \le \rho \le 1 - 3\rho_n^\star$ and $c_\star > 1$ it then immediately follows that Eq. (7.12) holds true, and this with a probability larger than $1 - c_0 n e^{-c_1 n}$ for some constants $0 < c_0, c_1 < \infty$. Thus, by Borel-Cantelli Lemma, it holds on a subset of Ω of full measure, for all but a finite number of indices n.

One proves (7.13) in a similar way. We skip the (simple but lenghty) details.

When $4 \le k \le k_n^\star$ we do not need such a refined control on $S_n(k)$: we simply write

$$S_n(k) \le k \sum_{B\subset\mathcal{V}_n} \mathbb{1}_{\{\exists x\in B: \chi_n^\rho(x)\prod_{y\in B\setminus x}\chi_n^{\star,\rho}(y)=1\}} \prod_{z\in\partial B}(1-\chi_n(z)), \tag{7.22}$$

where the sum is over all subsets $B \subset \mathcal{V}_n$ such that $|B| = k$, and such that the graph $G(B)$ is connected. Since the number of such sets is bounded above by $(k-1)!n^{k-1}2^n$, $\mathbb{E}S_n(k) \le k!n^{k-1}n^{-c_\star(k-1)}2^{n(1-\rho)}$, and a first order order Tchebychev inequality yields

$$\mathbb{P}\left(S_n(k) \ge n^{-1}\mathbb{E}S_n^0(2)\right) \le k!nn^{-(c_\star-1)(k-2)}. \tag{7.23}$$

One easily checks that if $c_\star > 2$ then, for all $m \ge 3$ and all $K \le n$,

$$\sum_{k=m}^K k!n^{-(c_\star-1)(k-2)} \le (m!+1)n^{-(m-2)(c_\star-1)}. \tag{7.24}$$

Therefore $\mathbb{P}\left(\cup_{4\le k\le k_n^\star}\{S_n(k) \ge n^{-1}\mathbb{E}S_n^0(2)\}\right) \le 25n^{-2(c_\star-1)+1}$, which is summable when $c_\star > 2$. By Borel-Cantelli Lemma we conclude that on a subset of Ω of full measure, for all but a finite number of indices n, (7.14) holds true for all $4 \le k \le k_n^\star$. This concludes the proof of the claim (7.12)-(7.14).

Now, by (7.12)-(7.14) and (6.6), on $\Omega_3 \equiv \Omega^\star \cap \Omega^{\star\star}$, for all large enough n,

$$\tfrac{1}{|\mathcal{V}_n^\circ|}\sum_{k=2}^{k_n^\star} S_n(k) \le (1+o(1))(1+(k_n^\star/n))n^{-c_\star+1}2^{-n\rho} = n^{-c_\star+1}2^{-n\rho}(1+o(1)), \tag{7.25}$$

where the last equality follows from (7.7). Inserting (7.25) in (7.8) and in (7.9) yields (7.2) and (7.3), respectively. The proof of Lemma 7.1 is done. □

7.2. **Elementary properties of the chains $J_n^\dagger$ and J_n°.** For easy reference we gather here a few elementary properties of the chains $J_n^\dagger$ and J_n°. We state them without proof: recalling that $J_n^\circ(i) \equiv J_n(T_{n,i}^\circ)$ and $J_n^\dagger(i) \equiv J_n(T_{n,i}^\dagger)$ they are immediate consequences from the definitions of the sequences $\left(T_{n,j}^\dagger\right)$ and $\left(T_{n,j}^\circ\right)$ (see (3.3)-(3.10)).

Lemma 7.3. *To each $j \ge 0$ there corresponds a unique $i \le j$ such that:*
(i) $J_n^\dagger(j) \notin \mathcal{V}_n^\circ \Leftrightarrow T_{n,j-1}^\dagger = T_{n,i-1}^\circ < T_{n,j}^\dagger = T_{n,i-1}^\circ + 1 < T_{n,j+1}^\dagger = T_{n,i}^\circ$,
(ii) $J_n^\dagger(j) \in \mathcal{V}_n^\circ \Leftrightarrow T_{n,j}^\dagger = T_{n,i}^\circ$.

From Lemma 7.3, (i), we derive two descriptions of the event $\{J_n^\dagger(j) \in C_{n,l}^\star\}$, $j > 0$, $1 \le l \le L^\star$. The first consists in saying that a visit of $J_n^\dagger$ to $C_{n,l}^\star$ must be immediately preceded and followed by a visit to $\partial C_{n,l}^\star$.

Corollary 7.4. $\{J_n^\dagger(j) \in C_{n,l}^\star\} = \{J_n^\dagger(j-1) \in \partial C_{n,l}^\star, J_n^\dagger(j) \in C_{n,l}^\star, J_n^\dagger(j+1) \in \partial C_{n,l}^\star\}$.

The second expresses the fact that when $J_n^\dagger(j)$ enters $C_{n,l}^\star$, $J_n^\circ(i)$ straddles over it.

Corollary 7.5. *To each $j \geq 0$ there corresponds a unique $i \leq j$ such that*

$$\{J_n(T^\dagger_{n,j}) \in C^\star_{n,l}\} = \{J_n(T^\circ_{n,i-1}) \in \partial C^\star_{n,l}, J_n(T^\circ_{n,i}) \in \partial C^\star_{n,l}\}. \tag{7.26}$$

Note finally that by Lemma 7.3, (ii), the chain $J^\dagger_n$ observed only when it visits $\mathcal{V}^\circ_n$ is nothing but the chain J°_n itself:

Corollary 7.6. $\{J^\dagger_n(j) : \exists i > 0 \text{ s.t. } T^\dagger_{n,j} = T^\circ_{n,i}, j \geq 0\} \stackrel{d}{=} \{J^\circ_n(i), i \geq 0\}$.

7.3. **Proof of Theorem 3.1.** Theorem 3.1 is a rough estimate. By (3.15),

$$0 \leq k^\dagger_n(t) - k^\circ_n(t) = \textstyle\sum_{j=0}^{k^\dagger_n(t)-1} \mathbb{1}_{\{J^\dagger_n(j) \notin \mathcal{V}^\circ_n\}}. \tag{7.27}$$

We now want to replace the chain $J^\dagger_n$ and the quantity $k^\dagger_n(t)$ in the right hand side of (7.27) by, respectively, J°_n and $k^\circ_n(t)$. Note that by Corollary 7.4, for each $j \geq 1$,

$$\{J^\dagger_n(j) \notin \mathcal{V}^\circ_n\} = \cup_{1 \leq l \leq L^\star} \{J^\dagger_n(j) \in C^\star_{n,l}\} \subseteq \{J^\dagger_n(j-1) \in \partial(\cup_{1 \leq l \leq L^\star} C^\star_{n,l})\}. \tag{7.28}$$

From this and the fact that $J^\dagger_n(0) = J^\circ_n(0) \in \mathcal{V}^\circ_n$ (indeed $J^\dagger_n$ starts in π°_n), we deduce that,

$$\begin{aligned} \textstyle\sum_{j=0}^{k^\dagger_n(t)-1} \mathbb{1}_{\{J^\dagger_n(j) \notin \mathcal{V}^\circ_n\}} &\leq \textstyle\sum_{j=1}^{k^\dagger_n(t)-1} \mathbb{1}_{\{J^\dagger_n(j-1) \in \partial(\cup_{1 \leq l \leq L^\star} C^\star_{n,l})\}} &(7.29)\\ &\stackrel{d}{=} \textstyle\sum_{i=1}^{k^\circ_n(t)-1} \mathbb{1}_{\{J^\circ_n(i-1) \in \partial(\cup_{1 \leq l \leq L^\star} C^\star_{n,l})\}}, &(7.30)\end{aligned}$$

where the last equality follows from Corollary 7.6 and the definition of $k^\circ_n(t)$ (see (3.19)). It remains to bound the last sum in (7.30). Since $k^\circ_n(t) = \lfloor a_n t \rfloor$ is deterministic, a first order Tchebychev inequality entails that for all $c_\circ > 0$,

$$P_{\pi^\circ_n}\left(\textstyle\sum_{i=1}^{\lfloor a_n t \rfloor - 1} \mathbb{1}_{\{J^\circ_n(i-1) \in \partial(\cup_{1 \leq l \leq L^\star} C^\star_{n,l})\}} \geq n^{-c_\circ} \lfloor a_n t \rfloor\right) \leq n^{c_\circ} \pi^\circ_n\left(\partial(\cup_{1 \leq l \leq L^\star} C^\star_{n,l})\right).$$

Inserting (7.4) of Lemma 7.2 in the right hand side above, and combining the resulting bound with (7.27) and (7.30), we get that on $\Omega^\star$, for all but a finite number of indices n,

$$P_{\pi^\circ_n}\left(k^\dagger_n(t) \geq k^\circ_n(t)\left(1 + n^{-c_\circ}\right)\right) \leq n^{-2(c_\star - 1) + c_\circ}(1 + \mathcal{O}(n^{-(c_\star - 1)})). \tag{7.31}$$

This readily implies the claim of Theorem 3.1.

7.4. **Proof of Theorem 3.3.** By definition of the Skorohod topology on $D[0, \infty)$, it is enough to show this result with ρ_∞ replaced by ρ_r, the Skorohod metric on $D[0, r]$, for $r > 0$ arbitrary. Choosing $r = 1$ for convenience we get

$$\mathcal{P}_{\pi^\circ_n}\left(\rho_1\big(S_n(\cdot), S^\circ_n(\cdot)\big) > \epsilon\right) \leq \mathcal{P}_{\pi^\circ_n}\big(\sup_{0 \leq t \leq 1} \widehat{S}_n(t) > \epsilon\big). \tag{7.32}$$

Theorem 3.3 then is an immediate consequence of the lemma below.

Lemma 7.7. *Assume that $c_\star > 2$ and that $\beta > \beta_c(\varepsilon)$. Then $\mathbb{P}$-almost surely, for all $\epsilon > 0$,*

$$\limsup_{n \to \infty} \mathcal{P}_{\pi^\circ_n}\big(\sup_{0 \leq t \leq 1} \widehat{S}_n(t) > \epsilon\big) = 0. \tag{7.33}$$

Proof of Lemma 7.7. Since $\widehat{S}_n$ is nondecreasing

$$\mathcal{P}_{\pi^\circ_n}\big(\sup_{0 \leq t \leq 1} \widehat{S}_n(t) > \epsilon\big) \leq \mathcal{P}_{\pi^\circ_n}\big(\widehat{S}_n(1) > \epsilon\big). \tag{7.34}$$

Introducing the event

$$\mathcal{A} \equiv \left\{\forall_{0 \leq j \leq k^\dagger_n(1)-1} \forall_{1 \leq l \leq L^\star} \widehat{\Lambda}^\dagger_n(J^\dagger_n(j)) \mathbb{1}_{\{J^\dagger_n(j) \in C^\star_{n,l}\}} \leq n \bar{\varrho}_{n,l}(0)\right\} \tag{7.35}$$

we have, by Corollary 4.2, that on $\Omega_0 \cap \Omega^\star$, for all but a finite number of indices n,

$$\mathcal{P}_{\pi^\circ_n}\big(\widehat{S}_n(1) > \epsilon\big) \leq e^{-n} + n^{-2(c_\star - 1) + c_\circ} + \mathcal{P}_{\pi^\circ_n}\big(\widehat{S}_n(1) > \epsilon, \mathcal{A}\big). \tag{7.36}$$

where $c_\circ > 0$ is arbitrary. From the definitions (3.27) and (3.12) of $\widehat{S}_n$ and $\Lambda_n^\dagger(J_n^\dagger(i))$, and since $\Lambda_n^\dagger(J_n^\dagger(i))$ is non zero if and only if $J_n^\dagger(i) \in \cup_{1\le l\le L^\star} C_{n,l}^\star$, we see that on $\mathcal{A}$,

$$\begin{aligned}\widehat{S}_n(1) &= c_n^{-1}\textstyle\sum_{l=1}^{L^\star}\sum_{j=1}^{k_n^\dagger(1)-1}\widehat{\Lambda}_n^\dagger(J_n^\dagger(j))\mathbb{1}_{\{J_n^\dagger(j)\in C_{n,l}^\star\}} &(7.37)\\ &\le c_n^{-1}\textstyle\sum_{l=1}^{L^\star}\sum_{j=1}^{k_n^\dagger(1)-1} n\bar{\varrho}_{n,l}(0)\mathbb{1}_{\{J_n^\dagger(j)\in C_{n,l}^\star\}}. &(7.38)\end{aligned}$$

Therefore,

$$\mathcal{P}_{\pi_n^\circ}\big(\widehat{S}_n(1)>\epsilon,\mathcal{A}\big)\le \mathcal{P}_{\pi_n^\circ}\big(nc_n^{-1}\textstyle\sum_{l=1}^{L^\star}\bar{\varrho}_{n,l}(0)\sum_{j=1}^{k_n^\dagger(1)-1}\mathbb{1}_{\{J_n^\dagger(j)\in C_{n,l}^\star\}}>\epsilon\big). \tag{7.39}$$

The problem we still face is that the quantity $\bar{\varrho}_{n,l}(0)$ appearing in (7.39) can be very large compared to c_n. However, sets $C_{n,l}^\star$ such that this happens will typically not be visited. More precisely, for $C_{n,l}^\star(\rho)$ as in (7.1), one may choose the parameter $0<\rho<1$ in a such a way that the event

$$\widetilde{\mathcal{A}} \equiv \Big\{\forall_{1\le j\le k_n^\dagger(1)-1} J_n^\dagger(j)\notin\big(\cup_{1\le l\le L^\star} C_{n,l}^\star(\rho)\big)\Big\}, \tag{7.40}$$

has probability close to one. Indeed

$$\begin{aligned}\mathcal{P}_{\pi_n^\circ}\big(\widetilde{\mathcal{A}}^c\big)=P_{\pi_n^\circ}\big(\widetilde{\mathcal{A}}^c\big) &= \textstyle\sum_{1\le l\le L^\star} E_{\pi_n^\circ}\sum_{j=0}^{k_n^\dagger(1)-1}\mathbb{1}_{\{J_n^\dagger(j)\in C_{n,l}^\star\}} &(7.41)\\ &= \textstyle\sum_{1\le l\le L^\star} E_{\pi_n^\circ}\sum_{i=1}^{k_n^\circ(1)-1}\mathbb{1}_{\{J_n^\circ(i-1)\in\partial C_{n,l}^\star, J_n^\circ(i)\in\partial C_{n,l}^\star\}} &(7.42)\end{aligned}$$

where (7.42) follows from Corollary 7.5. Note that for all $x\in\partial C_{n,l}^\star$,

$$P_{\pi_n^\circ}\big(J_n^\circ(i-1)=x, J_n^\circ(i)\in\partial C_{n,l}^\star\big)=\pi_n^\circ(x)P_x(J_n^\circ(1)\in\partial C_{n,l}^\star)=\pi_n^\circ(x)m_{n,l}^\star(x), \tag{7.43}$$

where $m_{n,l}^\star(x)$ is defined in (6.13). Inserting this in (7.42), it follows from (7.3) of Lemma 7.1 that on Ω_3, for all but a finite number of indices n,

$$\begin{aligned}\mathcal{P}_{\pi_n^\circ}\big(\widetilde{\mathcal{A}}^c\big) &\le k_n^\circ(t)\textstyle\sum_{1\le l\le L^\star}\sum_{x\in\partial C_{n,l}^\star(\rho)}\pi_n^\circ(x)m_{n,l}^\star(x) &(7.44)\\ &\le k_n^\circ(t)n^{-c_\star+1}2^{-n\rho}(1+o(1))=n2^{-n\rho}2^{n\varepsilon_n-n\rho_n^\star}(1+o(1)). &(7.45)\end{aligned}$$

where we wrote $\varepsilon_n\equiv\frac{\log a_n}{n\log 2}$; thus by 1.19, $\lim_{n\to\infty}\varepsilon_n=\varepsilon$, $0<\varepsilon<1$. Assume from now on that $\omega\in\Omega_3$ and take $\rho\equiv\varepsilon_n-\rho_n^\star/2$. Then

$$\begin{aligned}\mathcal{P}_{\pi_n^\circ}\big(\widehat{S}_n(1)>\epsilon,\mathcal{A}\big) &\le \mathcal{P}_{\pi_n^\circ}\big(\widetilde{\mathcal{A}}^c\big)+\mathcal{P}_{\pi_n^\circ}\big(\widehat{S}_n(1)>\epsilon,\mathcal{A},\widetilde{\mathcal{A}}\big)\\ &\le n2^{-n\rho_n^\star/2}(1+o(1))+\mathcal{P}_{\pi_n^\circ}\big(\widehat{\mathcal{A}}\big), &(7.46)\end{aligned}$$

where, recalling from (2.1) that $V_n(\varepsilon_n-\rho_n^\star/2)=\{x\in\mathcal{V}_n\mid w_n(x)\ge r_n(\varepsilon_n-\rho_n^\star/2)\}$,

$$\widehat{\mathcal{A}}\equiv\Big\{nc_n^{-1}\textstyle\sum_{1\le l\le L^\star: C_{n,l}^\star\cap V_n(\varepsilon_n-\rho_n^\star/2)=\emptyset}\bar{\varrho}_{n,l}(0)\sum_{j=1}^{k_n^\dagger(1)-1}\mathbb{1}_{\{J_n^\dagger(j)\in C_{n,l}^\star\}}>\epsilon\Big\}. \tag{7.47}$$

Again, we wish to express this event in terms of the chain J_n° and the quantity $k_n^\circ(t)$ rather than $J_n^\dagger$ and $k_n^\dagger(t)$. For this note that by Corollary 7.5, Corollary 7.6, the definition (3.15) of $k_n^\dagger(t)$ and the definition (3.19) of $k_n^\circ(t)$, for each $1\le l\le L^\star$,

$$\textstyle\sum_{j=1}^{k_n^\dagger(1)-1}\mathbb{1}_{\{J_n^\dagger(j)\in C_{n,l}^\star\}}\stackrel{d}{=}\sum_{i=1}^{k_n^\circ(1)-1}\mathbb{1}_{\{J_n^\circ(i-1)\in\partial C_{n,l}^\star, J_n^\circ(i)\in\partial C_{n,l}^\star\}}. \tag{7.48}$$

Then, by Tchebychev inequality, (7.48), and (7.43),

$$\mathcal{P}_{\pi_n^\circ}\big(\widehat{\mathcal{A}}\big) \le \frac{n\lfloor a_n\rfloor}{\epsilon c_n}\sum_{\substack{1\le l\le L^\star:\\ C_{n,l}^\star\cap V_n(\varepsilon_n-\rho_n^\star/2)=\emptyset}}\max_{x\in C_{n,l}^\star}w_n(x)\sum_{x\in\partial C_{n,l}^\star}\pi_n^\circ(x)m_{n,l}^\star(x). \tag{7.49}$$

We next decompose the sum in (7.49) according to the size of $\max_{x\in C^\star_{n,l}} w_n(x)$: given $K>0$ to be chosen later define, for $0\le k\le K$,

$$\mathcal{I}_k \equiv \Big\{1\le l\le L^\star \mid r_n\big(\varepsilon_n-\tfrac{k+2}{2}\rho_n^\star\big)\le \max_{x\in C^\star_{n,l}} w_n(x)\le r_n\big(\varepsilon_n-\tfrac{k+1}{2}\rho_n^\star\big)\Big\}. \quad (7.50)$$

By this and the choices of a_n and c_n from Theoreom 1.1, (7.49) becomes

$$\mathcal{P}_{\pi_n^\circ}\big(\widehat{\mathcal{A}}\big)\le \epsilon^{-1}n\left(\textstyle\sum_{0\le k\le K} Q_{n,k}+R_n\right), \quad (7.51)$$

where

$$\begin{aligned} Q_{n,k} &= 2^{n\varepsilon_n} r_n^{-1}(\varepsilon_n) r_n\big(\varepsilon_n-\tfrac{k+1}{2}\rho_n^\star\big)\textstyle\sum_{l\in\mathcal{I}_k}\sum_{x\in\partial C^\star_{n,l}}\pi_n^\circ(x)m^\star_{n,l}(x), && (7.52)\\ R_n &= 2^{n\varepsilon_n} r_n^{-1}(\varepsilon_n) r_n\big(\varepsilon_n-\tfrac{K+2}{2}\rho_n^\star\big)\textstyle\sum_{1\le l\le L^\star}\sum_{x\in\partial C^\star_{n,l}}\pi_n^\circ(x)m^\star_{n,l}(x). && (7.53)\end{aligned}$$

Now,

$$\begin{aligned}\textstyle\sum_{l\in\mathcal{I}_k}\sum_{x\in\partial C^\star_{n,l}}\pi_n^\circ(x)m^\star_{n,l}(x) &\le \textstyle\sum_{1\le l\le L^\star}\sum_{x\in\partial C^\star_{n,l}\left(\varepsilon_n-\frac{k+2}{2}\rho_n^\star\right)}\pi_n^\circ(x)m^\star_{n,l}(x)\\ &\le n^{-c_\star+1}2^{-n\left(\varepsilon_n-\frac{k+2}{2}\rho_n^\star\right)}(1+o(1)) && (7.54)\end{aligned}$$

where the last inequality is (7.3) of Lemma 7.1. Inserting (7.54) in (7.52),

$$Q_{n,k}\le n2^{\frac{kn}{2}\rho_n^\star} r_n^{-1}(\varepsilon_n) r_n\big(\varepsilon_n-\tfrac{k+1}{2}\rho_n^\star\big). \quad (7.55)$$

Using (2.19), the bound $\sqrt{1-x}-1\le -\frac12 x(1+\frac14 x)$, $0<x<1$, and the assumption that $\beta>\beta_c(\varepsilon)$, so that $\alpha(\varepsilon_n)\equiv\beta_c(\varepsilon_n)/\beta<1$ for large enough n, it follows from (7.55) that

$$Q_{n,k}\le c_0 n2^{-n\rho_n^\star/\alpha(\varepsilon_n)}2^{-n\rho_n^\star(1/\alpha(\varepsilon_n)-1)\frac{k}{2}} \quad (7.56)$$

for some constant $0<c_0\equiv c_0(\varepsilon_n,\beta)<\infty$. Similarly, by (7.3) with $\rho=\rho_n^\star$,

$$\textstyle\sum_{1\le l\le L^\star}\sum_{x\in\partial C^\star_{n,l}}\pi_n^\circ(x)m^\star_{n,l}(x)\le n^{-2c_\star+1}(1+o(1)) \quad (7.57)$$

and

$$R_n\le n2^{n\varepsilon_n-2n\rho_n^\star} r_n^{-1}(\varepsilon_n) r_n\big(\varepsilon_n-\tfrac{K+2}{2}\rho_n^\star\big). \quad (7.58)$$

Now choose $K=\big\lceil 2\varepsilon_n\big(1-\frac{1}{16}\big)/\rho_n^\star\big\rceil$. Then $\frac{K+2}{2}\rho_n^\star\ge\varepsilon_n\big(1-\frac{1}{16}\big)$ and, using (2.19),

$$R_n\le n2^{n\varepsilon_n-2n\rho_n^\star} r_n^{-1}(\varepsilon_n) r_n\big(\varepsilon_n/16\big)\le n2^{-n\varepsilon_n/4-2n\rho_n^\star} \quad (7.59)$$

for all $\beta>\beta_c(\varepsilon_n)$. Inserting (7.56) and (7.59) in (7.51),

$$\mathcal{P}_{\pi_n^\circ}\big(\widehat{\mathcal{A}}\big)\le 2\epsilon^{-1}\left(c_0 n^{2-c_\star/\alpha(\varepsilon_n)}+n^{-2(c_\star-1)}2^{-n\varepsilon_n/4}\right) \quad (7.60)$$

for all n large enough. Finally, combining (7.36), (7.46), and (7.60), we obtain that for all $\beta>\beta_c(\varepsilon)$, on $\Omega_0\cap\Omega^\star\cap\Omega_3$, for all but a finite number of indices n,

$$\mathcal{P}_{\pi_n^\circ}\big(\widehat{S}_n(1)>\epsilon\big)\le e^{-n}+n^{-2(c_\star-1)+c_\circ}+2n^{-(c_\star-2)/2}+2\epsilon^{-1}\left(c_0 n^{2-c_\star/\alpha(\varepsilon_n)}+n^{-2(c_\star-1)}2^{-n\varepsilon_n/4}\right)$$

for all $\epsilon>0$, where $c_\circ>0$ is arbitrarily small, and where $\lim_{n\to\infty}\varepsilon_n=\varepsilon$, $0<\varepsilon<1$. This yields the claim of Lemma 7.7 since by assumption $c_\star>2$. □

The proof of Theorem 3.3 is now complete.

8. Convergence of the front end clock process: Proof of Theorem 3.2

The proofs of Theorem 3.2 and Theorem 3.4 rely on a method developped by Durrett and Resnick [22] that provides sufficient conditions for partial sum processes to converge to Lévy processes. We use their results in a specialized form suitable for our applications which is taken from [26], where this method was first applied to the study of clock processes in random environment; see also [14] where it was implemented in more generality.

8.1. **A convergence theorem for FECP.** Consider the rescaled front end clock process (3.26),

$$S_n^\circ(t) = c_n^{-1}\widetilde{S}_n^\circ(\lfloor a_n t\rfloor), \quad t \geq 0. \tag{8.1}$$

Theorem 8.1 below is the corner stone of the proof of Theorem 3.2. It deduces convergence of S_n° to a subordinator from a set of four conditions which we now formulate. Note that these conditions refer to given sequences of numbers a_n and c_n, as well as a given realization of the random environment. For $t > 0$ and $u > 0$ define

$$h_n^u(y) = \sum_{x\in\mathcal{V}_n^\circ} p_n^\circ(y,x)\exp\{-uc_n\lambda_n(x)\}\,,\ y \in \mathcal{V}_n^\circ, \tag{8.2}$$

and, recalling the notation $k_n^\circ(t) = \lfloor a_n t\rfloor$,

$$\nu_n^{J_n^\circ,t}(u,\infty) = \sum_{j=0}^{k_n^\circ(t)-1} h_n^u(J_n^\circ(j)), \tag{8.3}$$

$$\sigma_n^{J_n^\circ,t}(u,\infty) = \sum_{j=0}^{k_n^\circ(t)-1} [h_n^u(J_n^\circ(j))]^2\,. \tag{8.4}$$

Condition (C0). For all $v > 0$,

$$\textstyle\sum_{x\in\mathcal{V}_n^\circ} \pi_n^\circ(x)e^{-vc_n\lambda_n(x)} = o(1)\,. \tag{8.5}$$

Condition (C1). There exists a σ-finite measure ν° on $(0,\infty)$ satisfying $\int_0^\infty (1\wedge u)\nu^\circ(du) < \infty$ such that, for all $t > 0$ and all $u > 0$,

$$P_{\pi_n^\circ}^\circ\left(\left|\nu_n^{J_n^\circ,t}(u,\infty) - t\nu^\circ(u,\infty)\right| < \epsilon\right) = 1 - o(1)\,, \quad \forall\epsilon > 0\,. \tag{8.6}$$

Condition (C2). For all $u > 0$ and all $t > 0$,

$$P_{\pi_n^\circ}^\circ\left(\sigma_n^{J_n^\circ,t}(u,\infty) < \epsilon\right) = 1 - o(1)\,, \quad \forall\epsilon > 0\,. \tag{8.7}$$

Condition (C3). For all $t > 0$,

$$\lim_{\epsilon\downarrow 0}\limsup_{n\uparrow\infty} k_n^\circ(t)\mathcal{E}_{\pi_n^\circ}^\circ \mathbb{1}_{\{\lambda_n^{-1}(J_n^\circ(0))e_0^\circ \leq c_n\epsilon\}} c_n^{-1}\lambda_n^{-1}(J_n^\circ(0))e_0^\circ = 0. \tag{8.8}$$

Theorem 8.1. *Let the initial distribution of J_n° be its invariant measure π_n°. For all sequences a_n and c_n for which Conditions (C0), (C1), (C2), and (C3) are verified $\mathbb{P}$-almost surely,*

$$S_n^\circ \Rightarrow_{J_1} S_\infty^\circ \tag{8.9}$$

$\mathbb{P}$-almost surely, where S_∞° is the Lévy subordinator with zero drift and Lévy measure ν°.

Proof. This is a restatement of Theorem 1.2 of [14] specialized to the case where θ_n, the "bloc length", is equal to one. (Theorem 1.2 of [14] is itself a generalization of Theorem 1.1 of [26] with a more workable Condition (C3).) □

To verify the conditions of Theorem 8.1 we follow a by now well established two-step strategy that was first proposed in [25], and was used later in [14]. The first step consists in using the mixing property and mean local time estimates of Proposition 6.1 and Proposition 6.2, respectively, to prove an almost sure ergodic theorem for the quantities (8.3) and (8.4). This is done in Subsection 8.2 (see Theorem 8.2). It then enables us to reduce Conditions (C1) and (C2) of Theorem 8.1 to laws of large numbers in the random environment. This second step is carried out in Subsection 8.3 (see Proposition 8.4). The proof is completed in Subsection 8.4.

8.2. **An ergodic theorem for FECP.** Let $\pi_n^{J_n^\circ,t}(x)$ denote the average number of visits of J_n° to x during the first $k_n^\circ(t)$ steps,

$$\pi_n^{J_n^\circ,t}(x) = (k_n^\circ(t))^{-1} \sum_{j=0}^{k_n^\circ(t)-1} \mathbb{1}_{\{J_n^\circ(j)=x\}}\,, \quad x \in \mathcal{V}_n^\circ\,. \tag{8.10}$$

Then (8.3) and (8.4) can be rewritten as

$$\nu_n^{J_n^\circ,t}(u,\infty) \;=\; k_n^\circ(t) \sum_{y\in\mathcal{V}_n^\circ} \pi_n^{J_n^\circ,t}(y) h_n^u(y), \tag{8.11}$$

$$\sigma_n^{J_n^\circ,t}(u,\infty) \;=\; k_n^\circ(t) \sum_{y\in\mathcal{V}_n^\circ} \pi_n^{J_n^\circ,t}(y) \left[h_n^u(y)\right]^2. \tag{8.12}$$

One readily sees, using reversibility, that

$$E_{\pi_n^\circ}^\circ\left[\nu_n^{J_n^\circ,t}(u,\infty)\right] = k_n^\circ(t) \sum_{x\in\mathcal{V}_n^\circ} \pi_n^\circ(x) h_n^u(x) = (k_n^\circ(t)/a_n)\nu_n^\circ(u,\infty), \tag{8.13}$$

$$E_{\pi_n^\circ}^\circ\left[\sigma_n^{J_n^\circ,t}(u,\infty)\right] = k_n^\circ(t) \sum_{x\in\mathcal{V}_n^\circ} \pi_n^\circ(x) \left[h_n^u(x)\right]^2 = (k_n^\circ(t)/a_n)\sigma_n^\circ(u,\infty), \tag{8.14}$$

where

$$\nu_n^\circ(u,\infty) = \frac{a_n}{|\mathcal{V}_n^\circ|} \sum_{x\in\mathcal{V}_n^\circ} e^{-uc_n\lambda_n(x)}, \tag{8.15}$$

$$\sigma_n^\circ(u,\infty) = \frac{a_n}{|\mathcal{V}_n^\circ|} \sum_{x\in\mathcal{V}_n^\circ} \sum_{x'\in\mathcal{V}_n^\circ} p_n^{\circ,2}(x,x') e^{-uc_n(\lambda_n(x)+\lambda_n(x'))}. \tag{8.16}$$

Here $p_n^{\circ,2}(\cdot,\cdot)$ denotes the 2-steps transition probabilities of J_n°. Note that since $H(x) = 0$ for all $x \in \mathcal{V}_n^\circ \setminus I_n^\star$, where $I_n^\star$ is the set of isolated vertices in the partition (2.9), we have

$$\lambda_n(x) = \begin{cases} e^{\beta H_n(x)}, & \text{if } x \in I_n^\star, \\ 1, & \text{if } x \in \mathcal{V}_n^\circ \setminus I_n^\star. \end{cases} \tag{8.17}$$

Theorem 8.2. *Assume that $c_\star > 3$. Let $\rho_n^\circ > 0$ be a decreasing sequence satisfying $\rho_n^\circ \downarrow 0$ as $n \uparrow \infty$. There exists a sequence of subsets $\Omega_n^{EG} \subset \Omega$ with $\mathbb{P}\left[(\Omega_n^{EG})^c\right] < \ell_n^\circ/(\rho_n^\circ a_n)$, and such that on Ω_n^{EG} the following holds for all large enough n: for all $t > 0$, all $u > 0$, and all $\epsilon > 0$,*

$$P_{\pi_n^\circ}^\circ\left(\left|\nu_n^{J_n^\circ,t}(u,\infty) - (k_n^\circ(t)/a_n)\nu_n^\circ(u,\infty)\right| \geq \epsilon\right) \leq \epsilon^{-2}[C_1 t\Theta_{n,1}(u) + t^2\Theta_{n,2}(u)] \tag{8.18}$$

for some constant $0 < C_1 < \infty$, where

$$\Theta_{n,1}(u) \equiv \ell_n^\circ e^{-uc_n}[1 + \nu_n^\circ(u,\infty)] + \sigma_n^\circ(u,\infty) + \frac{\nu_n^\circ(2u,\infty)}{n\log n} + \rho_n^\circ\left[\mathbb{E}\nu_n^\circ(u,\infty)\right]^2, \tag{8.19}$$

$$\Theta_{n,2}(u) \equiv 2^{-n}\left[\nu_n^\circ(u,\infty)\right]^2. \tag{8.20}$$

Moreover, for all $t > 0$, all $u > 0$, and all $\epsilon' > 0$,

$$P_{\pi_n^\circ}^\circ\left(\sigma_n^{J_n^\circ,t}(u,\infty) \geq \epsilon'\right) \leq \frac{t}{\epsilon'}(1 + o(1))\sigma_n^\circ(u,\infty). \tag{8.21}$$

Proof of Theorem 8.2 . The upper bound (8.21) simply results from a first order Tchebychev inequality and (8.14). The proof of (8.18) is more involved. It relies on a second order Tchebychev inequality, that is, using (8.13), we bound the left hand side of (8.18) from above by

$$\epsilon^{-2}(k_n^\circ(t))^2 \sum_{x\in\mathcal{V}_n^\circ}\sum_{y\in\mathcal{V}_n^\circ} h_n^u(x)h_n^u(y)E_{\pi_n^\circ}^\circ\left(\pi_n^{J_n^\circ,t}(x)-\pi_n^\circ(x)\right)\left(\pi_n^{J_n^\circ,t}(y)-\pi_n^\circ(y)\right) . \quad (8.22)$$

In view of (8.10), setting $\Delta_{ij}(x,y) = P_{\pi_n^\circ}^\circ\left(J_n^\circ(i)=x, J_n^\circ(j)=y\right) - \pi_n^\circ(x)\pi_n^\circ(y)$, the expectation in (8.22) may be rewritten as

$$E_{\pi_n^\circ}^\circ\left(\pi_n^{J_n^\circ,t}(x)-\pi_n^\circ(x)\right)\left(\pi_n^{J_n^\circ,t}(y)-\pi_n^\circ(y)\right) = \sum_{i=0}^{k_n^\circ(t)-1}\sum_{j=0}^{k_n^\circ(t)-1}\Delta_{ij}(x,y). \quad (8.23)$$

For ℓ_n° defined in (6.1) we now break the sum in the r.h.s. of (8.23) into three terms:

$$\begin{aligned} I_1^{(1)} &= 2\sum_{0\le i\le k_n^\circ(t)-1}\sum_{i+\ell_n^\circ\le j\le k_n^\circ(t)-1}\Delta_{ij}(x,y)\,,\\ I_2^{(1)} &= \sum_{0\le i\le k_n^\circ(t)-1}\mathbb{1}_{\{i=j\}}\Delta_{ij}(x,y)\,,\\ I_3^{(1)} &= 2\sum_{0\le i\le k_n^\circ(t)-1}\sum_{i<j<i+\ell_n^\circ}\Delta_{ij}(x,y)\,. \end{aligned} \quad (8.24)$$

Consider first $I_1^{(1)}$. By Proposition 6.1,

$$I_1^{(1)} \le \delta_n(k_n^\circ(t))^2\pi_n^\circ(x)\pi_n^\circ(y) \le 2^{-n}(k_n^\circ(t))^2\pi_n^\circ(x)\pi_n^\circ(y). \quad (8.25)$$

Turning to the term $I_2^{(1)}$, we have,

$$I_2^{(1)} = \sum_{1\le i\le k_n^\circ(t)}\Delta_{ii}(x,x)\mathbb{1}_{\{x=y\}} = k_n^\circ(t)\pi_n^\circ(x)(1-\pi_n^\circ(x))\mathbb{1}_{\{x=y\}}, \quad (8.26)$$

where we used that $P_{\pi_n^\circ}^\circ(J_n^\circ(i)=x) = \pi_n^\circ(x)$. Finally,

$$\begin{aligned} I_3^{(1)} &\le 2\sum_{i=0}^{k_n^\circ(t)-1}\sum_{l=1}^{\ell_n^\circ-1}P_{\pi_n^\circ}^\circ\left(J_n^\circ(i)=x, J_n^\circ(i+l)=y\right)\\ &= 2k_n^\circ(t)\pi_n^\circ(x)\sum_{l=1}^{\ell_n^\circ-1}p_n^{\circ,l}(x,y) \end{aligned} \quad (8.27)$$

where $p_n^{\circ,l}(\cdot,\cdot)$ denote the l-steps transition probabilities of J_n°. Combining our bounds on $(\overline{I}), I_2^{(1)}$, and $I_3^{(1)}$ with (8.22) we get that, for all $\epsilon>0$,

$$P_{\pi_n^\circ}^\circ\left(\left|\nu_n^{J_n^\circ,t}(u,\infty) - E_{\pi_n^\circ}^\circ\left[\nu_n^{J_n^\circ,t}(u,\infty)\right]\right| \ge \epsilon\right) \le \epsilon^{-2}[I_1^{(2)}+I_2^{(2)}+I_3^{(2)}]\,, \quad (8.28)$$

where

$$\begin{aligned} I_1^{(2)} &= 2^{-n}(k_n^\circ(t))^2\sum_{x\in\mathcal{V}_n^\circ}\sum_{y\in\mathcal{V}_n^\circ}h_n^u(x)h_n^u(y)\pi_n^\circ(x)\pi_n^\circ(y)\,,\\ I_2^{(2)} &= k_n^\circ(t)\sum_{x\in\mathcal{V}_n^\circ}\sum_{y\in\mathcal{V}_n^\circ}h_n^u(x)h_n^u(y)\pi_n^\circ(x)(1-\pi_n^\circ(x))\mathbb{1}_{\{x=y\}}\,,\\ I_3^{(2)} &= 2k_n^\circ(t)\sum_{x\in\mathcal{V}_n^\circ}\sum_{y\in\mathcal{V}_n^\circ}h_n^u(x)h_n^u(y)\pi_n^\circ(x)\sum_{l=1}^{\ell_n^\circ-1}p_n^{\circ,l}(x,y). \end{aligned} \quad (8.29)$$

In view of (8.13)-(8.14),

$$I_1^{(2)} \le 2^{-n}\left(k_n^\circ(t)/a_n\right)^2\left[\nu_n^\circ(u,\infty)\right]^2, \quad (8.30)$$

$$I_2^{(2)} \le (k_n^\circ(t)/a_n)\sigma_n^\circ(u,\infty). \quad (8.31)$$

To deal with the third term in (8.29) note first that by (8.2),

$$\sum_{y\in\mathcal{V}_n^\circ}p_n^{\circ,l}(x,y)h_n^u(y) = \sum_{z\in\mathcal{V}_n^\circ}p_n^{\circ,l+1}(x,z)e^{-uc_n\lambda_n(z)}, \quad (8.32)$$

so that

$$\sum_{x\in\mathcal{V}_n^\circ}\pi_n^\circ(x)h_n^u(x)p_n^{\circ,l+1}(x,z) = \sum_{y\in\mathcal{V}_n^\circ}e^{-uc_n\lambda_n(y)}\sum_{x\in\mathcal{V}_n^\circ}\pi_n^\circ(x)p_n^\circ(x,y)p_n^{\circ,l+1}(x,z)$$
$$= \sum_{y\in\mathcal{V}_n^\circ}e^{-uc_n\lambda_n(y)}\pi_n^\circ(y)p_n^{\circ,l+2}(y,z)\,, \tag{8.33}$$

where the last equality follows by reversibility. Hence,

$$I_3^{(2)} = 2k_n^\circ(t)\sum_{l=1}^{\ell_n^\circ-1}\sum_{z\in\mathcal{V}_n^\circ}\Big[\sum_{x\in\mathcal{V}_n^\circ}\pi_n^\circ(x)h_n^u(x)p_n^{\circ,l+1}(x,z)\Big]e^{-uc_n\lambda_n(z)}\,,$$
$$= 2\sum_{l=1}^{\ell_n^\circ-1}k_n^\circ(t)\sum_{z\in\mathcal{V}_n^\circ}\sum_{y\in\mathcal{V}_n^\circ}\pi_n^\circ(y)e^{-uc_n(\lambda_n(y)+\lambda_n(z))}p_n^{\circ,l+2}(y,z)$$
$$\equiv 2(k_n^\circ(t)/a_n)\sum_{z\in\mathcal{V}_n^\circ}\sum_{y\in\mathcal{V}_n^\circ}f_n(y,z) \tag{8.34}$$

where the last line defines $f_n(y,z)$. In view of (8.17), we have

$$\sum_{z\in\mathcal{V}_n^\circ\setminus I_n^\star}\sum_{y\in\mathcal{V}_n^\circ\setminus I_n^\star}f_n(y,z)\le\ell_n^\circ e^{-2uc_n}, \tag{8.35}$$

$$\sum_{z\in\mathcal{V}_n^\circ\setminus I_n^\star}\sum_{y\in I_n^\star}f_n(y,z)=\sum_{z\in I_n^\star}\sum_{y\in\mathcal{V}_n^\circ\setminus I_n^\star}f_n(y,z)\le\ell_n^\circ e^{-uc_n}\nu_n^\circ(u,\infty), \tag{8.36}$$

where the equality above is reversibility. It thus remains to bound the term

$$I^{(3)}\equiv 2\frac{k_n^\circ(t)}{a_n}\sum_{z\in I_n^\star}\sum_{y\in I_n^\star}f_n(y,z)=2\frac{k_n^\circ(t)}{a_n}\sum_{l=1}^{\ell_n^\circ-1}[I_{1,l}^{(3)}+I_{2,l}^{(3)}], \tag{8.37}$$

where, distinguishing the cases $z=y$ and $z\neq y$,

$$I_{1,l}^{(3)}\equiv\sum_{z\in I_n^\star}a_n\pi_n^\circ(z)e^{-2uc_n\lambda_n(z)}p_n^{\circ,l+2}(z,z), \tag{8.38}$$

$$I_{2,l}^{(3)}\equiv\sum_{z\in I_n^\star}\sum_{y\in I_n^\star:y\neq z}a_n\pi_n^\circ(y)e^{-uc_n(\lambda_n(y)+\lambda_n(z))}p_n^{\circ,l+2}(y,z). \tag{8.39}$$

By Proposition 6.2 we readily have that, on Ω^{SRW}, for all but a finite number of indices n,

$$\sum_{l=1}^{\ell_n^\circ-1}I_{1,l}^{(3)}\le\frac{C_\circ}{n\log n}\nu_n^\circ(2u,\infty). \tag{8.40}$$

for some constant $0<C_\circ<\infty$. The next lemma is designed to deal with (8.39).

Lemma 8.3. *Let $\rho_n^\circ>0$ be a decreasing sequence satisfying $\rho_n^\circ\downarrow 0$ as $n\uparrow\infty$. There exists a sequence of subsets $\Omega_{2,n}^{(3)}\subset\Omega$ with $\mathbb{P}\big(\Omega_{2,n}^{(3)}\big)\ge 1-\ell_n^\circ/(\rho_n^\circ a_n)$ such that on $\Omega_{2,n}^{(3)}$,*

$$\sum_{l=1}^{\ell_n^\circ-1}I_{2,l}^{(3)}<\rho_n^\circ\left[\mathbb{E}\nu_n^\circ(u,\infty)\right]^2\,. \tag{8.41}$$

Proof. By definition of $I_n^\star$, $\mathrm{dist}(y,z) \geq 2$ for all $y \in I_n^\star$ and $z \in I_n^\star$ such that $y \neq z$. Thus

$$\mathbb{E}\left(e^{-uc_n(\lambda_n(y)+\lambda_n(z))}\mathbb{1}_{\{y\in I_n^\star, z\in I_n^\star\}}\right)\mathbb{1}_{\{x\neq z\}} \leq \mathbb{E}\left(e^{-uc_n(\lambda_n(y)+\lambda_n(z))}\right)\mathbb{1}_{\{\mathrm{dist}(y,z)\geq 2\}} \quad (8.42)$$

$$\leq \left(\mathbb{E}e^{-uc_n\lambda_n(y)}\right)\left(\mathbb{E}e^{-uc_n\lambda_n(z)}\right) \quad (8.43)$$

$$= \left[a_n^{-1}\mathbb{E}\nu_n^\circ(u,\infty)\right]^2 \quad (8.44)$$

where we used independence in the second line. Therefore, by a first order Tchebychev inequality, for all $\eta > 0$,

$$\mathbb{P}\big(\textstyle\sum_{l=1}^{\ell_n^\circ-1} I_{2,l}^{(3)} \geq \eta\big) \leq \frac{1}{\eta a_n}\left[\mathbb{E}\nu_n(u,\infty)\right]^2 \sum_{l=1}^{\ell_n^\circ-1}\sum_{y\in\mathcal{V}_n^\circ}\pi_n^\circ(y)\sum_{z\in\mathcal{V}_n^\circ}p_n^{\circ,l+2}(y,z) \quad (8.45)$$

$$\leq \frac{\ell_n^\circ}{\eta a_n}\left[\mathbb{E}\nu_n^\circ(u,\infty)\right]^2. \quad (8.46)$$

The lemma now easily follows. □

Gathering our bounds we conclude that under the assumptions and with the notations of Proposition 6.2 and Lemma 8.3, on $\Omega^{\mathsf{SRW}} \cap \Omega_{2,n}^{(3)}$, for all but a finite number of indices n,

$$I_3^{(2)} \leq 2\frac{k_n^\circ(t)}{a_n}\left[\ell_n^\circ e^{-uc_n}[1+\nu_n^\circ(u,\infty)] + C_\circ\frac{\nu_n^\circ(2u,\infty)}{n\log n} + \rho_n^\circ\left[\mathbb{E}\nu_n^\circ(u,\infty)\right]^2\right] \quad (8.47)$$

for some constant $0 < C_\circ < \infty$. Inserting the bounds (8.30), (8.31), and (8.47) in (8.28) now yields (8.18)-(8.20). The proof of Theorem 8.2 is done. □

8.3. **Almost sure convergence of ν_n° and σ_n°.** Theorem 8.2 enables us to replace the chain dependant quantities $\nu_n^{j_n^\circ,t}$ and $\sigma_n^{j_n^\circ,t}$ by quantities, ν_n° and σ_n°, that now only depend on the randomness of the environment. Our next step consists in proving laws of large numbers for ν_n° and σ_n°.

Proposition 8.4. *Under the assumptions and with the notation of Theorem 1.1 there exists a subset $\Omega^{\mathsf{LLN}} \subset \Omega$ with $\mathbb{P}(\Omega^{\mathsf{LLN}}) = 1$ such that, on Ω^{LLN}, the following holds: for all $u > 0$,*

$$\lim_{n\to\infty}\nu_n^\circ(u,\infty) = \nu(u,\infty), \quad (8.48)$$

$$\lim_{n\to\infty} n\sigma_n^\circ(u,\infty) = \nu(2u,\infty). \quad (8.49)$$

We prove the proposition by comparing ν_n° and σ_n° to their counterpart, ν_n^{REM} and σ_n^{REM}, in the random hopping dynamics of the non truncated REM. To define ν_n^{REM} and σ_n^{REM} recall the definition of $w_n(x)$ from (1.15) and set

$$\gamma_n(x) = w_n(x)/c_n. \quad (8.50)$$

Then, for all $u > 0$,

$$\nu_n^{\mathsf{REM}}(u,\infty) = \frac{a_n}{|\mathcal{V}_n|}\sum_{x\in\mathcal{V}_n} e^{-u/\gamma_n(x)}, \quad (8.51)$$

$$\sigma_n^{\mathsf{REM}}(u,\infty) = \frac{a_n}{|\mathcal{V}_n|}\sum_{x\in\mathcal{V}_n}\sum_{x'\in\mathcal{V}_n} p_n^{\mathsf{SRW},(2)}(x,x')e^{-u(1/\gamma_n(x)+1/\gamma_n(x'))}, \quad (8.52)$$

where $p_n^{\mathsf{SRW},(2)}(\cdot,\cdot)$ denotes the 2-steps transition probabilities of J_n^{SRW} (see (6.55)). For later use (namely, for the treatment of Condition (C3)) we also define, for all $\epsilon > 0$,

$$\eta_n^{\mathsf{REM}}(\epsilon) = \frac{a_n}{|\mathcal{V}_n|}\sum_{x\in\mathcal{V}_n}\gamma_n(x)\left(1-e^{-\epsilon/\gamma_n(x)}\right). \quad (8.53)$$

The functions ν_n^{REM}, σ_n^{REM}, and η_n^{REM} are well understood. We know in particular that:

Proposition 8.5. *Given $0 < \varepsilon < 1$ let a_n and c_n be as in Theorem 1.1. Let ν be as in (1.21) and assume that $\beta > \beta_c(\varepsilon)$. Then, there exists a subset $\Omega^{REM} \subset \Omega$ with $\mathbb{P}(\Omega^{REM}) = 1$ such that, on Ω^{REM}, the following holds:*

$$\lim_{n\to\infty} \nu_n^{REM}(u,\infty) = \nu(u,\infty), \quad \forall u > 0, \tag{8.54}$$

$$\lim_{n\to\infty} n\sigma_n^{REM}(u,\infty) = \nu(2u,\infty), \quad \forall u > 0, \tag{8.55}$$

and

$$\lim_{\epsilon\to 0}\lim_{n\to\infty} \eta_n^{REM}(\epsilon) = 0. \tag{8.56}$$

Throughout this section we set $\varepsilon_n \equiv \frac{\log a_n}{n\log 2}$; thus by 1.19, $\lim_{n\to\infty}\varepsilon_n = \varepsilon$, $0 < \varepsilon < 1$.

Proof. Eq. (8.54) and (8.55) are proved in Proposition 5.1, (i), in Section 5.1 of [25]. The proof of (8.56) is elementary: by simple Gaussian calculations, $\mathbb{E}\eta_n^{\mathsf{REM}}(\epsilon) \le c\epsilon^{1-\alpha(\varepsilon_n)} \downarrow 0$ as $n \uparrow \infty$ and $\epsilon \downarrow 0$, where $0 < c < \infty$ is a constant, and $\mathbb{P}\left(|\eta_n^{\mathsf{REM}}(\epsilon) - \mathbb{E}\eta_n^{\mathsf{REM}}(\epsilon)| > n^{-1}\right) \le n^3 a_n/|\mathcal{V}_n|$, which is summable under our assumptions on a_n. Since $\eta_n^{\mathsf{REM}}(\epsilon)$ is a monotonic function of $\epsilon > 0$, arguing e.g. as in (9.135) yields the claim (8.56). □

Our next lemma establishes that ν_n° and σ_n° are very close to ν_n^{REM} and σ_n^{REM}.

Lemma 8.6. *On Ω_3, for all but a finite number of indices n, for all $u > 0$,*

$$|\nu_n^\circ(u,\infty) - \nu_n^{REM}(u,\infty)| \le 2n^{-2c_\star+1}\nu_n^{REM}(u,\infty) + 2a_n e^{-un^2} + 2n^{-c_\star+1+2\alpha(\varepsilon_n)}, \tag{8.57}$$

$$|\sigma_n^\circ(u,\infty) - \sigma_n^{REM}(u,\infty)| \le 2n^{-2c_\star+1}\sigma_n^{REM}(u,\infty) + 4a_n e^{-un^2} + 2n^{-c_\star+1+2\alpha(\varepsilon_n)}. \tag{8.58}$$

Proof of Lemma 8.6 . The proof hinges on the observation that $c_n\lambda_n(x) = 1/\gamma_n(x)$ for all x in the subset $I_n^\star$ of the decomposition (2.9). This enables us to rewrite $\nu_n^\circ(u,\infty)$ as

$$\nu_n^\circ(u,\infty) = (|\mathcal{V}_n|/|\mathcal{V}_n^\circ|)\nu_n^{\mathsf{REM}}(u,\infty) + I_1 - I_2 - I_3 \tag{8.59}$$

where

$$I_3 \equiv (a_n/|\mathcal{V}_n^\circ|)\textstyle\sum_{x\in\cup_{l=1}^{L^\star} C_{n,l}^\star} e^{-u/\gamma_n(x)}, \tag{8.60}$$

$$I_1 \equiv (a_n/|\mathcal{V}_n^\circ|)\textstyle\sum_{x\in\mathcal{V}_n^\circ\setminus I_n^\star} e^{-uc_n\lambda_n(x)} \le a_n e^{-uc_n}, \tag{8.61}$$

$$I_2 \equiv (a_n/|\mathcal{V}_n^\circ|)\textstyle\sum_{x\in\mathcal{V}_n^\circ\setminus I_n^\star} e^{-u/\gamma_n(x)} \le a_n e^{-uc_n/r_n(\rho_n^\star)}. \tag{8.62}$$

The bounds on I_1 and I_2 follow from the fact that on $\mathcal{V}_n^\circ \setminus I_n^\star \equiv N_n^\star$, $\lambda_n(x) = 1$ and $w_n(x) < r_n(\rho_n^\star)$. In order to bound I_3 recall 2.1 and set $\mathcal{W}_n(\rho) \equiv (\cup_{l=1}^{L^\star} C_{n,l}^\star) \cap V(\rho)$ and $\mathcal{W}_n^c(\rho) \equiv (\cup_{l=1}^{L^\star} C_{n,l}^\star) \cap V^c(\rho)$ for some $\rho > 0$. Then, on $\mathcal{W}_n^c(\rho)$, by (2.19) of Lemma 2.4,

$$\tfrac{w_n(x)}{c_n} \le \tfrac{r_n(\rho)}{r_n(\varepsilon_n)} = \exp\{n\beta\beta_c(1)(\sqrt{\varepsilon}_n - \sqrt{\rho}) - \tfrac{\beta\log n}{2\beta_c(1)}(\tfrac{1}{\sqrt{\varepsilon}_n} - \tfrac{1}{\sqrt{\rho}}) + o(1)\}, \tag{8.63}$$

so that choosing $\sqrt{\rho} = \sqrt{\varepsilon}_n - \frac{2\log n}{n\beta\beta_c(1)}$, we get

$$\tfrac{r_n(\rho)}{r_n(\varepsilon_n)} = n^2 \exp\left\{\tfrac{\log n}{n\beta\varepsilon_n 2\log 2}(1 + o(1))\right\} = n^2(1 + o(1)). \tag{8.64}$$

One also sees that for this choice of ρ, $4\rho_n^\star < \rho < 1 - 4\rho_n^\star$ for all $0 < \varepsilon < 1$ and large enough n. Therefore Lemma 7.1 applies, yielding

$$|\mathcal{W}_n(\rho)|\,/|\mathcal{V}_n^\circ| \le n^{-c_\star+1}2^{-n\rho}(1 + o(1)), \tag{8.65}$$

on Ω_3, for all n large enough. Assume from now on that $\omega \in \Omega_3$. By (8.64) and (8.65),

$$I_3 \le a_n e^{-un^2} + 2n^{-c_\star+1}2^{n(\varepsilon_n-\rho)} \le a_n e^{-un^2} + 2n^{-c_\star+1}n^{2\beta_c(\varepsilon_n)/\beta}, \tag{8.66}$$

where we used that $\varepsilon_n - \rho = (\sqrt{\varepsilon_n} - \sqrt{\rho})(\sqrt{\varepsilon_n} + \sqrt{\rho}) \le 2\sqrt{\varepsilon_n}\frac{2\log n}{n\beta\beta_c(1)}$. Eq. (8.57) now easily follows observing that, by (2.19) and (2.20) of Lemma 2.4, $c_n \gg c_n/r_n\big(\rho_n^\star\big) \gg n^2$, and using that $|\mathcal{V}_n|/|\mathcal{V}_n^\circ| = 1 + n^{-2c_\star+1}(1 + \mathcal{O}(n^{-(c_\star-1)}))$, as follows from (6.6).

The proof of (8.58) follows the same pattern, using the additionnal observation that $p_n^{\circ,2}(x,x') = p_n^{\mathsf{SRW},2}(x,x')$ for all x, x' in $I_n^\star \times I_n^\star$. This follows from Proposition 6.5 and the fact that, by construction, $I_n^\star \cap \partial C_{n,l}^\star = \emptyset$ for all $1 \le l \le L^\star$. We skip the details. □

Proof of Proposition 8.4 . The proposition is now an immediate consequence of Lemma 8.6 and Proposition 8.5. □

8.4. **Conclusion of the proof of Theorem 3.2.** We are now ready to show that under the assumptions of Theorem 1.1, taking for initial distribution the invariant measure π_n° of J_n°, the conditions of Theorem 8.1 are satisfied $\mathbb{P}$-almost surely. Firstly, by Theorem 8.2 and Proposition 8.4, Conditions (C1) and (C2) are satisfied $\mathbb{P}$-almost surely. That is, $\mathbb{P}$-almost surely the following holds: for all $u > 0$ and all $t > 0$,

$$\lim_{n\to\infty} \nu_n^{J_n^\circ,t}(u,\infty) = t\nu^\circ(u,\infty) \quad \text{in } \mathcal{P}^\circ\text{-probability}, \tag{8.67}$$

$$\lim_{n\to\infty} \sigma_n^{J_n^\circ,t}(u,\infty) = 0 \quad \text{in } \mathcal{P}^\circ\text{-probability}. \tag{8.68}$$

Next, in view of (8.15), (8.5) reads $\nu_n^\circ(v,\infty)/a_n = o(1)$, and so, by (8.48) of Proposition 8.4, Condition (C0) is satisfied. It remains to check Condition (C3). As in the proof of Proposition 8.4, we do this by comparing the quantity

$$\eta_n^\circ(\epsilon) \equiv \lfloor a_n \rfloor \mathcal{E}_{\pi_n^\circ}^\circ \mathbb{1}_{\{\lambda_n^{-1}(J_n^\circ(0))e_0^\circ \le c_n\epsilon\}} c_n^{-1}\lambda_n^{-1}(J_n^\circ(0))e_0^\circ \tag{8.69}$$

$$= \frac{\lfloor a_n \rfloor}{|\mathcal{V}_n^\circ|} \sum_{x\in\mathcal{V}_n^\circ} c_n^{-1}\lambda_n^{-1}(x)\left(1 - e^{-\epsilon c_n \lambda_n(x)}\right) \tag{8.70}$$

arising in (8.8), to its counterpart in the random hopping dynamics of the non truncated REM, $\eta_n^{\mathsf{REM}}(\epsilon)$, defined in (8.53). For this we simply write that since $\lambda_n(x) = 1$ on $\mathcal{V}_n^\circ \setminus I_n^\star$ and $c_n\lambda_n(x) = 1/\gamma_n(x)$ on $I_n^\star$,

$$\eta_n^\circ(\epsilon) \le \frac{\lfloor a_n \rfloor}{c_n} + \frac{\lfloor a_n \rfloor}{|\mathcal{V}_n^\circ|} \sum_{x\in I_n^\star} \gamma_n(x)\left(1 - e^{-\epsilon/\gamma_n(x)}\right) \le \frac{\lfloor a_n \rfloor}{c_n} + \frac{|\mathcal{V}_n|}{|\mathcal{V}_n^\circ|}\eta_n^{\mathsf{REM}}(\epsilon). \tag{8.71}$$

From this, (6.6), and (8.56) it follows that, under the assumptions of Proposition 8.5,

$$\lim_{\epsilon\to 0}\lim_{n\to\infty} \eta_n^\circ(\epsilon) = 0, \quad \mathbb{P}\text{-almost surely}. \tag{8.72}$$

Therefore Condition (C3) is satisfied $\mathbb{P}$-almost surely.

Since all four conditions (C0), (C1), (C2), and (C3) are satisfied $\mathbb{P}$-almost surely, it follows from Theorem 8.1 that, for our choices of a_n, c_n, β, and $c_\star$, $\mathbb{P}$-almost surely,

$$S_n^\circ \Rightarrow_{J_1} S_\infty^\circ \tag{8.73}$$

where S_∞° is a subordinator with zero drift and Lévy measure $\nu^\circ = \nu$ defined in (1.21). The proof of Theorem 3.2 is complete.

9. Convergence of the back end clock process below the critical temperature: proof of Theorem 3.4

9.1. **A convergence theorem for BECP.** Consider the rescaled process (3.16),

$$S_n^\dagger(t) = b_n^{-1}\widetilde{S}_n^\dagger(k_n^\dagger(t)), \quad t \ge 0. \tag{9.1}$$

Theorem 9.1 below parallels Theorem 8.1 for FECP, namely, it gives three sufficient conditions for the sequence $S_n^\dagger$ to converge to a subordinator when the initial distribution of $J_n^\dagger$ is the invariant measure π_n° of J_n°. As before these conditions refer to given sequences of numbers a_n and b_n, and a given realization of the random environment. For $u > 0$ define

$$\bar{h}_n^u(y) = \sum_{1\le l\le L^\star} \sum_{x\in C_{n,l}^\star} p_n(y,x) P_x(T_{n,l}^\star > b_n u)\,,\ y \in \mathcal{V}_n^\circ, \tag{9.2}$$

where $T_{n,l}^\star$ is the exit time (5.9). (Note that $\bar{h}_n^u(y) = 0$ unless $y \in \cup_{1\le l\le L^\star} \partial C_{n,l}^\star$.) For $k_n^\circ(t)$ as in (3.19) define, for $t > 0$ and $u > 0$,

$$\bar{\nu}_n^{J_n^\circ,t}(u,\infty) \;=\; \sum_{j=0}^{k_n^\circ(t)-1} \bar{h}_n^u(J_n^\circ(j)), \tag{9.3}$$

$$\bar{\sigma}_n^{J_n^\circ,t}(u,\infty) \;=\; \sum_{j=0}^{k_n^\circ(t)-1} \left[\bar{h}_n^u(J_n^\circ(j))\right]^2. \tag{9.4}$$

Condition (A1). There exists a σ-finite measure $\nu^\dagger$ on $(0,\infty)$ satisfying $\int_0^\infty (1\wedge u)\nu^\dagger(du) < \infty$ such that, for all $t > 0$ and all $u > 0$,

$$P_{\pi_n^\circ}^\circ\left(\left|\bar{\nu}_n^{J_n^\circ,t}(u,\infty) - t\nu^\dagger(u,\infty)\right| < \epsilon\right) = 1 - o(1)\,, \quad \forall \epsilon > 0\,. \tag{9.5}$$

Condition (A2). For all $u > 0$ and all $t > 0$,

$$P_{\pi_n^\circ}^\circ\left(\bar{\sigma}_n^{J_n^\circ,t}(u,\infty) < \epsilon\right) = 1 - o(1)\,, \quad \forall \epsilon > 0\,. \tag{9.6}$$

Condition (A3). For all $t > 0$,

$$\lim_{\epsilon\downarrow 0} \limsup_{n\uparrow\infty} \frac{k_n^\circ(t)}{|\mathcal{V}_n^\circ|} \sum_{1\le l\le L^\star} \sum_{x\in C_{n,l}^\star} E_x\left(\mathbb{1}_{\{b_n^{-1}T_{n,l}^\star \le \epsilon\}} b_n^{-1} T_{n,l}^\star\right) = 0. \tag{9.7}$$

Theorem 9.1. *Choose for initial distribution the invariant measure π_n° of J_n°. For all sequences a_n and b_n for which Conditions (A1), (A2), and (A3) are verified $\mathbb{P}$-almost surely,*

$$S_n^\dagger \Rightarrow_{J_1} S_\infty^\dagger \tag{9.8}$$

$\mathbb{P}$-almost surely, where $S_\infty^\dagger$ is the Lévy subordinator with zero drift and Lévy measure $\nu^\dagger$.

Proof of Theorem 9.1. The proof of Theorem 3.4 relies on Theorem 2.1 of [26], which is itself a specialization of Theorem 4.1 of [22] to processes with non-negative increments. Throughout we fix a realisation $\omega \in \Omega^\star$ of the random environment but do not make this explicit in the notation. With the notations of Subsection 3.2 define, for $i \ge 0$,

$$Z_{n,i} \equiv b_n^{-1} \Lambda_n^\dagger(J_n^\dagger(i))\,. \tag{9.9}$$

Thus, by (3.11) and (9.1),

$$S_n^\dagger(t) = \textstyle\sum_{i=0}^{k_n^\dagger(t)-1} Z_{n,i}. \tag{9.10}$$

In view of (3.15), $k_n^\dagger(t)$ is a stopping time for each $t > 0$. Furthermore, because $J_n^\dagger$ starts in π_n°, and because $\pi_n^\circ(\mathcal{V}_n \setminus \mathcal{V}_n^\circ) = 0$, it follows from (3.12) that $Z_{n,0} = 0$. We may thus apply Theorem 2.1 of [26] to the sum (9.10).

To this end let $\{\mathcal{F}_{n,i}^\dagger, n \ge 1, i \ge 0\}$ be the array of sub-sigma fields defined (with obvious notation) by $\mathcal{F}_{n,i}^\dagger = \sigma\left(J_n^\dagger(0),\ldots,J_n^\dagger(i)\right)$, for $i \ge 0$. Clearly, for each n and

$i \geq 1$, $Z_{n,i}$ is $\mathcal{F}^\dagger_{n,i}$ measurable and $\mathcal{F}^\dagger_{n,i-1} \subset \mathcal{F}^\dagger_{n,i}$. Next, observe that

$$\begin{aligned}\mathcal{P}^\dagger_{\pi^\circ_n}\big(Z_{n,i} > u \,\big|\, \mathcal{F}^\dagger_{n,i-1}\big) &= \textstyle\sum_{x\in\mathcal{V}_n} \mathcal{P}^\dagger_{\pi^\circ_n}\big(J^\dagger_n(i) = x, Z_{n,i} > u \,\big|\, J^\dagger_n(i-1)\big) \\ &= \textstyle\sum_{x\in\mathcal{V}_n} \mathcal{P}^\dagger_{\pi^\circ_n}\big(J^\dagger_n(i) = x, \Lambda^\dagger_n(x) > b_n u \,\big|\, J^\dagger_n(i-1)\big). \quad (9.11)\end{aligned}$$

By (3.12) and (5.9), $\Lambda^\dagger_n(x) = 0$ if $x \notin \cup_{1\leq l\leq L^\star} C^\star_{n,l}$, and $\Lambda^\dagger_n(x) = T^\star_{n,l}$ if $x \in C^\star_{n,l}$. Thus,

$$\mathcal{P}^\dagger_{\pi^\circ_n}\big(Z_{n,i} > u \,\big|\, \mathcal{F}^\dagger_{n,i-1}\big) = \sum_{1\leq l\leq L^\star} \sum_{x\in C^\star_{n,l}} \mathcal{P}^\dagger_{\pi^\circ_n}\big(J^\dagger_n(i) = x, \Lambda^\dagger_n(x) > b_n u \,\big|\, J^\dagger_n(i-1)\big). \quad (9.12)$$

Now for all $1 \leq l \leq L^\star$ and all $x \in C^\star_{n,l}$,

$$\mathcal{P}^\dagger_{\pi^\circ_n}\big(J^\dagger_n(i) = x, \Lambda^\dagger_n(x) > b_n u \,\big|\, J^\dagger_n(i-1)\big) = p^\dagger_n\big(J^\dagger_n(i-1), x\big) P_x\big(T^\star_{n,l} > b_n u\big) \quad (9.13)$$

where, by (3.13),

$$p^\dagger_n\big(J^\dagger_n(i-1), x\big) = p_n\big(J^\dagger_n(i-1), x\big) \mathbb{1}_{\{J^\dagger_n(i-1)\in\mathcal{V}^\circ_n\}} \quad (9.14)$$

(indeed, by definition of $J^\dagger_n$, $J^\dagger_n(i) \in C^\star_{n,l}$ if and only if $J^\dagger_n(i-1) \in \partial C^\star_{n,l} \subset \mathcal{V}^\circ_n$). In view of (9.2) it follows from (9.11), (9.12), (9.13) and (9.14) that

$$\textstyle\sum_{i=1}^{k^\dagger_n(t)} \mathcal{P}^\dagger_{\pi^\circ_n}\big(Z_{n,i} > u \,\big|\, \mathcal{F}^\dagger_{n,i-1}\big) = \sum_{i=1}^{k^\dagger_n(t)} \bar{h}^u_n\big(J^\dagger_n(i-1)\big) \mathbb{1}_{\{J^\dagger_n(i-1)\in\mathcal{V}^\circ_n\}}. \quad (9.15)$$

It remains to notice that the chain $J^\dagger_n$ observed only when it takes values in $\mathcal{V}^\circ_n$ is nothing but the chain J°_n, and that J°_n takes $k^\circ_n(t)$ steps when $J^\dagger_n$ takes $k^\dagger_n(t)$ steps (see (3.19)). Thus,

$$\textstyle\sum_{i=1}^{k^\dagger_n(t)} \bar{h}^u_n\big(J^\dagger_n(i-1)\big) \mathbb{1}_{\{J^\dagger_n(i-1)\in\mathcal{V}^\circ_n\}} \stackrel{d}{=} \sum_{i=1}^{k^\circ_n(t)} \bar{h}^u_n\big(J^\circ_n(i-1)\big) = \bar{\nu}^{J^\circ_n,t}_n(u,\infty), \quad (9.16)$$

where the first equality holds in distribution and the last is (9.3). Combining (9.15) and (9.16) now yields

$$\textstyle\sum_{i=1}^{k^\dagger_n(t)} \mathcal{P}^\dagger_{\pi^\circ_n}\big(Z_{n,i} > u \,\big|\, \mathcal{F}^\dagger_{n,i-1}\big) \stackrel{d}{=} \bar{\nu}^{J^\circ_n,t}_n(u,\infty). \quad (9.17)$$

Similarly, we get

$$\textstyle\sum_{i=1}^{k_n(t)-1} \Big[\mathcal{P}^\dagger_{\pi^\circ_n}\big(Z_{n,i} > u \,\big|\, \mathcal{F}^\dagger_{n,i-1}\big)\Big]^2 \stackrel{d}{=} \sum_{i=1}^{k^\dagger_n(t)} \big[\bar{h}^u_n\big(J^\circ_n(i-1)\big)\big]^2 = \bar{\sigma}^{J^\circ_n,t}_n(u,\infty). \quad (9.18)$$

From (9.17) and (9.18) it follows that Conditions (A2) and (A1) of Theorem 9.1 are exactly the Conditions (D1) and (D2) of Theorem 2.1 of [26]. To see that Condition (A3) implies Condition (D3) we have to establish that (9.7) implies

$$\lim_{\epsilon\to 0} \limsup_{n\to\infty} \mathcal{P}^\dagger_{\pi^\circ_n}\big(S^{\dagger,\epsilon}_n(t) > \epsilon\big) = 0\,, \quad (9.19)$$

where for $\epsilon \geq 0$, $S^{\dagger,\epsilon}_n(t) = \sum_{i=0}^{k^\dagger_n(t)-1} Z_{n,i}\mathbb{1}_{\{Z_{n,i}\leq\epsilon\}}$. By Theorem 3.1 with $c_\circ = 1$, on $\Omega^\star$, for all but a finite number of indices n, all $0 < t < \infty$, and all $\epsilon \geq 0$,

$$\mathcal{P}^\dagger_{\pi^\circ_n}\big(S^{\dagger,\epsilon}_n(t) > \epsilon\big) \leq \mathcal{P}^\dagger_{\pi^\circ_n}\Big(\textstyle\sum_{i=0}^{k_n(t)-1} Z_{n,i}\mathbb{1}_{\{Z_{n,i}\leq\epsilon\}} > \epsilon\Big) + n^{-2(c_\star-1)+1}(1+o(1)), \quad (9.20)$$

where $k_n(t) \equiv \lfloor k^\circ_n(t)(1+n^{-1})\rfloor = \lfloor\lfloor a_n t\rfloor(1+n^{-1})\rfloor$. By Tchebychev inequality,

$$\mathcal{P}^\dagger_{\pi^\circ_n}\Big(\textstyle\sum_{i=0}^{k_n(t)-1} Z_{n,i}\mathbb{1}_{\{Z_{n,i}\leq\epsilon\}} > \epsilon\Big) \leq \epsilon^{-1} \sum_{i=1}^{k_n(t)-1} \mathcal{E}^\dagger_{\pi^\circ_n} \mathbb{1}_{\{Z^n_i\leq\epsilon\}} Z^n_i, \quad (9.21)$$

and the right hand side of (9.21) is equal to

$$\textstyle\sum_{1\leq l\leq L^\star} \sum_{x\in C^\star_{n,l}} E_x\big(\mathbb{1}_{\{b_n^{-1}T^\star_{n,l}\leq\epsilon\}} b_n^{-1} T^\star_{n,l}\big) \sum_{i=1}^{k_n(t)-1} \mathcal{E}^\dagger_{\pi^\circ_n} \mathbb{1}_{\{J^\dagger_n(i)=x\}}. \quad (9.22)$$

By (7.28) and (3.13), for all $x \in C^\star_{n,l}$,

$$\begin{aligned}\textstyle\sum_{i=1}^{k_n(t)-1} \mathcal{E}^\dagger_{\pi^\circ_n} \mathbb{1}_{\{J^\dagger_n(i)=x\}} &= \textstyle\sum_{y\in\partial C^\star_{n,l}\cap\partial x} p_n(y,x) \sum_{i=1}^{k_n(t)-1} \mathcal{E}^\dagger_{\pi^\circ_n} \mathbb{1}_{\{J^\dagger_n(i-1)=y\}} \\ &\le \textstyle\sum_{y\in\partial C^\star_{n,l}\cap\partial x} p_n(y,x) \sum_{i=1}^{k_n(t)-1} \pi^\circ_n(y) \\ &\le (k_n(t)-1)/|\mathcal{V}^\circ_n| \end{aligned} \tag{9.23}$$

where the last inequality follows from (6.5) and the fact that, for $y \in \partial C^\star_{n,l} \cap \partial x$ and $x \in C^\star_{n,l}$, $p_n(y,x) = n^{-1}$. Combining (9.21), (9.22), and (9.23), the probability in the left hand side of (9.21) is bounded above by

$$\epsilon^{-1}(1+n^{-1})(k^\circ_n(t)/|\mathcal{V}^\circ_n|) \textstyle\sum_{1\le l\le L^\star} \sum_{x\in C^\star_{n,l}} E_x\big(\mathbb{1}_{\{b_n^{-1}T^\star_{n,l}\le\epsilon\}} b_n^{-1} T^\star_{n,l}\big). \tag{9.24}$$

Inserting this bound in (9.20) yields the claim that Condition (A3) implies Condition (D3) of Theorem 2.1 of [26]

Having established that, on $\Omega^\star$, the conditions of Theorem 2.1 of [26] are verified whenever those of Theorem 9.1 are verified, the proof of Theorem 9.1 is complete. □

9.2. **An ergodic theorem for BECP.** To prove that Conditions (A1) and (A2) of Theorem 9.1 are satisfied we closely follow the strategy of Subsection 8.2 and first prove an ergodic theorem for the quantities $\bar\nu_n^{J^\circ_n,t}(u,\infty)$ and $\bar\sigma_n^{J^\circ_n,t}(u,\infty)$ defined in (9.3) and (9.4). Clearly, for $\pi_n^{J^\circ_n,t}(x)$ as in (8.10), (9.3) and (9.4) can be rewritten as

$$\bar\nu_n^{J^\circ_n,t}(u,\infty) = k^\circ_n(t) \sum_{y\in\mathcal{V}^\circ_n} \pi_n^{J^\circ_n,t}(y) \bar h^u_n(y), \tag{9.25}$$

$$\bar\sigma_n^{J^\circ_n,t}(u,\infty) = k^\circ_n(t) \sum_{y\in\mathcal{V}^\circ_n} \pi_n^{J^\circ_n,t}(y) \left[\bar h^u_n(y)\right]^2. \tag{9.26}$$

Before stating our main theorem, let us express the mean values of (9.25) and (9.26) with respect to the law $P^\circ_{\pi^\circ_n}$. Given $x \in C^\star_{n,l}$, $1 \le l \le L^\star$, denote by

$$Q^u_{n,l}(x) \equiv P_x(T^\star_{n,l} > b_n u), \quad u > 0, \tag{9.27}$$

the tail distribution of the exit time $T^\star_{n,l}$ given that the set $C^\star_{n,l}$ is entered in x, and define

$$\bar\nu^\circ_n(u,\infty) = \frac{a_n}{2^n} \sum_{1\le l\le L^\star} \sum_{x\in C^\star_{n,l}} Q^u_{n,l}(x), \tag{9.28}$$

$$\bar\sigma^=_n(u,\infty) = \frac{a_n}{n2^n} \sum_{1\le l\le L^\star} \Big[\sum_{x\in C^\star_{n,l}} Q^u_{n,l}(x)\Big]^2, \tag{9.29}$$

$$\bar\sigma^\circ_n(u,\infty) = \frac{a_n}{2^n} \sum_{1\le l\le L^\star} \sum_{1\le l'\le L^\star} \sum_{x\in C^\star_{n,l}} \sum_{x'\in C^\star_{n,l'}} Q^u_{n,l}(x) Q^u_{n,l'}(x') \left(n^{-2}|\partial x\cap\partial x'|\right). \tag{9.30}$$

Lemma 9.2. *Assume that $c_\star > 2$. Then on $\Omega^\star$, for all but a finite number of indices n,*

$$E^\circ_{\pi^\circ_n}\big[\bar\nu_n^{J^\circ_n,t}(u,\infty)\big] = (1+o(1))(k^\circ_n(t)/a_n)\bar\nu^\circ_n(u,\infty), \tag{9.31}$$

$$E^\circ_{\pi^\circ_n}\big[\bar\sigma_n^{J^\circ_n,t}(u,\infty)\big] = (1+o(1))(k^\circ_n(t)/a_n)\bar\sigma^\circ_n(u,\infty). \tag{9.32}$$

The main theorem of this section controls the fluctuations of $\bar\nu_n^{J^\circ_n,t}$ around its mean value and provides an upper bound on $\bar\sigma_n^{J^\circ_n,t}$ in terms of the random (in the environment) quantities $\bar\nu^\circ_n$, $\bar\sigma^\circ_n$, and $\bar\sigma^=_n$.

Theorem 9.3. *Assume that $c_\star > 3$. Let $\bar\rho_n^\circ > 0$ be a decreasing sequence satisfying $\bar\rho_n^\circ \downarrow 0$ as $n \uparrow \infty$. There exists a sequence of subsets $\overline{\Omega}_n^{EG} \subset \Omega$ with $\mathbb{P}\big[\big(\overline{\Omega}_n^{EG}\big)^c\big] < 2^6 \ell_n^\circ/(\bar\rho_n^\circ n a_n)$, and such that on $\overline{\Omega}_n^{EG}$ the following holds for all large enough n: for all $t > 0$, all $u > 0$, and all $\epsilon > 0$,*

$$P_{\pi_n^\circ}^\circ\left(\left|\bar\nu_n^{J_n^\circ,t}(u,\infty) - E_{\pi_n^\circ}^\circ\big[\bar\nu_n^{J_n^\circ,t}(u,\infty)\big]\right| \ge \epsilon\right) \le \epsilon^{-2}[C_3 t\Theta_{n,3}(u) + t^2\Theta_{n,4}(u)] \tag{9.33}$$

for some constant $0 < C_3 < \infty$ and where, for $\varsigma_n^{\neq}(u)$ as in Lemma 9.4,

$$\Theta_{n,3}(u) \equiv \bar\sigma_n^\circ(u,\infty) + \bar\sigma_n^=(u,\infty) + \bar\rho_n^\circ\left[\varsigma_n^{\neq}(u)\right]^2, \tag{9.34}$$

$$\Theta_{n,4}(u) \equiv 2^{-n}\left[\bar\nu_n^\circ(u,\infty)\right]^2. \tag{9.35}$$

Moreover, for all $t > 0$, all $u > 0$, and all $\epsilon' > 0$,

$$P_{\pi_n^\circ}^\circ\left(\bar\sigma_n^{J_n^\circ,t}(u,\infty) \ge \epsilon'\right) \le \frac{t}{\epsilon'}(1+o(1))\bar\sigma_n^\circ(u,\infty). \tag{9.36}$$

We now prove, in this order, Lemma 9.2 and Theorem 9.3.

Proof of Lemma 9.2. By (9.2), (9.25), and (9.27),

$$\begin{aligned} E_{\pi_n^\circ}^\circ\big[\bar\nu_n^{J_n^\circ,t}(u,\infty)\big] &= k_n^\circ(t)\textstyle\sum_{y\in\mathcal{V}_n^\circ}\pi_n^\circ(y)\bar h_n^u(y) &(9.37)\\ &= k_n^\circ(t)\textstyle\sum_{y\in\mathcal{V}_n^\circ}\pi_n^\circ(y)\sum_{1\le l\le L^\star}\sum_{x\in C_{n,l}^\star}p_n(y,x)Q_{n,l}^u(x), &(9.38)\end{aligned}$$

and since both x and y belong to $\mathcal{V}_n^\circ$, $p_n(y,x) = n^{-1}$ if $\mathrm{dist}(x,y) = 1$ and is zero else. Thus $\sum_{y\in\mathcal{V}_n^\circ}p_n(y,x) = n^{-1}|\partial x \cap \partial C_{n,l}^\star|$ and

$$E_{\pi_n^\circ}^\circ\big[\bar\nu_n^{J_n^\circ,t}(u,\infty)\big] = (k_n^\circ(t)/|\mathcal{V}_n^\circ|)\textstyle\sum_{1\le l\le L^\star}\sum_{x\in C_{n,l}^\star}n^{-1}|\partial x\cap\partial C_{n,l}^\star|Q_{n,l}^u(x). \tag{9.39}$$

The claim of (9.31) now follows from (2.16) and (6.6). Eq. (9.32) is proved in a similar way. We skip the details. □

Proof of Theorem 9.3 . A first order Tchebychev inequality and (9.32) readily yield (9.36). As in Theorem 8.2, proving concentration of $\bar\nu_n^{J_n^\circ,t}(u,\infty)$ is more involved. Since (9.25) is nothing but (8.11) with h_n^u replaced by $\bar h_n^u$, the proof naturally starts in the same way as the proof of (8.18) of Theorem 8.2. More precisely, substituting $\bar h_n^u$ for h_n^u in the definition (8.29) of the quantities $I_i^{(2)}$, $1 \le i \le 3$, we get that for all $\epsilon > 0$,

$$P_{\pi_n^\circ}^\circ\left(\left|\bar\nu_n^{J_n^\circ,t}(u,\infty) - E_{\pi_n^\circ}^\circ\big[\bar\nu_n^{J_n^\circ,t}(u,\infty)\big]\right| \ge \epsilon\right) \le \epsilon^{-2}[I_1^{(2)} + I_2^{(2)} + I_3^{(2)}]. \tag{9.40}$$

We are thus left to bound $I_i^{(2)}$, $1 \le i \le 3$. By (9.25) and (9.31),

$$I_1^{(2)} = 2^{-n}\left[E_{\pi_n^\circ}^\circ\big[\bar\nu_n^{J_n^\circ,t}(u,\infty)\big]\right]^2 \le 2^{-n}\left(k_n^\circ(t)/a_n\right)^2\left[\bar\nu_n^\circ(u,\infty)\right]^2, \tag{9.41}$$

and by (9.26) and (9.32),

$$I_2^{(2)} \le E_{\pi_n^\circ}^\circ\big[\bar\sigma_n^{J_n^\circ,t}(u,\infty)\big] \le (k_n^\circ(t)/a_n)\bar\sigma_n^\circ(u,\infty). \tag{9.42}$$

The term $I_3^{(2)}$ is a little more involved. We may write it in the form

$$I_3^{(2)} \equiv 2(k_n^\circ(t)/a_n)(a_n/|\mathcal{V}_n^\circ|)\sum_{m=1}^{\ell_n^\circ-1}[I_{1,m}^{(3)} + I_{2,m}^{(3)}], \tag{9.43}$$

where, setting $f_n^{\circ,m}(x,x';y,y') \equiv p_n(y,x)p_n(y',x')p_n^{\circ,m}(y,y')$,

$$I_{1,m}^{(3)} \equiv \sum_{1\le l\le L^\star} \sum_{x\in C_{n,l}^\star} \sum_{x'\in C_{n,l}^\star} \sum_{y\in\partial C_{n,l}^\star} \sum_{y'\in\partial C_{n,l}^\star} Q_{n,l}^u(x)Q_{n,l}^u(x')f_n^{\circ,m}(x,x';y,y'), \tag{9.44}$$

$$I_{2,m}^{(3)} \equiv \sum_{1\le l,l'\le L^\star: l\neq l'} \sum_{x\in C_{n,l}^\star} \sum_{x'\in C_{n,l'}^\star} \sum_{y\in\partial C_{n,l}^\star} \sum_{y'\in\partial C_{n,l}^\star} Q_{n,l}^u(x)Q_{n,l'}^u(x')f_n^{\circ,m}(x,x';y,y'). \tag{9.45}$$

Consider $I_{1,m}^{(3)}$ first. It follows from Proposition 6.2, (ii) that on Ω^{SRW}, for all but a finite number of indices n, for all $x,x'\in C_{n,l}^\star$,

$$\sum_{m=1}^{\ell_n^\circ-1} \sum_{y\in\partial C_{n,l}^\star} \sum_{y'\in\partial C_{n,l}^\star} f_n^{\circ,m}(x,x';y,y') \le \frac{C'_\circ}{n} \sum_{y\in\partial C_{n,l}^\star} \sum_{y'\in\partial C_{n,l}^\star} p_n(y,x)p_n(y',x') \le \frac{C'_\circ}{n}. \tag{9.46}$$

(Here we used that $p_n(y,x) = p_n(y,x)$ if both x and y belong to $\mathcal{V}_n^\circ$.) From this and Proposition 6.3 we readily get that if $c_\star > 2$ then on $\Omega^{\mathsf{SRW}}\cap\Omega^\star$, for large enough n,

$$(a_n/|\mathcal{V}_n^\circ|)\textstyle\sum_{m=1}^{\ell_n^\circ-1} I_{1,m}^{(3)} \le C'_\circ(1+o(1))\bar\sigma_n^=(u,\infty). \tag{9.47}$$

The next lemma bounds the contribution to (9.43) coming from $I_{2,m}^{(3)}$. Its proof is given in Subsection 9.3.3.

Lemma 9.4. *Assume that $c_\star > 2$. Let $\bar\rho_n^\circ > 0$ be a decreasing sequence satisfying $\bar\rho_n^\circ \downarrow 0$ as $n\uparrow\infty$. There exists a sequence of subsets $\overline{\Omega}_{2,n}^{(3)} \subset \Omega$ with $\mathbb{P}\big(\overline{\Omega}_{2,n}^{(3)}\big) \ge 1 - 2^6\ell_n^\circ/(\bar\rho_n^\circ n a_n)$ such that on $\overline{\Omega}_{2,n}^{(3)}\cap\Omega^\star$, for all n large enough,*

$$(a_n/|\mathcal{V}_n^\circ|)\textstyle\sum_{m=1}^{\ell_n^\circ-1} I_{2,m}^{(3)} < \bar\rho_n^\circ\left[\varsigma_n^{\neq}(u)\right]^2. \tag{9.48}$$

where $\varsigma_n^{\neq}(u)$ is a positive decreasing function of $u>0$ that satisfies

$$\lim_{n\to\infty}\varsigma_n^{\neq}(u) = \nu^\dagger(u,\infty), \quad \forall u>0. \tag{9.49}$$

Equipped with 9.47 and Lemma 9.4 we conclude that under the assumptions and with the notations of Proposition 6.2, Lemma 9.4, and Proposition 6.3, on $\Omega^{\mathsf{SRW}}\cap\overline{\Omega}_{2,n}^{(3)}\cap\Omega^\star$, for all but a finite number of indices n,

$$I_3^{(2)} \le 2\frac{k_n^\circ(t)}{a_n}\left(C''_\circ\bar\sigma_n^=(u,\infty) + \bar\rho_n^\circ\left[\varsigma_n^{\neq}(u)\right]^2\right) \tag{9.50}$$

for some constant $0 < C''_\circ < \infty$. Inserting the bounds (9.41), (9.42), and (9.50) in (9.40) then yields (9.33)-(9.35). The proof of Theorem 9.3 is done. □

9.3. **Almost sure convergence of $\bar\nu_n^\circ$, $\bar\sigma_n^\circ$, and $\bar\sigma_n^=$.** As in Subsection 8.3 our next step consists in proving strong laws of large numbers for the random (but now chain independant) quantities $\bar\nu_n^\circ$, $\bar\sigma_n^\circ$, and $\bar\sigma_n^=$ defined in (9.28), (9.30), and (9.29), respectively. However, the complexity of these objects (note in particular that they are sums of correlated random variables) makes this task much more arduous than in FECP.

Proposition 9.5. *Given $0<\varepsilon<1$ let the sequences a_n and b_n be defined through*

$$\lim_{n\to\infty}\frac{\log a_n}{n\log 2} = \varepsilon, \quad \sqrt{na_n}\mathbb{P}(w_n(x)\ge(n-1)b_n) \sim 1, \tag{9.51}$$

Assume that $c_\star > 2$ and let $\nu^\dagger$ be as in (1.23). Then, there exists a subset $\overline{\Omega}^{LLN} \subset \Omega$ with $\mathbb{P}(\overline{\Omega}^{LLN}) = 1$ such that, on $\overline{\Omega}^{LLN}$, the following holds: for all $u > 0$,

$$\lim_{n\to\infty} \bar\nu_n^\circ(u,\infty) = \nu^\dagger(u,\infty), \tag{9.52}$$

$$\lim_{n\to\infty} n\bar\sigma_n^\circ(u,\infty) = \lim_{n\to\infty} n\bar\sigma_n^=(u,\infty) = 2\nu^\dagger(2u,\infty). \tag{9.53}$$

To prove Proposition 9.5 we first establish control over the mean values of $\bar\nu_n^\circ(u,\infty)$, $\bar\sigma_n^\circ(u,\infty)$, and $\bar\sigma_n^=(u,\infty)$ (see Lemmata 9.8 and 9.10), and then prove that these quantities concentrate around their means (in Lemmata 9.9, 9.11). Both these steps rely on the following key lemma. Given sequences $\bar a_n$, $\bar b_n$, and two distinct vertices $x, y \in \mathcal{V}_n$, set

$$v_n(u;\bar a_n,\bar b_n) = \bar a_n^2 \mathbb{E}\left[\exp\left(-\tfrac{u\bar b_n}{\min\{w_n(x),w_n(y)\}}\right)\mathbb{1}_{\min\{w_n(x),w_n(y)\}\ge r_n(\rho_n^\star)\}}\right]. \tag{9.54}$$

Lemma 9.6. *If the sequences $\bar a_n$ and $\bar b_n$ satisfy $\bar a_n \mathbb{P}(w_n(x) \ge \bar b_n) \sim 1$, $\lim_{n\to\infty} \frac{\bar a_n}{2^n} = 0$, and $\lim_{n\to\infty} \frac{\log \bar a_n}{n\log 2} = \bar\varepsilon$ for some $\bar\varepsilon > 0$, then*

$$\lim_{n\to\infty} v_n(u;\bar a_n,\bar b_n) = u^{-2\alpha(\bar\varepsilon)} 2\alpha(\bar\varepsilon)\Gamma(2\alpha(\bar\varepsilon)). \tag{9.55}$$

The rest of Subsection 9.3 is organized as follows. We prove Lemma 9.6 in Subsection 9.3.1. Equipped with this result we establish the convergence properties of $\bar\nu_n^\circ(u,\infty)$ in Subsection 9.3.2, and those of $\bar\sigma_n^\circ(u,\infty)$ and $\bar\sigma_n^=(u,\infty)$ in Subsection 9.3.3. Subsection 9.3.3 also contains the proof of of Lemma 9.4. The proof of Proposition 9.5 is then completed in Subsection 9.3.4.

9.3.1. *Proof of Lemma 9.6.* The proof uses the following lemma, taken from [25]. Set

$$F_n(v) = \bar a_n \mathbb{P}(w_n(x) \ge v\bar b_n), \quad v \ge 0. \tag{9.56}$$

Lemma 9.7 (Lemma 2.1 of [25])**.** *Let the sequences $\bar a_n$ and $\bar b_n$ be as in Lemma 9.6.*
(i) For each fixed $\zeta > 0$ and all n sufficiently large so that $\zeta > \bar b_n^{-1}$ we have, for all v such that $\zeta \le v < \infty$,

$$F_n(v) = v^{-\alpha_n}(1 + o(1)), \tag{9.57}$$

where $0 \le \alpha_n = \alpha(\bar\varepsilon) + o(1)$.
(ii) Let $0 < \delta < 1$. Then, for all v such that $\bar b_n^{-\delta} \le v \le 1$ and all large enough n,

$$v^{-\alpha_n}(1 + o(1)) \le F_n(v) \le \tfrac{1}{1-\delta} v^{-\alpha_n(1-\frac{\delta}{2})}(1 + o(1)), \tag{9.58}$$

where α_n is as before.

Proof of Lemma 9.6. For fixed $u > 0$ set $f(y) = e^{-u/y}$. Integrating by part in (9.54), and using the fact that $\bar a_n^2 \mathbb{P}\left(\min\{w_n(x), w_n(y)\} \ge y\bar b_n\right) = F_n^2(y)$, we get

$$\mathbb{E}[v_n(u;\bar a_n,\bar b_n)] = F_n^2(r_n(\rho_n^\star)/\bar b_n) e^{-u\bar b_n/r_n(\rho_n^\star)} + \textstyle\int_{r_n(\rho_n^\star)/\bar b_n}^\infty f'(y) F_n^2(y) dy. \tag{9.59}$$

Note that $F_n^2(r_n(\rho_n^\star)/\bar b_n) e^{-u\bar b_n/r_n(\rho_n^\star)} \le \bar a_n^2 e^{-u\bar b_n/r_n(\rho_n^\star)} \downarrow 0$ as $n \uparrow \infty$ for all $u > 0$. To deal with the second term in the r.h.s. of (9.59) set $I_n(a,b) = \int_a^b f'(y) F_n^2(y) dy$, $a \le b$, and, given $0 < \varsigma < 1$ and $\zeta > 1$, write

$$I_n(r_n(\rho_n^\star)/\bar b_n, \infty) = I_n\big(r_n(\rho_n^\star)/\bar b_n, \varsigma\big) + I_n\big(\varsigma, \zeta\big) + I_n(\zeta, \infty). \tag{9.60}$$

Let us now show that, as $n \uparrow \infty$, for small enough ς and large enough ζ, the leading contribution to (9.60) comes from $I_n\big(\varsigma, \zeta\big)$. To do so we first use that by Lemma 9.7, (ii) with $\delta = 1/2$, $I_n\big(r_n(\rho_n^\star)/\bar b_n, \varsigma\big) \le 2(1 + o(1)) \int_0^\varsigma f'(y) y^{-(3/2)\alpha_n} dy$ for all $0 < \varsigma < 1$, where $0 \le \alpha_n = \alpha(\bar\varepsilon) + o(1)$. Now, there exists $\varsigma^* \equiv \varsigma^*(u) > 0$ such that, for all

$\varsigma < \varsigma^*$, $f'(y)y^{-(3/2)\alpha_n}$ is strictly increasing on $[0,\varsigma]$. Hence, for all $\varsigma < \min(1,\varsigma^*)$, $I_n\big(r_n(\rho_n^\star)/\bar b_n,\varsigma\big) \le 2(1+o(1))u\varsigma^{-1+(3/2)[\alpha(\bar\varepsilon)+o(1)]}e^{-u/\varsigma}$, implying that

$$\lim_{n\to\infty} I_n\big(r_n(\rho_n^\star)/\bar b_n,\varsigma\big) \le 2u\varsigma^{-1+(3/2)\alpha(\bar\varepsilon)}e^{-u/\varsigma}\,, \quad \varsigma<\min(1,\varsigma^*)\,. \tag{9.61}$$

To deal with $I_n\big(\varsigma,\zeta\big)$ note that by Lemma 9.7, (i), $F_n(y)\to y^{-\alpha(\bar\varepsilon)}$, $n\to\infty$, where the convergence is uniform in $\varsigma\le y\le\zeta$ since, for each n, $F_n(y)$ is a monotone function, and since the limit, $y^{-\alpha(\bar\varepsilon)}$, is continuous. Hence,

$$\lim_{n\to\infty} I_n\big(\varsigma,\zeta\big) = \lim_{n\to\infty}\int_\varsigma^\zeta f'(y)F_n^2(y)dy = \int_\varsigma^\zeta f'(y)y^{-2\alpha(\bar\varepsilon)}dy\,. \tag{9.62}$$

It remains to bound $I_n(\zeta,\infty)$. By (9.57) of Lemma 9.7, $I_n(\zeta,\infty)=\int_\zeta^\infty f'(y)F_n^2(y)dy = (1+o(1))\int_\zeta^\infty f'(y)y^{-2\alpha_n}dy$, where again $0\le\alpha_n=\alpha(\bar\varepsilon)+o(1)$. Thus, for $0<\delta<1$ arbitrary we have, taking n large enough, that for all $y\ge\zeta>1$, $f'(y)y^{-2\alpha_n}\le f'(y)y^{-2\alpha(\bar\varepsilon)+2\delta}\le u/y^{2(1-\delta)}$. Therefore $I_n(\zeta,\infty)\le(1+o(1))\frac{1}{1-\delta}\zeta^{-(1-2\delta)}$ and, choosing e.g. $\delta=1/4$,

$$\lim_{n\to\infty} I_n(\zeta,\infty) \le 4u\zeta^{-1/2}\,. \tag{9.63}$$

Collecting (9.59)-(9.63), we obtain that for all $\zeta>1$ and $\varsigma<\min(1,\varsigma^*)$,

$$\mathbb{E}[v_n(u;\bar a_n,\bar b_n)] = \int_\varsigma^\zeta f'(y)y^{-2\alpha(\bar\varepsilon)}dy + \mathcal{R}(\varsigma,\zeta)\,, \tag{9.64}$$

where $0\le\mathcal{R}(\varsigma,\zeta)\le 2u\varsigma^{-1+(3/2)\alpha(\bar\varepsilon)}e^{-u/\varsigma}+4u\zeta^{-1/2}$. Finally, passing to the limit $\varsigma\to0$ and $\zeta\to\infty$ in (9.64) yields

$$\mathbb{E}[v_n(u;\bar a_n,\bar b_n)] = \int_0^\infty f'(y)y^{-2\alpha(\bar\varepsilon)}dy = u^{-2\alpha(\bar\varepsilon)}2\alpha(\bar\varepsilon)\Gamma(2\alpha(\bar\varepsilon)), \tag{9.65}$$

where we used in the last equality that $2\alpha(\bar\varepsilon)>0$ since $\bar\varepsilon>0$. Lemma 9.6 is proven. □

9.3.2. *Convergence properties of $\bar\nu_n^\circ$.* As stated in the next two lemmata $\bar\nu_n^\circ$ concentrates around its mean, and the mean as a limit.

Lemma 9.8. *Assume that $c_\star>2$. If a_n and b_n satisfy (9.51) for some $0<\varepsilon<1$, then*

$$\lim_{n\to\infty}\mathbb{E}[\bar\nu_n^\circ(u,\infty)] = \nu^\dagger(u,\infty), \quad \forall u>0. \tag{9.66}$$

Lemma 9.9. *For all $L_1>0$ and $L_2\ge0$ such that $na_nL_2/2^n=o(1)$, for all $u>0$,*

$$\mathbb{P}\big(|\bar\nu_n^\circ(u,\infty)-\mathbb{E}[\bar\nu_n^\circ(u,\infty)]|\ge\phi_n(u,L_1,L_2)\big)\le 2ne^{-L_2}+2L_1, \tag{9.67}$$

where $\phi_n(u,L_1,L_2)\equiv\sqrt{8na_nL_2V_{n,2}(2u)/2^n}+4n^{-(c_\star-1)}V_{n,1}(u)/L_1$, and where for each n, $V_{n,1}(u)$ and $V_{n,2}(u)$ are positive decreasing functions, while for each $u>0$, under the assumptions of Lemma 9.8,

$$\lim_{n\to\infty}V_{n,1}(u)=\lim_{n\to\infty}V_{n,2}(u)=\nu^\dagger(u,\infty). \tag{9.68}$$

Proof of Lemma 9.8. Write $\bar\nu_n^\circ(u,\infty)=\sum_{k\ge2}\bar\nu_n^{\circ,(k)}(u,\infty)$ where

$$\bar\nu_n^{\circ,(k)}(u,\infty)\equiv\frac{a_n}{2^n}\sum_{1\le l\le L^\star}\mathbb{1}_{\{|C_{n,l}^\star|=k\}}\sum_{x\in C_{n,l}^\star}Q_{n,l}^u(x). \tag{9.69}$$

We saw in Subsection 7.1 (see (7.7)) that on $\Omega^\star$, $k_n^\star\equiv\max_{2\le l\le L^\star}|C_{n,l}^\star(\rho)|\le\frac{n}{(c_\star-2)\log n}$ for all large enough n. We may thus restrict the range of k to $2\le k\le k_n^\star$. Let now $\mathcal{G}_k$ be the collection of all vertex sets $\mathcal{C}\subset\mathcal{V}_n$ of size k such that $G(\mathcal{C})$ forms a connected subgraph of $G(\mathcal{V}_n)$,

$$\mathcal{G}_k=\{\mathcal{C}=\{x_1,\dots,x_k\}:\forall_{1\le i\le k}\exists_{1\le j\le k}\ \text{such that}\ \ \mathrm{dist}(x_i,x_j)=1\}. \tag{9.70}$$

Then (9.69) can be written as

$$\bar\nu_n^{\circ,(k)}(u,\infty) \equiv \frac{a_n}{2^n}\sum_{\mathcal{C}\in\mathcal{G}_k}\prod_{x\in\mathcal{C}}\chi_n(x)\prod_{x'\in\partial\mathcal{C}}\overline{\chi}_n(x')\sum_{x\in\mathcal{C}}Q^u_{n,\mathcal{C}}(x), \tag{9.71}$$

where $Q^u_{n,\mathcal{C}}(x)$ stands for $Q^u_{n,l}(x)$ with $C^\star_{n,l}\equiv\mathcal{C}$, and where $\chi_n(x)\equiv\mathbb{1}_{\{w_n(x)\geq r_n(\rho^\star_n)\}}$, $\overline{\chi}_n(x)\equiv 1-\chi_n(x)$, are Bernoulli variables r.v.'s with $\mathbb{P}\left(\chi_n(x)=1\right)=n^{-c_\star}$. To further express $\bar\nu_n^{\circ,(k)}$ we distinguish the case $k=2$ from the case $3\leq k\leq k^\star_n$.

• ***The case*** $k=2$**.** Here $\mathcal{G}_2$ is the set of undirected edges of $G(\mathcal{V}_n)$ and $Q^u_{n,\mathcal{C}}(x)$ is given by Proposition 5.1, (i) : observing that $Q^u_{n,\mathcal{C}}(x)=Q^u_{n,\mathcal{C}}(y)$ on $\mathcal{C}=\{x,y\}$, and that, by (5.2), $\varrho_{n,l}(0)=\min_{x\in\mathcal{C}}w_n(x)$ when $C^\star_{n,l}=\mathcal{C}$, we obtain

$$\bar\nu_n^{\circ,(2)}(u,\infty)\equiv 2\frac{a_n}{2^n}\sum_{\mathcal{C}\in\mathcal{G}_2}\prod_{x\in\mathcal{C}}\chi_n(x)\prod_{x'\in\partial\mathcal{C}}\overline{\chi}_n(x')\left(1-\frac{1}{1+\frac{\min_{x\in\mathcal{C}}w_n(x)}{(n-1)}}\right)^{\lceil b_n u\rceil}. \tag{9.72}$$

From this we easily derive the bounds

$$\bar\nu_n^{\circ,(2),-}(u,\infty)(1-s_n)\leq\bar\nu_n^{\circ,(2)}(u,\infty)\leq\bar\nu_n^{\circ,(2),+}(u,\infty) \tag{9.73}$$

where $s_n=\frac{n-1}{r_n(\rho^\star_n)}$ and where, setting $b_n^\pm=b_n(n-1)(1-s_n)^{\pm1}$ and

$$\gamma_n^\pm(\mathcal{C})=\min_{x\in\mathcal{C}}w_n(x)/b_n^\pm, \tag{9.74}$$

$$\bar\nu_n^{\circ,(2),\pm}(u,\infty)\equiv 2\frac{a_n}{2^n}\sum_{\mathcal{C}\in\mathcal{G}_2}\prod_{x'\in\partial\mathcal{C}}\overline{\chi}_n(x')e^{-u/\gamma_n^\pm(\mathcal{C})}\mathbb{1}_{\{\gamma_n^\pm(\mathcal{C})\geq r_n(\rho^\star_n)/b_n^\pm\}}. \tag{9.75}$$

By Lemma 9.6,

$$\lim_{n\to\infty}\mathbb{E}\bar\nu_n^{\circ,(2),-}(u,\infty)=\lim_{n\to\infty}\mathbb{E}\bar\nu_n^{\circ,(2),+}(u,\infty)=\nu^\dagger(u,\infty). \tag{9.76}$$

To see this note that, setting $a_n^+=a_n^-=\sqrt{na_n}(1-n^{-c_\star})^{n-1}$,

$$\mathbb{E}[\bar\nu_n^{\circ,(2),\pm}(u,\infty)]=v_n(u;a_n^\pm,b_n^\pm). \tag{9.77}$$

One then readily checks that for a_n, b_n as in (9.51), $\lim_{n\to\infty}\frac{\log a_n^\pm}{n\log 2}=\varepsilon/2$, $\lim_{n\to\infty}\frac{a_n^\pm}{2^n}=0$, and $a_n^\pm\mathbb{P}(w_n(x)\geq b_n^\pm)\sim 1$. The conditions of Lemma 9.6 are thus satisfied with $\bar\varepsilon=\varepsilon/2$, yielding (9.76). Since clearly $\lim_{n\to\infty}s_n=0$, it follows from (9.73) that

$$\lim_{n\to\infty}\mathbb{E}\bar\nu_n^{\circ,(2)}(u,\infty)=\nu^\dagger(u,\infty). \tag{9.78}$$

• ***The case*** $3\leq k\leq k^\star_n$**.** Recall that $Q^u_{n,\mathcal{C}}(x)$ in (9.71) stands for $Q^u_{n,l}(x)$ with $C^\star_{n,l}\equiv\mathcal{C}$. Similarly, denote by $\varrho_{n,\mathcal{C}}(0)$, $\varrho_{n,\mathcal{C}}(1)$, and $\theta^\star_{n,\mathcal{C}}$ the quantities $\varrho_{n,l}(0)$, $\varrho_{n,l}(1)$, and $\theta^\star_{n,l}$ from (5.2)-(5.4) with $C^\star_{n,l}\equiv\mathcal{C}$. Let us now split the sum over $\mathcal{C}\in\mathcal{G}_k$ in (9.71) according to whether $b_n<n\theta^\star_{n,\mathcal{C}}$ or not, and write $\bar\nu_n^{\circ,(k)}(u,\infty)=\bar\nu_n^{\circ,(k),-}(u,\infty)+\bar\nu_n^{\circ,(k),+}(u,\infty)$ where

$$\bar\nu_n^{\circ,(k),+}(u,\infty)\equiv\frac{a_n}{2^n}\sum_{\mathcal{C}\in\mathcal{G}_k}\prod_{x\in\mathcal{C}}\chi_n(x)\prod_{x'\in\partial\mathcal{C}}\overline{\chi}_n(x')\sum_{x\in\mathcal{C}}Q^u_{n,\mathcal{C}}(x)\mathbb{1}_{\{b_n\geq n\theta^\star_{n,\mathcal{C}}\}}, \tag{9.79}$$

and $\bar\nu_n^{\circ,(k),-}(u,\infty)$ is the sum over the complement, i.e. over terms such that $b_n<n\theta^\star_{n,\mathcal{C}}$. The point of doing this is that if $b_n\geq n\theta^\star_{n,\mathcal{C}}$ then, for each fixed $u>0$, $\lceil b_nu\rceil\gg\theta^\star_{n,\mathcal{C}}$ for all x in $\mathcal{C}$ and all n large enough, so that $\bar\nu_n^{\circ,(k),+}(u,\infty)$ can be bounded using Proposition 5.1, (ii). More precisely, on $\Omega_0\cap\Omega^\star$, for all but a finite number of indices n, by (5.6),

$$Q^u_{n,\mathcal{C}}(x)\mathbb{1}_{\{b_n\geq n\theta^\star_{n,\mathcal{C}}\}}\leq e^{-\{u(n-1)b_n/\varrho_{n,\mathcal{C}}(0)\}}(1+o(1)),\quad\forall x\in\mathcal{C}, \tag{9.80}$$

(since for $k \geq 3$ and large enough n, $k(n-1)/n(1-o(1)) > 1$). Note that by (5.2),

$$e^{-\{u(n-1)b_n/\varrho_{n,\mathcal{C}}(0)\}} = \max_{\{x,y\}\in G(\mathcal{C})} e^{-\{u(n-1)b_n/\min(w_n(y),w_n(x))\}} \quad (9.81)$$

$$\leq \sum_{\{x,y\}\in G(\mathcal{C})} e^{-u/\bar{\gamma}_n(\{x,y\})}, \quad (9.82)$$

where we now set

$$\bar{\gamma}_n(\mathcal{C}') = \min_{x\in\mathcal{C}'} w_n(x)/(n-1)b_n, \quad \mathcal{C}' \in \mathcal{G}_2. \quad (9.83)$$

Combining these observations yields the bound

$$\bar{\nu}_n^{\circ,(k),+}(u,\infty) \leq k\frac{a_n}{2^n} \sum_{\mathcal{C}'\in\mathcal{G}_2} \sum_{\mathcal{C}\in\mathcal{G}_k:\mathcal{C}'\subset\mathcal{C}} \prod_{x\in\mathcal{C}} \chi_n(x) e^{-u/\bar{\gamma}_n(\mathcal{C}')}, \quad (9.84)$$

valid on $\Omega_0 \cap \Omega^\star$, for all but a finite number of indices n, and this in turns implies that

$$\mathbb{E}[\bar{\nu}_n^{\circ,(k),+}(u,\infty)] \leq k(k-2)! n^{-(c_\star-1)(k-2)} v_n(u;\bar{a}_n,\bar{b}_n), \quad (9.85)$$

where $\bar{a}_n \equiv \sqrt{na_n}$, $\bar{b}_n \equiv (n-1)b_n$. Again one sees that these sequences (that differ but slightly from the choices made in (9.77)) satisfy the conditions of Lemma 9.6 with $\bar{\varepsilon} = \varepsilon/2$. Thus

$$\lim_{n\to\infty} \mathbb{E}\left[v_n(u;\bar{a}_n,\bar{b}_n)\right] = \nu^\dagger(u,\infty). \quad (9.86)$$

Since by assumption $c_\star > 2$ we may use (7.24) to sum (9.85) over k, which gives

$$\sum_{3\leq k\leq k_n^\star} \mathbb{E}[\bar{\nu}_n^{\circ,(k),+}(u,\infty)] \leq 4n^{-(c_\star-1)} \mathbb{E}\left[v_n(u;\bar{a}_n,\bar{b}_n)\right]. \quad (9.87)$$

It remains to bound the term $\bar{\nu}_n^{\circ,(k),-}(u,\infty)$ (see the line above (9.79)). For this we use the trite bound $Q^u_{n,\mathcal{C}}(x) \leq 1$ and observe that, by (5.2)-(5.4) and the rightmost inequality in Lemma 4.5, $\mathbb{1}_{\{b_n<n\theta^\star_{n,\mathcal{C}}\}} \prod_{x\in\mathcal{C}} \chi_n(x) \leq \mathbb{1}_{\{\varrho_{n,\mathcal{C}}(0)>r_n(\rho_n^\star)b_n/(2\beta n^2k^5)\}} \prod_{x\in\mathcal{C}} \chi_n(x)$. Next, by (5.2) and (9.83),

$$\{\varrho_{n,\mathcal{C}}(0) > r_n(\rho_n^\star)b_n/(2\beta n^2k^5)\} \subseteq \cup_{\{x,y\}\in G(\mathcal{C})}\{\bar{\gamma}_n(\{x,y\}) > r_n(\rho_n^\star)/\beta n^8\} \quad (9.88)$$

for large enough n. Thus

$$\bar{\nu}_n^{\circ,(k),-}(u,\infty) \leq k\frac{a_n}{2^n} \sum_{\mathcal{C}'\in\mathcal{G}_2} \sum_{\mathcal{C}\in\mathcal{G}_k:\mathcal{C}'\subset\mathcal{C}} \prod_{x\in\mathcal{C}} \chi_n(x) \mathbb{1}_{\{\bar{\gamma}_n(\mathcal{C}')>r_n(\rho_n^\star)/\beta n^8\}}, \quad (9.89)$$

and averaging out,

$$\mathbb{E}[\bar{\nu}_n^{\circ,(k),-}(u,\infty)] \leq k(k-2)! n^{-(c_\star-1)(k-2)} \left[\bar{a}_n \mathbb{P}(w_n(0) > \bar{b}_n r_n(\rho_n^\star)/\beta n^8)\right]^2, \quad (9.90)$$

where $\bar{a}_n$, $\bar{b}_n$ are as before (see the line below (9.85)). A simple Gaussian tail estimate gives $\left[\bar{a}_n \mathbb{P}(w_n(0) > \bar{b}_n r_n(\rho_n^\star)/\beta n^8)\right]^2 \leq (\beta n^8/r_n(\rho_n^\star))^{2\alpha(\varepsilon/2)(1+o(1))}$. Again, the assumption that $c_\star > 2$ enables us to use (7.24), yielding

$$0 \leq \sum_{3\leq k\leq k_n^\star} \mathbb{E}[\bar{\nu}_n^{\circ,(k),-}(u,\infty)] \leq 4n^{-(c_\star-1)} \left(\beta n^8/r_n(\rho_n^\star)\right)^{2\alpha(\varepsilon/2)(1+o(1))}. \quad (9.91)$$

Now set

$$\Delta_n(u) \equiv \sum_{k\geq 3} \bar{\nu}_n^{\circ,(k)}(u,\infty) = \hat{\nu}_n^\circ(u,\infty) - \bar{\nu}_n^{\circ,(2)}(u,\infty) > 0. \quad (9.92)$$

Combining (9.87), (9.91), and using (2.20) to bound $r_n(\rho_n^\star)$, we obtain that under the assumptions of the lemma, for all $u > 0$,

$$0 \leq \mathbb{E}\Delta_n(u) \leq 4n^{-(c_\star-1)} \left(\mathbb{E}\left[v_n(u;\bar{a}_n,\bar{b}_n)\right] + e^{-\beta\sqrt{8\varepsilon n\log n}}\right), \quad (9.93)$$

and so $\lim_{n\to\infty} \mathbb{E}\Delta_n(u) = 0$. But this and (9.78) yield (9.66). The proof of Lemma 9.8 is complete. □

Proof of Lemma 9.9. As in the proof of Lemma 9.8 we separate the contribution of $\bar\nu_n^{\circ,(2)}$ from those of $\bar\nu_n^{\circ,(k)}$, $k \geq 3$ (see (9.71) and (9.72) for their definitions). Namely, we write

$$\bar\nu_n^\circ(u,\infty) - \mathbb{E}[\bar\nu_n^\circ(u,\infty)] = \bar\nu_n^{\circ,(2)}(u,\infty) - \mathbb{E}[\bar\nu_n^{\circ,(2)}(u,\infty)] + \{\Delta_n(u) - \mathbb{E}[\Delta_n(u)]\} \quad (9.94)$$

where $\Delta_n(u)$ is defined in (9.92), and we take

$$V_{n,1}(u) \equiv \mathbb{E}\left[v_n(u;\bar a_n,\bar b_n)\right] + e^{-\beta\sqrt{8\varepsilon n\log n}}, \quad (9.95)$$

$$V_{n,2}(u) \equiv \mathbb{E}\left[v_n(u;a_n^+,b_n^+)\right], \quad (9.96)$$

where $v_n(u;\bar a_n,\bar b_n)$ and $v_n(u;a_n^+,b_n^+)$ are as in (9.93) and (9.77), respectively. Eq. (9.68) then follows from (9.86) and (9.78). Note that $\Delta_n(u) > 0$. Thus, by (9.93),

$$\mathbb{P}\left(|\Delta_n(u) - \mathbb{E}[\Delta_n(u)]| \geq 4n^{-(c_\star-1)}V_{n,1}(u)/L_1\right) \leq 2L_1, \quad \forall L_1 > 0. \quad (9.97)$$

Let us now establish that for all $L_2 \geq 0$ such that $na_nL_2/2^n = o(1)$,

$$\mathbb{P}\left(\left|\bar\nu_n^{\circ,(2)}(u,\infty) - \mathbb{E}[\bar\nu_n^{\circ,(2)}(u,\infty)]\right| \geq \sqrt{8na_nL_2V_{n,2}(2u)/2^n}\right) \leq e^{-L_2}. \quad (9.98)$$

Eq. (9.72) prompts us to set $\mathcal{S}_n \equiv \sum_{\mathcal{C}\in\mathcal{G}_2} X(\mathcal{C})$, where $X(\mathcal{C}) \equiv Y(\mathcal{C}) - \mathbb{E}Y(\mathcal{C})$ and

$$Y(\mathcal{C}) \equiv \prod_{x\in\mathcal{C}}\chi_n(x)\prod_{x'\in\partial\mathcal{C}}\overline{\chi}_n(x')\Big(1 - \Big[1 + \tfrac{\min_{x\in\mathcal{C}} w_n(x)}{(n-1)}\Big]^{-1}\Big)^{\lceil b_nu\rceil}, \quad \mathcal{C}\in\mathcal{G}_2. \quad (9.99)$$

Then

$$\mathbb{P}\left(\left|\bar\nu_n^{\circ,(2)}(u,\infty) - \mathbb{E}[\bar\nu_n^{\circ,(2)}(u,\infty)]\right| \geq \theta\right) = \mathbb{P}\left(|\mathcal{S}_n| \geq 2^{n-1}a_n^{-1}\theta\right). \quad (9.100)$$

Observe that $\mathcal{S}_n$ is a sum of dependent random variables. To get round this difficulty we split it into $2n$ disjoint sums as follows. Write $\mathcal{G}_2 = \cup_{1\leq j\leq n}\mathcal{G}_2^j$ where, for each $1 \leq j \leq n$, $\mathcal{G}_2^j \equiv \{\{x,y\}\in\mathcal{G}_2 \mid x_j = -y_j\}$ is the set of neighboring vertices that differ in exactly the j-th coordinate. Subdivide each $\mathcal{G}_2^j$ into 4 disjoint sets, $\mathcal{G}_2^j = \cup_{1\leq i\leq 4}\mathcal{G}_2^{j,i}$,

$$\mathcal{G}_2^{j,i} \equiv \left\{\{x,y\}\in\mathcal{G}_2^j \mid \mathrm{dist}(\mathbf{1},\{x,y\}) = 4m + i - 1, m \geq 0\right\}, \quad (9.101)$$

where $\mathbf{1}$ denotes the vertex of $\mathcal{V}_n$ all of whose coordinates are 1. Then,

$$\mathcal{S}_n = \sum_{i=1}^4\sum_{j=1}^n \mathcal{S}_n^{j,i}, \quad \mathcal{S}_n^{j,i} \equiv \sum_{\mathcal{C}\in\mathcal{G}_2^{j,i}} X(\mathcal{C}). \quad (9.102)$$

Each $\mathcal{S}_n^{j,i}$ now is a sum of independent random variables, and can be controlled using Bennett's bound [9] for the tail behavior of sums of random variables, which we specialize as follows: if $(X(i),\, i\in\mathcal{I})$ is a family of i.i.d. centered random variables that satisfies $\max_i|X(i)| \leq A$ then, setting $\tilde B^2 = \sum_{i\in\mathcal{I}}\mathbb{E}X^2(i)$, for all $B^2 \geq \tilde B^2$, for all $t < B^2/(2A)$,

$$\mathbb{P}\left(\left|\sum_{i\in\mathcal{I}}X(i)\right| \geq t\right) \leq \exp\left\{-t^2/4B^2\right\}. \quad (9.103)$$

Since $|X(\mathcal{C})| \leq 1$ and since $\sum_{\mathcal{C}\in\mathcal{G}_2^{j,i}}\mathbb{E}X^2(\mathcal{C}) \leq (4na_n)^{-1}2^{n-1}V_{n,2}(2u)$, as follows from (9.73)-(9.77), we may choose $A = 1$ and $B^2 = (4na_n)^{-1}2^{n-1}V_{n,2}(2u)$ in (9.103), and we get that for all $L_2 > 0$,

$$\mathbb{P}\left(\left|\mathcal{S}_n^{j\pm}\right| \geq 2^{n-1}(4na_n)^{-1}\theta\right) = \mathbb{P}\left(|\mathcal{S}_n^{j\pm}| \geq \sqrt{2^{n-1}L_2V_{n,2}(2u)/(4na_n)}\right) \leq e^{-L_2}, \quad (9.104)$$

where we chose $\theta^2 = 4na_nL_22^{-n+1}V_{n,2}(2u)$. This choice is permissible provided that $\theta \leq V_{n,2}(2u)/2$. In view of (9.76) this will be verified for all n large enough whenever $\theta\downarrow 0$ as $n\uparrow\infty$, i.e. whenever $na_nL_2/2^n = o(1)$. Eq. (9.104) holds true for each $1 \leq j \leq n$ and $1 \leq i \leq 4$, and combined with (9.102) yields

$$\mathbb{P}\left(|\mathcal{S}_n| \geq 4n\sqrt{2^{n-1}L_2V_{n,2}(2u)/(4na_n)}\right) \leq 4ne^{-L_2}, \quad (9.105)$$

which, by (9.100), is tantamount to (9.98). Combining (9.97) and (9.98) then yields (9.67) and concludes the proof of Lemma 9.9. □

9.3.3. *Convergence properties of $\bar\sigma_n^\circ$ and related functions.* We have:

Lemma 9.10. *Under the assumption and with the notation of Lemma 9.8,*

$$\lim_{n\to\infty} n\mathbb{E}[\bar\sigma_n^\circ(u,\infty)] = \lim_{n\to\infty} n\mathbb{E}[\bar\sigma_n^=(u,\infty)] = 2\nu^\dagger(2u,\infty), \quad \forall u>0. \tag{9.106}$$

Lemma 9.11. *For all $L_1, L_3 > 0$ and $L_2 \geq 0$ such that $na_nL_2/2^n = o(1)$, for all $u>0$,*

$$\mathbb{P}\big(|\bar\sigma_n^=(u,\infty) - \mathbb{E}[\bar\sigma_n^=(u,\infty)]| \geq \tilde\phi_n(u,L_1,L_2)\big) \leq 2ne^{-L_2} + 4L_1, \tag{9.107}$$

$$\mathbb{P}\big(|\bar\sigma_n^\circ(u,\infty) - \mathbb{E}[\bar\sigma_n^\circ(u,\infty)]| \geq \psi_n(u,L_1,L_2,L_3)\big) \leq 2ne^{-L_2} + 2L_1 + 2L_3, \tag{9.108}$$

where $\tilde\phi_n(u,L_1,L_2) \equiv \sqrt{a_nL_2W_{n,2}(2u)/2^n} + 4n^{-c_\star}W_{n,1}(u)/L_1$, $\psi_n(u,L_1,L_2,L_3) \equiv \tilde\phi_n(u,L_1,L_2) + 2^8W_{n,3}^2(u)/(a_nL_3)$, and where for each n and $1\leq i\leq 3$, $W_{n,i}(u)$ are positive decreasing functions, while for each $u>0$, under the assumptions of Lemma 9.8,

$$\lim_{n\to\infty} W_{n,1}(u) = \lim_{n\to\infty} W_{n,2}(u) = 2\nu^\dagger(2u,\infty), \tag{9.109}$$

$$\lim_{n\to\infty} W_{n,3}^2(u) = [\nu^\dagger(u,\infty)]^2. \tag{9.110}$$

We prove Lemmata 9.10 and 9.11 simultaneously.

Proof of Lemma 9.10 and Lemma 9.11. Write $\bar\sigma_n^\circ(u,\infty) = \bar\sigma_n^=(u,\infty) + \bar\sigma_n^{\neq}(u,\infty)$ where

$$\bar\sigma_n^=(u,\infty) = \frac{a_n}{n2^n}\sum_{1\leq l\leq L^\star}\Big[\sum_{x\in C_{n,l}^\star} Q_{n,l}^u(x)\Big]^2, \tag{9.111}$$

$$\bar\sigma_n^{\neq}(u,\infty) = \frac{a_n}{n^22^n}\sum_{1\leq l\neq l'\leq L^\star}\sum_{x\in C_{n,l}^\star}\sum_{x'\in C_{n,l'}^\star} Q_{n,l}^u(x)Q_{n,l'}^u(x')|\partial x\cap\partial x'|. \tag{9.112}$$

Comparing (9.111) to (9.28), we see that $n\bar\sigma_n^=(u,\infty)$ differs from $\bar\nu_n^\circ(u,\infty)$ in that the term in square brackets is squared. However, examining the proof of 9.8 (see (9.72)-(9.75) and (9.79)-(9.84)) we also see that $n\bar\sigma_n^=(u,\infty)$ can be controlled in exactly the same way as $\bar\nu_n^\circ(u,\infty)$, substituting $|C_{n,l}^\star|^2Q_{n,l}^{2u}(x)$ for $[\sum_{x\in C_{n,l}^\star} Q_{n,l}^u(x)]^2$. This yields

$$\lim_{n\to\infty} n\mathbb{E}[\bar\sigma_n^=(u,\infty)] = 2\nu^\dagger(2u,\infty), \quad \forall u>0. \tag{9.113}$$

Similarly, a rerun of the proof of Lemma 9.9 yields, under the same assumptions on L_1, L_2,

$$\mathbb{P}\big(|\bar\sigma_n^=(u,\infty) - \mathbb{E}[\bar\sigma_n^=(u,\infty)]| \geq \phi_n(u,L_1,L_2)\big) \leq 2ne^{-L_2} + 4L_1, \tag{9.114}$$

where $\phi_n(u,L_1,L_2) \equiv \sqrt{a_nL_2W_{n,2}(2u)/2^n} + 4n^{-c_\star}W_{n,1}(u)/L_1$, and where $W_{n,1}(u)$ and $W_{n,2}(u)$ enjoy the same properties, under the same assumptions, as $V_{n,1}(u)$ and $V_{n,2}(u)$, but with (9.68) replaced by (9.109).

Let us now establish that

$$\mathbb{E}[\bar\sigma_n^{\neq}(u,\infty)] \leq 2^8a_n^{-1}[W_{n,3}(u)]^2 \tag{9.115}$$

for some positive decreasing function $W_{n,3}(u)$ of $u>0$, that satisfies

$$\lim_{n\to\infty} W_{n,3}(u) = \nu^\dagger(u,\infty), \quad \forall u>0. \tag{9.116}$$

For this write $\bar\sigma_n^{\neq}(u,\infty) = \sum_{2\leq k,k'\leq k_n^\star}\hat\sigma_n^{\neq,(k,k')}(u,\infty)$ where, with the notation of (9.71),

$$\hat\sigma_n^{\neq,(k,k')}(u,\infty) \equiv \frac{a_n}{n^22^n}\sum_{\mathcal{C},\mathcal{C}'}^{(1)}\phi(\mathcal{C},\mathcal{C}')\sum_{x,x'}^{(2)} Q_{n,\mathcal{C}}^u(x)Q_{n,\mathcal{C}'}^u(x'). \tag{9.117}$$

Here the first sum, $\Sigma^{(1)}$, is over all $\mathcal{C} \in \mathcal{G}_k$ and $\mathcal{C}' \in \mathcal{G}_{k'}$ such that $\mathrm{dist}(\mathcal{C}, \mathcal{C}') = 2$, the second one, $\Sigma^{(2)}$, is over all $x \in \mathcal{C}$ and $x' \in \mathcal{C}'$ such that $\mathrm{dist}(x, x') = 2$, and $\phi(\mathcal{C}, \mathcal{C}') \equiv \prod_{y \in \mathcal{C} \cup \mathcal{C}'} \chi_n(y) \prod_{y' \in \partial\mathcal{C} \cup \partial\mathcal{C}'} \overline{\chi}_n(y')$. Thus $\mathcal{C} \cap \mathcal{C}' = \emptyset$, so that that $Q^u_{n,\mathcal{C}}(x)$ and $Q^u_{n,\mathcal{C}'}(x')$ are independant random variables for all $x \in \mathcal{C}$, $x' \in \mathcal{C}'$, and averaging out,

$$\mathbb{E} \sum_{2 \le k,k' \le k^\star_n} \hat{\sigma}^{\neq,(k,k')}_n(u, \infty) \le a_n^{-1} [W_{n,3}(u)]^2 s^{(3)}_n s^{(4)}_n, \tag{9.118}$$

where

$$W_{n,3}(u) \equiv \mathbb{E}\left[v_n(u; \bar{a}_n, \bar{b}_n)\right] + e^{-\beta\sqrt{8\varepsilon n \log n}} \tag{9.119}$$

for some sequences $\bar{a}_n$, $\bar{b}_n$ chosen as in Lemma 9.6, and where

$$s^{(i)}_n = \sum_{2 \le k \le k^\star_n} k^i (k-1)! n^{-(c_\star - 1)(k-2)}, \quad i \ge 1. \tag{9.120}$$

To see this, reason that there are at most $2^n(k-1)!n^{k-1}$ sets $\mathcal{C} \in \mathcal{G}_k$, that for each $\mathcal{C} \in \mathcal{G}_k$ there are at most $n^2 k'(k'-1)! n^{k'-1}$ sets $\mathcal{C}' \in \mathcal{G}_{k'}$ such that $\mathrm{dist}(\mathcal{C}, \mathcal{C}') = 2$, that $\Sigma^{(2)}$ contains at most kk' terms, and that, proceeding as in (9.80)-(9.84), respectively (9.89)-(9.91), to bound the terms $Q^u_{n,\mathcal{C}}$ when $k > 2$, depending on whether $b_n < n\theta^\star_{n,\mathcal{C}}$ or $b_n \ge n\theta^\star_{n,\mathcal{C}}$, and proceeding as in (9.73)-(9.76) when $k = 2$, we have

$$\mathbb{E}\left[\phi(\mathcal{C}, \mathcal{C}') Q^u_{n,\mathcal{C}}(x) Q^u_{n,\mathcal{C}'}(x')\right] \le (kk')^2 (na_n)^{-2} n^{-c_\star[(k-2)+(k'-2)]} [W_{n,3}(u)]^2, \tag{9.121}$$

for some sequences $\bar{a}_n$, $\bar{b}_n$ for which the assumptions of Lemma 9.6 are verified, and all $k, k' \ge 2$. Now for $c_\star > 2$, by (7.24), $s^{(i)}_n \le 2^i(1 + 2^{i+1} n^{-2(c_\star - 1)})$. Inserting this in (9.118) and using Lemma 9.6 proves the claim (9.115)-(9.116). This immediately implies that

$$\lim_{n \to \infty} n\mathbb{E}[\bar{\sigma}^{\neq}_n(u, \infty)] = 0, \quad \forall u > 0, \tag{9.122}$$

and that, under the assumptions and with the notation of (9.115)-(9.116), for all $u > 0$,

$$\mathbb{P}\left(\left|\bar{\sigma}^{\neq}_n(u, \infty) - \mathbb{E}[\bar{\sigma}^{\neq}_n(u, \infty)]\right| \ge 2^8 [W_{n,3}(u)]^2 / (a_n L_3)\right) \le 2L_3. \tag{9.123}$$

Lemma 9.10 now follows from (9.113) and (9.122), and Lemma 9.11 follows from (9.114), (9.116), and (9.123). □

We now prove Lemma 9.4.

Proof of Lemma 9.4. Let us establish first that for all $m \ge 1$, if $c_\star > 2$,

$$\mathbb{E}\big[I^{(3)}_{2,m}\big] \le n^{-1} a_n^{-2} 2^n [W_{n,3}(u)]^2 2^5 (1 + 2^4 n^{-2(c_\star - 1)})^2, \tag{9.124}$$

where $W_{n,3}(u) > 0$ is a decreasing function satisfying $\lim_{n \to \infty} W_{n,3}(u) = \nu^\dagger(u, \infty)$ for all $u > 0$. For this note that $I^{(3)}_{2,m}$ in (9.45) is very similar to the quantity $\bar{\sigma}^{\neq}_n(u, \infty)$ defined in (9.112). This prompts us to write

$$I^{(3)}_{2,m} = \sum_{2 \le k,k' \le k^\star_n} I^{(3),(k,k')}_{2,m} \tag{9.125}$$

where, for $\phi(\mathcal{C}, \mathcal{C}')$ as in (9.117),

$$I^{(3),(k,k')}_{2,m} \equiv \sum^{(1)}_{\mathcal{C},\mathcal{C}'} \phi(\mathcal{C}, \mathcal{C}') \sum^{(2)}_{x,x'} \sum^{(3)}_{y,y'} Q^u_{n,l}(x) Q^u_{n,l'}(x') f^{\circ,m}_n(x, x'; y, y'). \tag{9.126}$$

Here the first sum, $\Sigma^{(1)}$, is over all $\mathcal{C} \in \mathcal{G}_k$ and $\mathcal{C}' \in \mathcal{G}_{k'}$ such that $\mathcal{C} \cap \mathcal{C}' = \emptyset$, the second one, $\Sigma^{(2)}$, is over all $x \in \mathcal{C}$ and $x' \in \mathcal{C}'$, and the third one, $\Sigma^{(3)}$, is over all $y \in \partial\mathcal{C}$ and $y \in \partial\mathcal{C}'$. Since $\mathcal{C} \cap \mathcal{C}' = \emptyset$, $Q^u_{n,\mathcal{C}}(x)$ and $Q^u_{n,\mathcal{C}'}(x')$ are independant random variables for all $x \in \mathcal{C}$, $x' \in \mathcal{C}'$. Thus we see, using (9.121), that for all $k, k' \ge 2$,

$$\begin{aligned} &\mathbb{E}\big[I^{(3),(k,k')}_{2,m}\big] \\ &\le (kk')^2 (na_n)^{-2} n^{-c_\star[(k-2)+(k'-2)]} [W_{n,3}(u)]^2 \textstyle\sum^{(1)}_{\mathcal{C},\mathcal{C}'} \sum^{(2)}_{x,x'} \sum^{(3)}_{y,y'} f^{\circ,m}_n(x, x'; y, y'), \end{aligned} \tag{9.127}$$

where $W_{n,3}(u)$ is given by (9.119) for some sequences $\bar{a}_n$, $\bar{b}_n$ for which the assumptions of Lemma 9.6 are verified – hence it has the properties claimed in the line below (9.124). To deal with the sums in (9.127), observe that given any $\mathcal{C} \in \mathcal{G}_k$, $x \in \mathcal{C}$, and $y \in \partial\mathcal{C}$,

$$f_n^{\circ,m}(x;y) \equiv \sum_{\mathcal{C}' \in \mathcal{G}_{k'}} \sum_{x' \in \mathcal{C}'} \sum_{y' \in \partial\mathcal{C}'} f_n^{\circ,m}(x,x';y,y') \tag{9.128}$$

$$= p_n(x,y) \sum_{y' \in \mathcal{V}_n^\circ} \sum_{\mathcal{C}'}^{(4)} \sum_{x' \in \mathcal{C}'} p_n(y',x') p_n^{\circ,m}(y,y') \tag{9.129}$$

where the sum $\Sigma^{(4)}$ is over all $\mathcal{C}' \in \mathcal{G}_{k'}$ such that $\mathcal{C}' \cap \partial y' \neq \emptyset$. Indeed if $\mathcal{C}' \cap \partial y' = \emptyset$ then $p_n(y',x') = 0$ for all $x' \in \mathcal{C}'$. Now $\sum_{x' \in \mathcal{C}'} p_n(y',x') \leq 1$ while the number of terms in $\Sigma^{(4)}$ is at most $k'! n^{k'}$. Thus

$$f_n^{\circ,m}(x;y) \leq k'! n^{k'} p_n(x,y) \sum_{y' \in \mathcal{V}_n^\circ} p_n^{\circ,m}(y,y') \leq k'! n^{k'} p_n(x,y), \tag{9.130}$$

From this we readily get

$$\sum_{\mathcal{C},\mathcal{C}'}^{(1)} \sum_{x,x'}^{(2)} \sum_{y,y'}^{(3)} f_n^{\circ,m}(x,x';y,y') \leq (k-1)! n^{k-1} k'! n^{k'}, \tag{9.131}$$

and inserting this bound in (9.127) and (9.125) successively yields

$$\mathbb{E}\big[I_{2,m}^{(3)}\big] \leq n^{-1} a_n^{-2} 2^n [W_{n,3}(u)]^2 s_n^{(2)} s_n^{(3)}, \tag{9.132}$$

where $s_n^{(i)}$ is defined in (9.120) and obeys $s_n^{(i)} \leq 2^i(1 + 2^{i+1} n^{-2(c_\star - 1)})$ whenever $c_\star > 2$. Eq. (9.124) now immediately follows. Invoking (6.6) of Proposition 6.3 we get that on $\Omega^\star$, for all but a finite number of indices n, if $c_\star > 2$,

$$(a_n/|\mathcal{V}_n^\circ|) \sum_{m=1}^{\ell_n^\circ - 1} \mathbb{E}\big[I_{2,m}^{(3)}\big] \leq 2^5(1 + o(1)) \frac{\ell_n^\circ}{n a_n} [W_{n,3}(u)]^2. \tag{9.133}$$

The lemma now follows by a first order Tchebychev inequality. □

9.3.4. *Proof of Proposition 9.5.* The proof of Proposition 9.5 is now a mere formality. Recall that $c_\star > 2$ and that a_n obeys (9.51) for some $0 < \varepsilon < 1$. Choose $L_1 = n^{-1-(c_\star - 2)/2}$ and $L_2 = 4 \log n$ in Lemma 9.9. Then $n^2 a_n L_2 / 2^n = o(1)$, $\lim_{n\to\infty} \phi_n(u, L_1, L_2) \to 0$, and $\sum_n (2ne^{-L_2} + 2L_1) < \infty$, so that by Lemma 9.8, Lemma 9.9, and Borel-Cantelli Lemma,

$$\lim_{n\to\infty} \bar{\nu}_n^\circ(u,\infty) = \nu^\dagger(u,\infty), \quad \mathbb{P} - \text{almost surely}, \tag{9.134}$$

for all $u > 0$. Because $\bar{\nu}_n^\circ(u,\infty)$ is a sequence of monotonic functions of $u > 0$ whose limit, $\nu^\dagger(u,\infty)$, is continuous, (9.134) entails the existence of a subset $\overline{\Omega}_1^{\mathsf{LLN}} \subset \Omega$ with the property that $\mathbb{P}(\overline{\Omega}_1^{\mathsf{LLN}}) = 1$, and such that on $\overline{\Omega}_1^{\mathsf{LLN}}$,

$$\lim_{n\to\infty} \bar{\nu}_n^\circ(u,\infty) = \nu^\dagger(u,\infty), \quad \forall u > 0. \tag{9.135}$$

We prove in the same way, using the monotonicity of $\bar{\sigma}_n^\circ(u,\infty)$, Lemma 9.10, and Lemma 9.11 (with L_1 and L_2 as above and $L_3 = n^{-2}$, so that $\lim_{n\to\infty} n\psi_n(u, L_1, L_2, L_3) = 0$ and $\sum_n (2ne^{-L_2} + 2L_1 + 2L_2) < \infty$) that there exists a subset $\overline{\Omega}_2^{\mathsf{LLN}} \subset \Omega$ of full measure such that, on $\overline{\Omega}_2^{\mathsf{LLN}}$,

$$\lim_{n\to\infty} n\bar{\sigma}_n^\circ(u,\infty) = 2\nu^\dagger(2u,\infty), \quad \forall u > 0. \tag{9.136}$$

Taking $\overline{\Omega}^{\mathsf{LLN}} = \overline{\Omega}_1^{\mathsf{LLN}} \cap \overline{\Omega}_2^{\mathsf{LLN}}$ completes the proof of Proposition 9.5 .

9.4. **Conclusion of the proof of Theorem 3.4.** It suffices to prove that under the assumptions of Theorem 3.4, Conditions (A1), (A2), and (A3) of Theorem 9.1 are verified $\mathbb{P}$-almost surely when $\nu^\dagger$ in Condition (A1) is as in (1.23).

9.4.1. *Verification of Conditions (A1) and (A2).* It immediately follows Lemma 9.2, Theorem 9.3, and Proposition 9.5 that under the assumptions therein, $\mathbb{P}$-almost surely, for all $u > 0$ and all $t > 0$,

$$\lim_{n\to\infty} \bar{\nu}_n^{J_n^\circ,t}(u,\infty) = t\nu^\dagger(u,\infty) \text{ in } P^\circ\text{-probability}, \tag{9.137}$$

$$\lim_{n\to\infty} \bar{\sigma}_n^{J_n^\circ,t}(u,\infty) = 0 \text{ in } P^\circ\text{-probability}. \tag{9.138}$$

Conditions (A1) and (A2) are thus satisfied $\mathbb{P}$-almost surely.

9.4.2. *Verification of Condition (A3).* This still requires a little work. Given $\epsilon > 0$, define

$$\eta_{n,k}(\epsilon) \equiv \frac{a_n}{2^n} \sum_{1\le l\le L^\star} \mathbb{1}_{\{|C^\star_{n,l}|=k\}} \sum_{x\in C^\star_{n,l}} A_{n,l}(x), \quad k \ge 2, \tag{9.139}$$

where, given $x \in C^\star_{n,l}$, $1 \le l \le L^\star$,

$$A_{n,l}(x) \equiv E_x\big(\mathbb{1}_{\{b_n^{-1}T^\star_{n,l}\le\epsilon\}} b_n^{-1} T^\star_{n,l}\big). \tag{9.140}$$

One readily sees that Condition (A3) will be verified $\mathbb{P}$-almost surely if

$$\lim_{\epsilon\downarrow 0} \limsup_{n\uparrow\infty} \sum_{k\ge 2} \eta_{n,k}(\epsilon) = 0, \quad \mathbb{P}\text{-a.s.}. \tag{9.141}$$

Note that $\eta_{n,k}(\epsilon)$ is of the form (9.69) with $Q^u_{n,l}(x)$ replaced by $A_{n,l}(x)$ and hence, as in (9.71), may be written as

$$\eta_{n,k}(\epsilon) \equiv \frac{a_n}{2^n} \sum_{\mathcal{C}\in\mathcal{G}_k} \prod_{x\in\mathcal{C}} \chi_n(x) \prod_{x'\in\partial\mathcal{C}} \overline{\chi}_n(x') \sum_{x\in\mathcal{C}} A_{n,\mathcal{C}}(x), \tag{9.142}$$

where $A_{n,\mathcal{C}}(x)$ stands for $A_{n,l}(x)$ with $C^\star_{n,l} \equiv \mathcal{C}$. As in the proof of Lemma 9.8 we note that on $\Omega^\star$, $k^\star_n \le \frac{n}{(c_\star-2)\log n}$ for all large enough n, and treat the terms $k = 2$ and $3 \le k \le k^\star_n$ separately. Throughout the proof we set $\bar{a}_n = \sqrt{na_n}$, $\bar{b}_n = b_n(n-1)$, and define

$$\gamma_n(\mathcal{C}') = \min_{x\in\mathcal{C}} w_n(x)/\bar{b}_n, \quad \mathcal{C}' \in \mathcal{G}_2. \tag{9.143}$$

• ***The term*** $k = 2$**.** Let us establish that for all large enough n and small enough ϵ, the mean and variance of $\eta_{n,2}(\epsilon)$ obey

$$\mathbb{E}\eta_{n,2}(\epsilon) \le 2^{-n/6} + 2\epsilon^{1-\left(2\alpha_c(\frac{\varepsilon}{2})+o(1)\right)}, \tag{9.144}$$

$$\mathbb{E}\left(\eta_{n,2}(\epsilon) - \mathbb{E}\eta_{n,2}(\epsilon)\right)^2 \le 2\frac{a_n}{2^n}\left[2^{-n/6} + 4\epsilon^{2-\left(2\alpha_c(\frac{\varepsilon}{2})+o(1)\right)}\right]. \tag{9.145}$$

We first prove (9.144). By Proposition 5.1, (i), and integration by parts, for all $x \in \mathcal{C}$,

$$A_{n,\mathcal{C}}(x) \le \tfrac{1}{b_n}\sum_{i=0}^{\lfloor b_n\epsilon\rfloor}\left(1 - \tfrac{1}{1+\min_{x\in\mathcal{C}} w_n(x)/(n-1)}\right)^i \tag{9.146}$$

$$\le [1+o_{n,1}(1)]\varphi_\epsilon\left([1+o_{n,2}(1)]\min_{x\in\mathcal{C}} w_n(x)/\bar{b}_n\right) \tag{9.147}$$

where $|o_{n,i}(1)| \le \mathcal{O}(r_n^{-1}(\rho^\star_n))$, $i = 1, 2$, and

$$\varphi_\epsilon(y) = y(1 - e^{-\epsilon/y}), \quad y \ge 0. \tag{9.148}$$

Plugging (9.147) in (9.142) yields

$$\mathbb{E}\eta_{n,2}(\epsilon) \le [1+o_{n,1}(1)]\mathbb{E}\left[\bar{a}_n^2\varphi_\epsilon\left([1+o_{n,2}(1)]\gamma_n(\mathcal{C})\right)\mathbb{1}_{\{\gamma_n(\mathcal{C})\ge r_n(\rho^\star_n)/\bar{b}_n\}}\right]. \tag{9.149}$$

Now for $\epsilon > r_n(\rho^\star_n)/\bar{b}_n$ split $\mathbb{1}_{\{\gamma_n(\mathcal{C})\ge r_n(\rho^\star_n)/\bar{b}_n\}}$ into $\mathbb{1}_{\{\gamma_n(\mathcal{C})\ge\epsilon\}} + \mathbb{1}_{\{r_n(\rho^\star_n)/\bar{b}_n\le\gamma_n(\mathcal{C})<\epsilon\}}$. On the one hand, observing that $\varphi_\epsilon(y) \le \epsilon$ for all $y > 0$, we have

$$\mathbb{E}\left[\bar{a}_n^2\varphi_\epsilon\left([1+o_{n,2}(1)]\gamma_n(\mathcal{C})\right)\mathbb{1}_{\{\gamma_n(\mathcal{C})\ge\epsilon\}}\right] \le \epsilon\bar{a}_n^2\mathbb{P}\left(\gamma_n(\mathcal{C}) \ge \epsilon\right) = \epsilon^{1-2\left(\alpha_c(\frac{\varepsilon}{2})+o(1)\right)}, \tag{9.150}$$

where the last equality follows from Lemma 9.7, (i). On the other hand $\varphi_\epsilon(y) \le y$ for all $y > 0$. From this, integration by part, and (9.56) we obtain,

$$\mathbb{E}\left[\bar{a}_n^2 \varphi_\epsilon\left([1+o_{n,2}(1)]\gamma_n(\mathcal{C})\right) \mathbb{1}_{\{r_n(\rho_n^\star)/\bar{b}_n \le \gamma_n(\mathcal{C}) < \epsilon\}}\right] \le [1+o_{n,2}(1)] \int_{r_n(\rho_n^\star)/\bar{b}_n}^{\epsilon} F_n^2(y)dy. \tag{9.151}$$

Given $0 < \delta < 1$, split the domain of integration in (9.151) into $[r_n(\rho_n^\star)/\bar{b}_n, \bar{b}_n^{-\delta}] \cup [\bar{b}_n^{-\delta}, \epsilon]$. Using that $F_n^2(y) \le \bar{a}_n^2$ on the first domain, and using Lemma 9.7, (ii), on the the second,

$$\int_{r_n(\rho_n^\star)/\bar{b}_n}^{\epsilon} F_n^2(y)dy \le \bar{a}_n^2 \bar{b}_n^{-\delta} + \tfrac{1+o(1)}{1-2(1-\frac{\delta}{2})\alpha_n}\left(\tfrac{1}{1-\delta}\right)^2 \epsilon^{1-2(1-\frac{\delta}{2})\alpha_n}, \tag{9.152}$$

where $0 \le \alpha_n = \alpha_c(\frac{\varepsilon}{2}) + o(1)$. By definition of $\bar{a}_n$, $\bar{b}_n$, (2.19), and the assumption that $\beta > 2\beta_c(\varepsilon/2)$, we get $\bar{a}_n^2 \bar{b}_n^{-\delta} \le \exp\{n\,[\beta_c^2(\varepsilon/2)(1-2\delta(1+o(1)))]\}$. Hence, choosing $\delta = 2/3$, $\int_{r_n(\rho_n^\star)/\bar{b}_n}^{\epsilon} F_n^2(y)dy \le 2^{-n/6} + \frac{9}{1-[2\alpha_c(\varepsilon/2)+o(1)](2/3)}\epsilon^{1-\frac{2}{3}[2\alpha_c(\varepsilon/2)+o(1)]}$. Collecting our bounds we arrive at (9.144).

Turning to the variance we have

$$\mathbb{E}\left(\eta_{n,2}(\epsilon) - \mathbb{E}\eta_{n,2}(\epsilon)\right)^2 = \left(\frac{a_n}{2^n}\right)^2 \mathbb{E}\left(\textstyle\sum_{\mathcal{C}\in\mathcal{G}_2}[Y_n(\mathcal{C}) - \mathbb{E}Y_n(\mathcal{C})]\right)^2, \tag{9.153}$$

where $Y_n(\mathcal{C}) \equiv \prod_{x\in\mathcal{C}} \chi_n(x) \prod_{x'\in\partial\mathcal{C}} \overline{\chi}_n(x') \sum_{x\in\mathcal{C}} A_{n,\mathcal{C}}(x)$, $\mathcal{C} \in \mathcal{G}_2$. Observing that $Y_n(\mathcal{C})$ and $Y_n(\mathcal{C}')$ are independent whenever $\mathcal{C} \neq \mathcal{C}'$ and $\partial\mathcal{C} \cap \partial\mathcal{C}' = \emptyset$, and that $Y_n(\mathcal{C})Y_n(\mathcal{C}') = 0$ whenever $\mathcal{C} \neq \mathcal{C}'$ and $\mathcal{C} \cap \mathcal{C}' \neq \emptyset$, we readily get that $\mathbb{E}\left(\eta_{n,2}(\epsilon) - \mathbb{E}\eta_{n,2}(\epsilon)\right)^2 \le I_n^{=} + I_n^{\neq}$,

$$I_n^{=} \equiv \left(\frac{a_n}{2^n}\right)^2 \textstyle\sum_{\mathcal{C}\in\mathcal{G}_2} \mathbb{E}[Y_n^2(\mathcal{C})], \tag{9.154}$$

$$I_n^{\neq} \equiv \left(\frac{a_n}{2^n}\right)^2 \textstyle\sum_{\mathcal{C},\mathcal{C}'}^{(1)} (\mathbb{E}[Y_n(\mathcal{C})Y_n(\mathcal{C}')] - [\mathbb{E}Y_n(\mathcal{C})][\mathbb{E}Y_n(\mathcal{C}')]), \tag{9.155}$$

where, as in (9.117), the sum $\Sigma^{(1)}$ is over all $\mathcal{C} \in \mathcal{G}_2$ and $\mathcal{C}' \in \mathcal{G}_{2'}$ such that $\mathrm{dist}(\mathcal{C}, \mathcal{C}') = 2$. We bound (9.154) in just the same way as $\mathbb{E}\eta_{n,2}(\epsilon)$, namely, using (9.147) in (9.154) gives

$$I_n^{=} \le 2\frac{a_n}{2^n}[1+o_{n,1}(1)]\mathbb{E}\left[\bar{a}_n^2 \varphi_\epsilon^2\left([1+o_{n,2}(1)]\gamma_n(\mathcal{C})\right) \mathbb{1}_{\{\gamma_n(\mathcal{C}) \ge r_n(\rho_n^\star)/\bar{b}_n\}}\right], \tag{9.156}$$

and proceeding as in (9.149)-(9.152) to evaluate (9.149), we obtain (9.145). To Bound $I_n^{\neq}$ note that

$$I_n^{\neq} = \left(\frac{a_n}{2^n}\right)^2 \textstyle\sum_{\mathcal{C},\mathcal{C}'}^{(1)} \mathbb{E}[Z_n(\mathcal{C})]\mathbb{E}[Z_n(\mathcal{C}')]\Delta_n(\mathcal{C}, \mathcal{C}') \tag{9.157}$$

where $Z_n(\mathcal{C}) \equiv \prod_{x\in\mathcal{C}} \chi_n(x) \sum_{x\in\mathcal{C}} A_{n,\mathcal{C}}(x)$, and

$$\begin{aligned} \Delta_n(\mathcal{C}, \mathcal{C}') &\equiv \mathbb{E}[\textstyle\prod_{y\in\partial\mathcal{C}\cup\partial\mathcal{C}'} \overline{\chi}_n(y)] - \mathbb{E}[\textstyle\prod_{x\in\partial\mathcal{C}} \overline{\chi}_n(x)]\mathbb{E}[\textstyle\prod_{x'\in\partial\mathcal{C}'} \overline{\chi}_n(x')] \\ &= (1-n^{-c_\star})^3(1-(1-n^{-c_\star})). \end{aligned} \tag{9.158}$$

Observing that the right hand side of (9.149) (and a fortiori the r.h.s. of (9.144)) is an upper bound on $\bar{a}_n^2\mathbb{E}[Z_n(\mathcal{C})]$, and that the sum $\Sigma^{(1)}$ contains at most $4!n^4 2^{n-1}$ terms, we obtain

$$I_n^{\neq} \le 4!(1-n^{-c_\star})^3 n^{2-c_\star} 2^{-n}\left(2^{-n/6} + 2\epsilon^{1-\left(2\alpha_c(\frac{\varepsilon}{2})+o(1)\right)}\right)^2. \tag{9.159}$$

Combining (9.156) and (9.159) now yields (9.145).

Since $n^2\sum_n a_n/2^n < \infty$ it follows from (9.144), Borel-Cantelli Lemma (through a second order Tchebychev inequality), and (9.145) that $\lim_{n\uparrow\infty} \eta_{n,2}(\epsilon) = 2\epsilon^{1-2\alpha_c(\varepsilon/2)}$ $\mathbb{P}$-almost surely, for all $\varepsilon > 0$. Observing that $\eta_{n,2}(\epsilon)$ is a monotonic function of ϵ, and arguing as in the proof of Proposition 9.5 (see (9.134)-(9.135)), we obtain that

$$\lim_{\epsilon\downarrow 0} \limsup_{n\uparrow\infty} \eta_{n,2}(\epsilon) = 0, \quad \mathbb{P}\text{- almost surely.} \tag{9.160}$$

• ***The terms*** $3 \le k \le k_n^\star$**.** Note that $\eta_n(\epsilon) - \eta_{n,2}(\epsilon) > 0$. Our strategy is to bound $\mathbb{E}(\eta_n(\epsilon) - \eta_{n,2}(\epsilon))$ from above and use a first order Tchebychev inequality to infer from it $\mathbb{P}$-a.s. convergence of $\eta_n(\epsilon) - \eta_{n,2}(\epsilon)$ to zero.

As in the proof of Lemma 9.8 we denote by $\varrho_{n,\mathcal{C}}(0)$, $\varrho_{n,\mathcal{C}}(1)$, and $\theta^\star_{n,\mathcal{C}}$ the quantities $\varrho_{n,l}(0)$, $\varrho_{n,l}(1)$, and $\theta^\star_{n,l}$ from (5.2)-(5.4) with $C^\star_{n,l} \equiv \mathcal{C}$. Similarly $T^\star_{n,\mathcal{C}}$ stands for $T^\star_{n,l}$ with $C^\star_{n,l} \equiv \mathcal{C}$. Decomposing the event $\{b_n^{-1} T^\star_{n,\mathcal{C}} \le \epsilon\}$ into $\mathcal{A}^{(1)} \cup \mathcal{A}^{(2)} \cup \mathcal{A}^{(2)}$ where $\mathcal{A}^{(1)} \equiv \{T^\star_{n,\mathcal{C}} \le b_n\epsilon < \theta^\star_{n,\mathcal{C}}\}$, $\mathcal{A}^{(2)} \equiv \{T^\star_{n,\mathcal{C}} < \theta^\star_{n,\mathcal{C}} \le b_n\epsilon\}$, and $\mathcal{A}^{(3)} \equiv \{\theta^\star_{n,\mathcal{C}} < T^\star_{n,\mathcal{C}} \le b_n\epsilon\}$, define

$$A^{(i)}_{n,\mathcal{C}}(x) \equiv E_x\big(b_n^{-1} T^\star_{n,\mathcal{C}} \mathbb{1}_{\mathcal{A}^{(i)}}\big), \tag{9.161}$$

and denote by $\eta^{(i)}_{n,k}(\epsilon)$ the quantity $\eta_{n,k}(\epsilon)$ where $A^{(i)}_{n,\mathcal{C}}(x)$ is substituted for $A_{n,\mathcal{C}}(x)$. Thus

$$0 < \eta_n(\epsilon) - \eta_{n,2}(\epsilon) = \textstyle\sum_{i=1}^3 \sum_{k=3}^{k_n^\star} \eta^{(i)}_{n,k}(\epsilon), \quad \eta_{n,k}(\epsilon) = \sum_{i=1}^3 \eta^{(i)}_{n,k}(\epsilon). \tag{9.162}$$

To bound $\eta_n(\epsilon) - \eta_{n,2}(\epsilon)$ it now suffices to bound each $\mathbb{E}\eta^{(i)}_{n,k}(\epsilon)$, $1 \le i \le 3$, $3 \le k \le k_n^\star$.

Using the trivial bound $A^{(1)}_{n,l}(x) \le \epsilon 1_{\theta^\star_{n,\mathcal{C}} \ge b_n\epsilon}$, the term $\mathbb{E}[\eta^{(1)}_{n,k}(\epsilon)]$ is bounded in exactly the same way as $\bar\nu_n^{\circ,(k),-}(u,\infty)$ in (9.89)-(9.91), namely

$$\mathbb{E}[\eta^{(1)}_{n,k}(\epsilon)] \le \epsilon k(k-2)! n^{-(c_\star-1)(k-2)} \left(2\beta n^2 k^5/\epsilon r_n(\rho_n^\star)\right)^{2\alpha(\varepsilon/2)(1+o(1))}. \tag{9.163}$$

From the bound $A^{(2)}_{n,\mathcal{C}}(x) \le b_n^{-1}\theta^\star_{n,\mathcal{C}} \mathbb{1}\{b_n^{-1}\theta^\star_{n,\mathcal{C}} \le \epsilon\}$, (5.2)-(5.4), and Lemma 4.5, we get

$$\begin{aligned} A^{(2)}_{n,\mathcal{C}}(x) &\le \varrho_{n,\mathcal{C}}(0) \tfrac{2\beta n^2 k^5}{\bar b_n r_n(\rho_n^\star)} \mathbb{1}_{\{\varrho_{n,\mathcal{C}}(0) \le \epsilon \bar b_n r_n(\rho_n^\star)/2\beta n^2 k^5\}} && (9.164)\\ &\le \textstyle\sum_{\{x,y\} \in G(\mathcal{C})} \gamma_n(\{x,y\}) \tfrac{2\beta n^2 k^5}{r_n(\rho_n^\star)} \mathbb{1}_{\{\gamma_n(\{x,y\}) \le \epsilon r_n(\rho_n^\star)/2\beta n^2 k^5\}}. && (9.165)\end{aligned}$$

Inserting this in (9.142) and, again, proceeding as in (9.89)-(9.91),

$$\mathbb{E}[\eta^{(2)}_{n,k}(\epsilon)] \le \frac{\beta n k^5 (k-2)!}{n^{(c_\star-1)(k-2)}} \mathbb{E}\left[\bar a_n^2(\gamma_n(\mathcal{C}')/r_n(\rho_n^\star)) \mathbb{1}_{\{1/\bar b_n \le \gamma_n(\mathcal{C}')/r_n(\rho_n^\star) \le \epsilon/\beta n^2\}}\right]. \tag{9.166}$$

An expectation similar to that appearing in (9.166) was estimated in (9.151). Observing that $r_n^{-1}(\rho_n^\star) \int_{r_n(\rho_n^\star)/\bar b_n}^{\epsilon r_n(\rho_n^\star)/\beta n^2} F_n^2(y)dy = \int_{1/\bar b_n}^{\epsilon/\beta n^2} [\tilde a_n \mathbb{P}(w_n(x) \ge v\tilde b_n)]^2 dy$ where $\tilde b_n = \bar b_n r_n(\rho_n^\star)$ and $\tilde a_n \mathbb{P}(w_n(x) \ge \tilde b_n) = 1$, and proceeding as in (9.151)- (9.144), we obtain

$$\mathbb{E}[\eta^{(2)}_{n,k}(\epsilon)] \le \frac{2\beta n k^6 (k-2)!}{n^{(c_\star-1)(k-2)}} \left[2^{-n/6} r_n(\rho_n^\star)^{-2/3} + 2(\epsilon/\beta n^2)^{1-\left(2\alpha_c(\frac{\varepsilon}{2})+o(1)\right)}\right]. \tag{9.167}$$

It remains to bound $\mathbb{E}[\eta^{(3)}_{n,k}(\epsilon)]$. Using Proposition 5.1, (ii) and proceeding as in (9.146)-(9.147), we get that on $\Omega_0 \cap \Omega^\star$, for all but a finite number of indices n, for all $x \in \mathcal{C}$,

$$\begin{aligned} A^{(3)}_{n,\mathcal{C}}(x) &\le [1+o(1)]\varphi_\epsilon\left([1+o(1)]k\varrho_{n,\mathcal{C}}(0)/\bar b_n\right) && (9.168)\\ &\le [1+o(1)]\left(\epsilon 1_{\{k\varrho_{n,\mathcal{C}}(0) < \bar b_n\epsilon\}} + k\varrho_{n,\mathcal{C}}(0) 1_{\{k\varrho_{n,\mathcal{C}}(0) \ge \bar b_n\epsilon\}}\right) && (9.169)\end{aligned}$$

where the last line follows from the bounds $\varphi_\epsilon(y) \le \epsilon$ and $\varphi_\epsilon(y) \le y$, both valid for all $y > 0$. Comparing the right hand side of (9.169) to the bounds

$$A^{(1)}_{n,l}(x) \le \epsilon \mathbb{1}_{\{\theta^\star_{n,\mathcal{C}} \ge b_n\epsilon\}}, \quad A^{(2)}_{n,\mathcal{C}}(x) \le b_n^{-1}\theta^\star_{n,\mathcal{C}} \mathbb{1}_{\{b_n^{-1}\theta^\star_{n,\mathcal{C}} \le \epsilon\}} \tag{9.170}$$

we just used, one sees that the contributions to $\mathbb{E}[\eta^{(3)}_{n,k}(\epsilon)]$ coming from the two summands in (9.169) can be bounded in the same way as $\mathbb{E}[\eta^{(1)}_{n,k}(\epsilon)]$ and $\mathbb{E}[\eta^{(2)}_{n,k}(\epsilon)]$, respectively, replacing $\theta^\star_{n,\mathcal{C}}$ by $k\varrho_{n,\mathcal{C}}(0)$ in (9.170). Doing this, it follows from (9.163) and (9.167) that

$$\mathbb{E}[\eta^{(3)}_{n,k}(\epsilon)] \le \frac{k(k-2)!}{\eta^{(c_\star-1)(k-2)}} \left[5\epsilon(nk/\epsilon)^{\left(2\alpha_c(\frac{\varepsilon}{2})+o(1)\right)} + k2^{-n/6} + k2\epsilon^{1-\left(2\alpha_c(\frac{\varepsilon}{2})+o(1)\right)}\right]. \quad (9.171)$$

Finally, collecting (9.163), (9.167), and (9.171), and using (7.24) to perform the sum over k, we arrive at

$$\mathbb{E}[\textstyle\sum_{i=1}^3 \sum_{k=3}^{k^\star_n} \eta^{(i)}_{n,k}(\epsilon)] \le \frac{c_0}{n^{(c_\star-2)}}(\epsilon/n)^{1-\left(2\alpha_c(\frac{\varepsilon}{2})+o(1)\right)} + 2^{-n/6} \quad (9.172)$$

for some constant $0 < c_0 \equiv c_0(\beta,\varepsilon) < \infty$. Since by assumption $c_\star > 3$ and $1 - \left(2\alpha_c(\frac{\varepsilon}{2}) + o(1)\right) > 0$ for all n large enough, the r.h.s. of (9.172) is summabe in n. Combining this and (9.162) yields that $0 < \eta_n(\epsilon) - \eta_{n,2}(\epsilon) \to 0$ $\mathbb{P}$-a.s., for all $\varepsilon > 0$. Since each $\eta_{n,k}(\epsilon)$ in (9.142) is a monotonic function of ϵ, so is $\eta_n(\epsilon) - \eta_{n,2}(\epsilon)$, and thus, arguing again as in the proof of Proposition 9.5,

$$\lim_{\epsilon\downarrow 0} \limsup_{n\uparrow\infty} (\eta_n(\epsilon) - \eta_{n,2}(\epsilon)) = 0, \quad \mathbb{P}\text{-a.s.}. \quad (9.173)$$

Since (9.160) and (9.173) imply (9.141), Condition (A3) is verified $\mathbb{P}$-almost surely under the assumptions of Theorem 9.1.

Having established that all three conditions of Theorem 9.1 are satisfied $\mathbb{P}$-almost surely with $\nu^\dagger$ given by (1.23), the proof of Theorem 3.4 is done.

10. Convergence of the back end clock process below the critical line: proof of Theorem 3.5

Since BECP below the critical line has a deterministic limit, this case is simpler than that of Theorem 3.4. The proof proceeds in two steps. By (3.11) and (3.16),

$$S^\dagger_n(t) = b_n^{-1} \sum_{i=0}^{k^\dagger_n(t)-1} \Lambda^\dagger_n(J^\dagger_n(i)), \quad (10.1)$$

where by (3.12), $\Lambda^\dagger_n(J^\dagger_n(i))$ is non zero if and only if $J^\dagger_n(i) \in \cup_{1\le l\le L^\star} C^\star_{n,l}$. The first step consists in counting the number of distinct sets $C^\star_{n,l}$ that $J^\dagger_n$ visits along suitably chosen subsequences $0 \le i \le k^\dagger_n(t) - 1$, as well as the typical and maximal number of visits to each set. This information is then used to extract sums of independent random variables from the sum (10.1), and a remainder. In a second step, and after suitable truncation, each of these sums is controlled by a classical mean-variance calculation, and the remainder is shown to be sub-leading.

10.1. **Step one.** For simplicity we assume throughout that $\lfloor k^\circ_n(t)/\ell^\circ_n \rfloor = k^\circ_n(t)/\ell^\circ_n$. We first decompose the index set $I = \{0,\dots,k^\dagger_n(t)-1 : J^\dagger_n(i) \in \mathcal{V}^\circ_n\}$ into ℓ°_n disjoint subsets, $I = \cup_{0\le j\le \ell^\circ_n-1} I_j$, defined through

$$I_j = \left\{ 0 \le i \le k^\dagger_n(t)-1 \,\middle|\, \exists 0 \le i'' \le \frac{k^\circ_n(t)}{\ell^\circ_n} - 1 \, s.t. \sum_{i'=0}^{i} \mathbb{1}_{\{J^\dagger_n(i')\in\mathcal{V}^\circ_n\}} = j + i''\ell^\circ_n \right\}. \quad (10.2)$$

The reason behind this choice is that, by definition of $J^\dagger_n$ and J°_n, (3.15), and (3.19),

$$\{J^\dagger_n(i), i \in I_j\} \stackrel{d}{=} \{J^\circ_n(j + i\ell^\circ_n), i = 0,\dots,\lfloor k^\circ_n(t)/\ell^\circ_n\rfloor\}, \quad 0 \le j \le \ell^\circ_n - 1, \quad (10.3)$$

(where equality holds in distribution) and by the mixing property of Proposition 6.1, the chain J_n° observed along such subsequences is well approximated by an i.i.d. sequence. Now clearly, visits of $J_n^\dagger$ to $\cup_{1\le l\le L^\star} C^\star_{n,l}$ can only occur along subsequences of the form $\{i+1 : i \in I_j\}$. For $0 \le j \le \ell_n^\circ - 1$, let $N_{n,j}$, respectively $M_{n,j}$, be the number of distinct sets $C^\star_{n,l}$, respectively the total number of sets $C^\star_{n,l}$ that the chain $J_n^\dagger$ visits along each subsequence $\{i+1 : i \in I_j\}$. Define

$$\mathcal{L}'_n \equiv \{1,\dots,L^\star\} \setminus \big(\cup_{2\le k\le k_n^\star} \mathcal{L}_n(k)\big) \equiv \cup_{2\le k\le k_n^\star} \mathcal{L}'_n(k) \tag{10.4}$$

where, for $2 \le k \le k_n^\star \equiv \max_{1\le l\le L^\star} |C^\star_{n,l}|$,

$$\mathcal{L}_n(k) \equiv \{1 \le l \le L^\star : |C^\star_{n,l}| = k, \partial C^\star_{n,l} \cap \partial C^\star_{n,l'} = \emptyset \,\forall l' \ne l, 1 \le l' \le L^\star\}, \tag{10.5}$$

$$\mathcal{L}'_n(k) \equiv \{l \in \mathcal{L}'_n : |C^\star_{n,l}| = k\}. \tag{10.6}$$

With these definitions $N_{n,j} = \sum_{k=2}^{k_n^\star} N_{n,j}(k) + N'_{n,j}$ and $M_{n,j} = \sum_{k=2}^{k_n^\star}(M_{n,j}(k) + M'_{n,j})$ where

$$N_{n,j}(k) = \sum_{l\in\mathcal{L}_n(k)} \mathbb{1}_{\{\exists i\in I_j : J_n^\dagger(i+1)\in C^\star_{n,l}\}}, \quad N'_{n,j} = \sum_{l\in\mathcal{L}'_n} \mathbb{1}_{\{\exists i\in I_j : J_n^\dagger(i+1)\in C^\star_{n,l}\}}, \tag{10.7}$$

$$M_{n,j}(k) = \sum_{l\in\mathcal{L}_n(k)} \sum_{i\in I_j} \mathbb{1}_{\{J_n^\dagger(i+1)\in C^\star_{n,l}\}}, \quad M'_{n,j}(k) = \sum_{l\in\mathcal{L}'_n(k)} \sum_{i\in I_j} \mathbb{1}_{\{J_n^\dagger(i+1)\in C^\star_{n,l}\}}. \tag{10.8}$$

Note that the number of sets $C^\star_{n,l}$, $l \in \mathcal{L}_n(k)$, that are visited at least twice along a given subsequence $\{i+1 : i \in I_j\}$ is at most $M_{n,j}(k) - N_{n,j}(k)$. Lastly, let $m_{n,j}$ be the maximum number of visits to a given $C^\star_{n,l}$:

$$m_{n,j} = \max_{1\le l\le L^\star} \sum_{i\in I_j} \mathbb{1}_{\{J_n^\dagger(i+1)\in C^\star_{n,l}\}}. \tag{10.9}$$

Proposition 10.1 below yields control over these various quantities. In order to state it we need the following definitions. Given constants $0 < c_k, c'_k < \infty$ and $c_2 = 1$, let the functions $\delta_n(2)$ and $\eta_n(k)$, $2 \le k \le k_n^\star$, be defined through.

$$\eta_n(k) = \frac{c_k k! \lfloor a_n t\rfloor}{\ell_n^\circ 2n^{k(c_\star-1)+1}}(1+o(1)), \quad \delta_n(2) = \frac{\lfloor a_n t\rfloor}{\ell_n^\circ 2^n}\eta_n(2), \tag{10.10}$$

We know (see (7.7)) that on $\Omega^\star$, for large enough n, $k_n^\star \le n/((c_\star-2)\log n)$. Given an integer $2 < K < \infty$ define, for $2 \le k \le K$,

$$\mathcal{A}^{1,-}_{n,j,k} = \big\{|N_{n,j}(k) - \eta_n(k)| < n^{2c_\star}\sqrt{\eta_n(k)}\big\}, \tag{10.11}$$

$$\mathcal{A}^{2,-}_{n,j,k} = \{M_{n,j}(k) - N_{n,j}(k) < (k/n)\eta_n(k)\}, \tag{10.12}$$

$$\mathcal{A}^{3,-}_{n,j,k} = \big\{M'_{n,j}(k) < c'_k n k_n^\star \eta_n(k+2)\big\}, \tag{10.13}$$

and for $K < k \le k_n^\star$,

$$\mathcal{A}^{1,+}_{n,j,k} = \big\{N_{n,j}(k) - n^2\eta_n(k) < n^{2c_\star+1}\sqrt{\eta_n(k)}\big\}, \tag{10.14}$$

$$\mathcal{A}^{2,+}_{n,j,k} = \{M_{n,j}(k) - N_{n,j}(k) < kn\eta_n(k)\}, \tag{10.15}$$

$$\mathcal{A}^{3,+}_{n,j,k} = \big\{M'_{n,j}(k) < c'_k n^3 k_n^\star \eta_n(k+2)\big\}, \tag{10.16}$$

Finally set $\mathcal{A}^0_{n,j} = \big\{|[M_{n,j}(2) - N_{n,j}(2)] - \delta_n(2)| < n^{2c_\star}\sqrt{\delta_n(2)}\big\}$.

Proposition 10.1. *Assume that $c_\star > 3$ and let $2 < K < \infty$ be a fixed integer. For all $\beta > 0$, there exists a subset $\Omega^{BCL} \subset \Omega$ with $\mathbb{P}(\Omega^{BCL}) = 1$ such that on $\Omega_1 \cap \Omega^\star \cap \Omega^{BCL}$ the following holds: for all $t \in [0,T]$, for all but a finite number of indices n,*

$$P^\dagger_{\pi^\circ_n}\left(\bigcap_{0\le j\le \ell^\circ_n-1}\mathcal{A}^0_{n,j}\bigcap_{i=1}^3\Big((\bigcap_{2\le k\le K}\mathcal{A}^{i,-}_{n,j,k})\cap(\bigcap_{K<k\le k^\star_n}\mathcal{A}^{i,+}_{n,j,k})\Big)\right)\ge 1-n^{-2(c_\star-1)+\bar c} \tag{10.17}$$

where $\bar c > 0$ is arbitrary and for some constants $0 < c_k, c'_k < \infty$ and $c_2 = 1$. Moreover,

$$P^\dagger_{\pi^\circ_n}\left(\max_{0\le j\le \ell^\circ_n-1} m_{n,j}\ge \tfrac{3}{1-\varepsilon}\right)\le 2^{-n}. \tag{10.18}$$

Proof of Proposition 10.1. The proof of Proposition 10.1 hinges on Lemma 10.2 below, that allows to substitute simpler variables for the variables $N_{n,j}(k)$ and $M_{n,j}(k)$. In order to define them let $\{\widetilde J^\circ_n(i), i\ge 0, n>0\}$ and $\big\{X_{n,i}(x), x\in(\cup_{1\le l\le L^\star}\partial C^\star_{n,l}), i\ge 1\big\}$ be two independent arrays of i.i.d. r.v.'s, defined on a common probability space $(\widetilde\Omega,\widetilde{\mathcal{F}},\widetilde P)$, such that $\widetilde P(\widetilde J^\circ_n(i)=x)=\pi^\circ_n(x)$, $x\in\mathcal{V}^\circ_n$, and, for each $x\in(\cup_{1\le l\le L^\star}\partial C^\star_{n,l})$,

$$\widetilde P(X_{n,i}(x)=1)=1-\widetilde P(X_{n,i}(x)=0)=n^{-1}|\partial x\cap(\cup_{1\le l\le L^\star}C^\star_{n,l})|. \tag{10.19}$$

Thus the $X_{n,i}$'s are Bernoulli r.v.'s. Using these arrays, define the quantities

$$\widetilde N_{n,j}(k)=\textstyle\sum_{l\in\mathcal{L}_n(k)}\mathbb{1}_{\{\exists 0\le i\le\lfloor k^\circ_n(t)/\ell^\circ_n\rfloor-1:\widetilde J^\circ_n(j+i\ell^\circ_n)\in\partial C^\star_{n,l},X_{n,i}(\widetilde J^\circ_n(j+i\ell^\circ_n))=1\}}, \tag{10.20}$$

$$\widetilde{M}_{n,j}(k)=\textstyle\sum_{l\in\mathcal{L}_n(k)}\sum_{i=0}^{k^\circ_n(t)/\ell^\circ_n-1}\mathbb{1}_{\{\widetilde J^\circ_n(j+i\ell^\circ_n)\in\partial C^\star_{n,l},X_{n,i}(\widetilde J^\circ_n(j+i\ell^\circ_n))=1\}}. \tag{10.21}$$

Similarly, let $\widetilde N'_{n,j}(k)$ and $\widetilde{M}'_{n,j}(k)$ be defined, respectively, as $\widetilde N_{n,j}(k)$ and $\widetilde{M}_{n,j}(k)$ with $\mathcal{L}_n(k)$ replaced by $\mathcal{L}'_n(k)$.

Lemma 10.2. *For all $0\le j\le\ell^\circ_n-1$, all $2\le k\le k^\star_n$, and any $\eta,\rho,\eta',\rho'>0$,*

$$P_{\pi^\circ_n}\left(|N_{n,j}(k)-\eta|\ge\rho\right)\ \stackrel{d}{=}\ (1+\delta_n)^{\lfloor k^\circ_n(t)/\ell^\circ_n\rfloor}\widetilde P\big(\big|\widetilde N_{n,j}(k)-\eta\big|\ge\rho\big), \tag{10.22}$$

$$P_{\pi^\circ_n}\left(|M_{n,j}(k)-\eta'|\ge\rho'\right)\ \stackrel{d}{=}\ (1+\delta'_n)^{\lfloor k^\circ_n(t)/\ell^\circ_n\rfloor}\widetilde P\big(\big|\widetilde M_{n,j}(k)-\eta'\big|\ge\rho'\big), \tag{10.23}$$

for some $0\le\delta_n,\delta'_n\le 2^{-n}$, and where equality holds in distribution. The same relations hold with $M_{n,j}(k)$ and $\widetilde M_{n,j}(k)$ replaced by $M'_{n,j}(k)$ and $\widetilde M'_{n,j}(k)$, respectively.

Proof. By (10.3) and Corollary 7.5, for $0\le j\le\ell^\circ_n-1$ and all $2\le k\le k^\star_n$,

$$N_{n,j}(k)\ \stackrel{d}{=}\ \textstyle\sum_{l\in\mathcal{L}_n(k)}\mathbb{1}_{\{\exists 0\le i\le k^\circ_n(t)/\ell^\circ_n-1:J^\circ_n(j+i\ell^\circ_n)\in\partial C^\star_{n,l},J^\circ_n(j+i\ell^\circ_n+1)\in\partial C^\star_{n,l}\}}, \tag{10.24}$$

$$M_{n,j}(k)\ \stackrel{d}{=}\ \textstyle\sum_{l\in\mathcal{L}_n(k)}\sum_{i=0}^{k^\circ_n(t)/\ell^\circ_n-1}\mathbb{1}_{\{J^\circ_n(j+i\ell^\circ_n)\in\partial C^\star_{n,l},J^\circ_n(j+i\ell^\circ_n+1)\in\partial C^\star_{n,l}\}}. \tag{10.25}$$

Next note that by the Markov property and the mixing property of Proposition 6.1 we have, for all $y_i\in\mathcal{V}^\circ_n$, $y'_i\in\mathcal{V}^\circ_n$, $i=0,\dots,\lfloor k^\circ_n(t)/\ell^\circ_n\rfloor$,

$$\begin{aligned}&P^\circ_{\pi^\circ_n}\left(\cap_{0\le i\le\lfloor k^\circ_n(t)/\ell^\circ_n\rfloor}\{J^\circ_n(j+i\ell^\circ_n)=y_i,J^\circ_n(j+i\ell^\circ_n+1)=y'_i\}\right)\\ =\ &(1+\delta_n)^{\lfloor k^\circ_n(t)/\ell^\circ_n\rfloor}\textstyle\prod_{0\le i\le\lfloor k^\circ_n(t)/\ell^\circ_n\rfloor}\pi^\circ_n(y_i)p^\circ_n(y_i,y'_i),\end{aligned} \tag{10.26}$$

for some $0\le\delta_n\le 2^{-n}$, where $p^\circ_n(\cdot,\cdot)$ are the transition probabilities (3.5). From this, (10.24)-(10.25), and the relations (6.14) and (6.15), the lemma easily follows. One proves in a similar way that (10.25) holds (for some other $0\le\delta'_n\le 2^{-n}$) with $M_{n,j}(k)$ and $\widetilde M_{n,j}(k)$ replaced by $M'_{n,j}(k)$ and $\widetilde M'_{n,j}(k)$, respectively. □

Now the probabilities in the right hand sides of (10.22) and (10.23) are not hard to evaluate. Set

$$\Delta_{n,j}(k) = \widetilde{M}_{n,j}(k) - \widetilde{N}_{n,j}(k). \tag{10.27}$$

Lemma 10.3. *Let $2 < K < \infty$ be a fixed integer. There exists a subset $\Omega^{BCL} \subset \Omega$ with $\mathbb{P}(\Omega^{BCL}) = 1$ such that, on Ω^{BCL}, for all but a finite number of indices n, the following holds: there exists $\delta_n(2)$, $\eta_n(k)$, $2 \le k \le k_n^\star$, satisfying (10.10) such that, for all $0 \le j \le \ell_n^\circ - 1$ and all $m_1, m_2 > 0$,*

$$\widetilde{P}\big(\big|\Delta_{n,j}(2) - \delta_n(2)\big| \ge n^{m_2}\sqrt{\delta_n(2)}\big) \le 2n^{-2m_2}, \tag{10.28}$$

$$\widetilde{P}\big(\big|\widetilde{N}_{n,j}(k) - \eta_n(k)\big| \ge n^{m_1}\sqrt{\eta_{n,j}(2)}\big) \le 4n^{-2m_1}, \quad 2 \le k \le K, \tag{10.29}$$

$$\widetilde{P}\big(\Delta_{n,j}(k) \ge (k/n)\eta_n(k)\big) \le e^{-c''n}, \quad 3 \le k \le K, \tag{10.30}$$

$$\widetilde{P}\big(\widetilde{M}'_{n,j}(k) \ge c'_k n k_n^\star \eta_n(2+k)\big) \le 2e^{-n^2/4}, \quad 2 \le k \le K, \tag{10.31}$$

and for all $0 \le j \le \ell_n^\circ - 1$, all $2 < K < k \le k_n^\star$, and all $m_1, m_2 > 0$,

$$\widetilde{P}\big(\widetilde{N}_{n,j}(k) - n^2\eta_n(k) \ge n^{m_1+1}\sqrt{\eta_{n,j}(2)}\big) \le 4n^{-2m_1}, \tag{10.32}$$

$$\widetilde{P}\big(\Delta_{n,j}(k) \ge nk\eta_n(k)\big) \le e^{-c''n}, \tag{10.33}$$

$$\widetilde{P}\big(\widetilde{M}'_{n,j}(k) \ge c'_k n^3 k_n^\star \eta_n(2+k)\big) \le 2e^{-n^2/4}. \tag{10.34}$$

Proof. First let $k = 2$. By (10.5), $|\partial x \cap (\cup_{1 \le l \le L^\star} C^\star_{n,l})| = 1$ for all $x \in \cup_{\mathcal{L}_n(2)} \partial C^\star_{n,l}$ so that

$$P\big(\widetilde{J}_n^\circ(j + i\ell_n^\circ) \in \partial C^\star_{n,l}, X_{n,i}(\widetilde{J}_n^\circ(j + i\ell_n^\circ)) = 1\big) = 2(1 - n^{-1})/|\mathcal{V}_n^\circ|. \tag{10.35}$$

The variables $\widetilde{N}_{n,j}(2)$ and $\widetilde{M}_{n,j}(2)$ can now be expressed using random allocation of balls into urns: each set $C^\star_{n,l}$, $l \in \mathcal{L}_n(2)$, stands for an urn and a ball is allocated to $C^\star_{n,l}$ if the event $\{\widetilde{J}_n^\circ(j + i\ell_n^\circ) \in \partial C^\star_{n,l}, X_{n,i}(\widetilde{J}_n^\circ(j + i\ell_n^\circ)) = 1\}$ occurs. The total number of balls is $\widetilde{M}_{n,j}(2)$ and the number of urns is $|\mathcal{L}_n(2)|$. Then $\widetilde{N}_{n,j}(2) = |\mathcal{L}_n(2)| - U_n$, where $U_n \equiv U_{n,j}(2)$ denotes the number of empty urns obtained by distributing $M_n \equiv \widetilde{M}_{n,j}(2)$ balls into $L_n \equiv |\mathcal{L}_n(2)|$ urns at random, uniformly. The next lemma is classical.

Lemma 10.4 ([17] Section 9.2, Theorem 3)**.** *Set $A_n \equiv \frac{M_n}{L_n}$ and $B_n \equiv (e^{A_n} - A_n - 1)^{1/2}$. If $M_n \to \infty$ and $A_n/L_n \to 0$ as $n \to \infty$ then, for all $\delta > 0$,*

$$P\left(|U_n - E\left[U_n\right]| > \delta V_n\right) \le \delta^{-2}, \tag{10.36}$$

where $E\left[U_n\right] = L_n e^{-A_n}(1 + \mathcal{O}(\frac{A_n}{L_n}))$ and $V_n^2 = L_n e^{-2A_n} B_n^2(1 + o(1))$.

It remains to evaluate the quantities entering in Lemma 10.4. The following lemma anticipates our needs.

Lemma 10.5. *Let $2 < K < \infty$ be a fixed integer. If $c_\star > 3$, there exists a subset $\Omega^{BCL} \subset \Omega$ with $\mathbb{P}(\Omega^{BCL}) = 1$ such that, on $\Omega^\star \cap \Omega^{BCL}$, for all but a finite number of indices n the following holds: for all $2 \le k \le K$, $|\mathcal{L}_n(k)| = c_k(k-1)!2^{n-1}n^{-k(c_\star-1)-1}(1 + o(1))$ and $|\mathcal{L}'_n(k)| \le c'_k(k+2)!2^{n-1}n^{-(k+2)(c_\star-1)-1}$ while for all $K < k \le k_n^\star$, $|\mathcal{L}_n(k)| \le c_k(k-1)!2^{n-1}n^{-k(c_\star-1)+1}(1 + o(1))$ and $|\mathcal{L}'_n(k)| \le c'_k(k+2)!2^{n-1}n^{-(k+2)(c_\star-1)+1}$ where $0 < c_k, c'_k < \infty$ are constants and $c_2 = 1$.*

Proof. If $k \le K < \infty$ proceed as in the proof of Lemma 9.9 (see (9.99)-(9.105)) with appropriate choices of the decomposition (9.101)-(9.102). Otherwise, use a first order Tchebychev inequality. These are by now routine calculations which we leave out. Note that working out an estimate on $|\mathcal{L}_n(k)|$ when $k \equiv k_n$ is diverging with n is a hard task,

while working out an upper bound on $\mathbb{E}|\mathcal{L}_n(k)|$ is elementary. This is what leads us we distinguish the cases $2 \le k \le K$ and $K < k \le k_n^\star$ in Proposition 10.1. □

Remark. It follows from Lemma 10.5 that if $c_\star > 3$, the collection $C_{n,l}^\star$, $1 \le l \le L^\star$, is dominated by "isolated pairs", namely, by sets of size two that are at distance larger than three from all other sets. This is not true for smaller values of $c_\star$.

In view of (10.8)-(10.9), $\widetilde{M}_{n,j}(2)$ is a sum of $|\mathcal{L}_n(2)|\,(\lfloor k_n^\circ(t)/\ell_n^\circ\rfloor - 1)$ i.i.d. Bernoulli r.v.'s with success probability $2(1-\frac{1}{n})/|\mathcal{V}_n^\circ|$. Thus, on Ω^{BCL}, for n large enough, $E\widetilde{M}_{n,j}(2) = 2\frac{|\mathcal{L}_n(2)|}{|\mathcal{V}_n^\circ|}(1-\frac{1}{n})(\lfloor\frac{k_n^\circ(t)}{\ell_n^\circ}\rfloor - 1) = \frac{\lfloor a_n t\rfloor}{\ell_n^\circ n^{2c_\star-1}}(1+o(1))$, while by Bennett's bound (9.103),

$$P\left(\left|\widetilde{M}_{n,j}(2) - E\widetilde{M}_{n,j}(2)\right| < n\big(E\widetilde{M}_{n,j}(2)\big)^{1/2}\right) \ge 1 - 2e^{-n^2/4}. \tag{10.37}$$

Conditionnal on the event appearing in the probability (10.37), $\widetilde{M}_{n,j}(2) \to \infty$ as $n \to \infty$ and $A_n \equiv \frac{M_{n,j}(2)}{|\mathcal{L}_n(2)|} = \frac{\lfloor a_n t\rfloor}{\ell_n^\circ 2^{n-1}}(1+o(1)) \le n^{-3}t2^{-n(1-\varepsilon)} \to 0$ for all $t \le T < \infty$ and $1 < \varepsilon < 1$. Thus $A_n/|\mathcal{L}_n(2)| \to 0$ as well. Thus the conditions of Lemma 10.4 are satisfied, and so, on Ω^{BCL}, for all but a finite number of indices n, for all $m_1, m_2 > 0$, setting $\Delta_{n,j}(2) \equiv \widetilde{M}_{n,j}(2) - \widetilde{N}_{n,j}(2)$,

$$\widetilde{P}\left(\left|\widetilde{N}_{n,j}(2) - \widetilde{E}\widetilde{N}_{n,j}(2)\right| \ge n^{m_1}\big(\widetilde{E}\widetilde{N}_{n,j}(2)\big)^{1/2}\right) \le 2n^{-2m_1} + 2e^{-n^2/4}, \tag{10.38}$$

$$\widetilde{P}\left(\left|\Delta_{n,j}(2) - \widetilde{E}\Delta_{n,j}(2)\right| \ge n^{m_2}\big(\widetilde{E}\Delta_{n,j}(2)\big)^{1/2}\right) \le 2n^{-2m_2} + 2e^{-n^2/4}, \tag{10.39}$$

where $\widetilde{E}\widetilde{N}_{n,j}(2) = \frac{\lfloor a_n t\rfloor}{\ell_n^\circ n^{2c_\star-1}}(1+o(1))$ $\widetilde{E}\Delta_{n,j}(2) = \frac{\lfloor a_n t\rfloor^2}{(\ell_n^\circ)^2 2^n n^{2c_\star-1}}(1+o(1))$. This proves (10.28) and (10.29) with $k = 2$.

If $k > 2$, $|\partial x \cap (\cup_{1\le l\le L^\star} C_{n,l}^\star)| = |\partial x \cap C_{n,l}^\star|$ for all $x \in \partial C_{n,l}^\star$, $l \in \mathcal{L}_n(k)$, but $|\partial x \cap C_{n,l}^\star|$ is not always one. However, at most k vertices of $\partial C_{n,l}^\star$ can have more than one neighbor in $C_{n,l}^\star$, and a given vertex cannot have more than k neighbors. Thus, denoting by $\partial C_{n,l}^{\star,+}$ and $\partial C_{n,l}^{\star,-}$ the subsets where $|\partial x \cap C_{n,l}^\star| = 1$ and $|\partial x \cap C_{n,l}^\star| > 1$, respectively, it follows from (2.15) that on $\Omega^\star$, $|\partial C_{n,l}^{\star,+}| \ge |\partial C_{n,l}^\star| - k > k(n-1)$ while $|\partial C_{n,l}^{\star,+}| \le k$. Then $\widetilde{N}_{n,j}^+(k) \le \widetilde{N}_{n,j}(k) \le \widetilde{N}_{n,j}^+(k) + \widetilde{N}_{n,j}^-(k)$ where $\widetilde{N}_{n,j}^\pm(k)$ are defined as $\widetilde{N}_{n,j}(k)$ with $\partial C_{n,l}^\star$ replaced by $\partial C_{n,l}^{\star,\pm}$. With obvious notation we write $\widetilde{M}_{n,j}^\pm(k)$ and $\widetilde{\Delta}_{n,j}^\pm(k)$.

At this point we must distinguish the cases $k \le K < \infty$ and $K < k \le k_n^\star$. Assume first that $k \le K < \infty$. To bound $\widetilde{N}_{n,j}^+(k)$ we proceed exactly as for $\widetilde{N}_{n,j}(2)$: the r.h.s. of (10.35) is now equal to $k(1-o(1))/|\mathcal{V}_n^\circ|$, and we get that $\widetilde{N}_{n,j}^+(k)$ obeys (10.37) - (10.39), under the same assumptions, with $\widetilde{E}\Delta_{n,j}^+(k) = \frac{c_k(k-1)!(k\lfloor a_n t\rfloor)^2}{(\ell_n^\circ)^2 2^{n+2} n^{k(c_\star-1)+1}}(1+o(1))$ and $\widetilde{E}\widetilde{N}_{n,j}^+(k) = \frac{c_k k!\lfloor a_n t\rfloor}{\ell_n^\circ 2^{n^{k(c_\star-1)+1}}}(1+o(1))$. Using that the l.h.s. of (10.35) is bounded above by $k^2(1-o(1))/n|\mathcal{V}_n^\circ|$ it also follows from (10.37) that

$$\widetilde{P}\left(\widetilde{M}_{n,j}^-(k) \ge \tfrac{k}{n}E\widetilde{M}_{n,j}(k) + n\big(E\widetilde{M}_{n,j}(k)\big)^{1/2}\right) \le 2e^{-n^2/4}. \tag{10.40}$$

Combining these estimates and the bound $\widetilde{\Delta}_{n,j}(k) \le \widetilde{\Delta}_{n,j}^+(k) + \widetilde{M}_{n,j}^-(k)$, yield (10.29) and (10.30) for all $3 \le k < K$. The same arguments apply to the case $K < k \le k_n^\star$ but since Lemma 10.5 now only provides an upper bound on $|\mathcal{L}_n(k)|$, they only yield one-sided inequalities, namely, (10.32) and (10.33).

It remains to bound $M'_{n,j}(k)$. For this we simply use that by (2.15), on $\Omega^\star$, the l.h.s. of (10.35) is bounded above by $P(\widetilde{J}_n^\circ(j+i\ell_n^\circ) \in \partial C_{n,l}^\star) \le nk_n^\star(1-o(1))/|\mathcal{V}_n^\circ|$ and, proceeding

as in the proof of (10.37), we obtain (10.31) and (10.34). The proof of Lemma 10.3 is done. □

Combining the estimates of Lemma 10.3 together with (10.22)-(10.23) and choosing $m_1 = m_2 = 2c_\star$ now easily yields (10.17). To prove (10.18) write

$$P^\dagger_{\pi^\circ_n}\left(m_{n,j} \geq \kappa\right) \leq \textstyle\sum_{1\leq l\leq L^\star} P^\dagger_{\pi^\circ_n}\left(\sum_{i\in I_j} \mathbb{1}_{\{J^\dagger_n(i+1)\in C^\star_{n,l}\}} \geq \kappa\right). \tag{10.41}$$

Using Corollary 7.5, the mixing property of Proposition 6.1, and observing that $k^\circ_n(t) = \lfloor a_n t\rfloor \leq \lfloor a_n T\rfloor$, one sees that (10.41) is bounded above by

$$(1+o(1)) \sum_{1\leq l\leq L^\star} P^\dagger_{\pi^\circ_n}\left(\textstyle\bigcup_{0\leq i_1<\cdots<i_\kappa\leq \lfloor a_nT\rfloor/\ell^\circ_n-1} \bigcap_{k=1}^\kappa \{\widetilde{J}^\circ_n(j+i_k\ell^\circ_n)\in \partial C^\star_{n,l},\}\right). \tag{10.42}$$

Using again that on $\Omega^\star$, $\widetilde{P}(\widetilde{J}^\circ_n(j+i\ell^\circ_n)\in\partial C^\star_{n,l}) \leq nk^\star_n(1-o(1))/|\mathcal{V}^\circ_n|$, and since $L^\star < 2^n$,

$$P^\dagger_{\pi^\circ_n}\left(m_{n,j}\geq\kappa\right) \leq L^\star\left(\tfrac{nk^\star_n\lfloor a_nT\rfloor}{\ell^\circ_n|\mathcal{V}^\circ_n|}\right)^\kappa \leq (1+o(1))\,(T/n)^\kappa\, 2^{-n\{\kappa(1-\varepsilon)-1\}}. \tag{10.43}$$

Choosing $\kappa = \frac{3}{1-\varepsilon}$ in (10.43) implies (10.18). The proof of Proposition 10.1 is done. □

10.2. **Step two.** Building on Step 1 we now aim at proving that:

Proposition 10.6. *Assume that $c_\star > 3$ and set $b_n = a_n \exp(n(\beta/2)^2)/(\beta\sqrt{\pi n})$. For all $0<\beta\leq 2\beta_c(\varepsilon/2) - 12\sqrt{c_\star \log n/n}$, all $t\in[0,T]$, and large enough n,*

$$\mathbb{P}\left\{P^\dagger_{\pi^\circ_n}\left(\left|S^\dagger_n(t)-t\right| \geq c_1\sqrt{t/n} + c_2 t n^{-(c_\star-1)}\right) \leq n^{-2}\right\} > 1-n^{-2} \tag{10.44}$$

for some constants $c_1, c_2 > 0$.

Proof. By definition of the sets I_j of Step 1, assuming again that $\lfloor k^\circ_n(t)/\ell^\circ_n\rfloor = k^\circ_n(t)/\ell^\circ_n$,

$$S^\dagger_n(t) = \frac{1}{\ell^\circ_n}\sum_{j=0}^{\ell^\circ_n-1} S^\dagger_{n,j}(t), \quad S^\dagger_{n,j}(t) \equiv \frac{\ell^\circ_n}{b_n}\sum_{i\in I_j}\Lambda^\dagger_n(J^\dagger_n(i+1)). \tag{10.45}$$

Note that each realisation of the sets $\{C^\star_{n,l}, 1\leq l\leq L^\star\}$ induces a partition of $\{1,\dots,L^\star\}$ into $\cup_{2\leq k\leq k^\star_n}(\mathcal{L}_n(k)\cup\mathcal{L}'_n(k))$ and, given this partition, each I_j can be decomposed into disjoint sets, $I_j = \cup_{2\leq k\leq k^\star_n}(I_{j,0}(k)\cup I_{j,1}(k)\cup I'_j(k))$, with the properties that:
(a) $\left\{J^\dagger_n(i+1), i\in I_{j,0}(k)\right\}$ consists of $N_{n,j}(k)$ visits to distinct sets $C^\star_{n,l}, l\in\mathcal{L}_n(k)$;
(b) $\left\{J^\dagger_n(i+1), i\in I_{j,1}(k)\right\}$ consists of $M_{n,j}(k)-N_{n,j}(k)$ visits to sets $C^\star_{n,l}, l\in\mathcal{L}_n(k)$, each $C^\star_{n,l}$ being visited at most $m_{n,j}$ times;
(c) $\left\{J^\dagger_n(i+1), i\in I'_j(k)\right\}$ consists of $M'_{n,j}(k)$ visits to sets $C^\star_{n,l}, l\in\mathcal{L}'_n(k)$, each $C^\star_{n,l}$ being visited at most $m_{n,j}$ times.

Denoting by $S^k_{n,j,0}$, $S^k_{n,j,1}$, and $S'^k_{n,j}$, $2\leq k\leq k^\star_n$, the restrictions of the sum in $S^\dagger_{n,j}$ to, respectively, $I_{j,0}(k)$, $I_{j,1}(k)$, and $I'_j(k)$, $2\leq k\leq k^\star_n$, we have

$$S^\dagger_{n,j}(t) = \textstyle\sum_{k=2}^{k^\star_n}\left[S^k_{n,j,0}(t) + S^k_{n,j,1}(t) + S'^k_{n,j}(t)\right]. \tag{10.46}$$

Observe that by (a), each $S^k_{n,j,0}$ is a sum of $N_{n,j}(k)$ independent rv's, and by (b) and (c) each $S^k_{n,j,1}$, respectively $S'^k_{n,j}$, is bounded above by $m_{n,j}$ times a sum of (at most) $M_{n,j}(k)-N_{n,j}(k)$, respectively $M'_{n,j}(k)$, independent rv's. Before making these bounds explicit we state two lemmata.

Let $\mathbb{P}_\mathcal{C}$ denote the conditional distribution $\mathbb{P}(\cdot \mid \{C^\star_{n,l}, 1\leq l\leq L^\star\})$.

Lemma 10.7. *Let $\bar a_n > 0$ be an increasing sequence such that $0 < \bar\varepsilon_n \equiv \frac{\log \bar a_n}{n \log 2} < 1$. Given $\{l_i, 1 \le i \le \lfloor \bar a_n t\rfloor\} \subseteq \mathcal{L}_n(2)$ let $(G_{n,i}, 1 \le i \le \lfloor \bar a_n t\rfloor)$ be independent geometric r.v.'s satisfying $P_G(G_{n,i} = k) = p_{n,i}(1-p_{n,i})^{k-1}, k \ge 1$, where $(p_{n,i})^{-1} = 1 + (n-1)^{-1}\varrho_{n,l_i}(0)$ for $\varrho_{n,l_i}(0)$ as in (5.2). Let P_G be the joint law of $(G_{n,i}, 1 \le i \le \lfloor \bar a_n t\rfloor)$. Set $\bar\beta = \beta/\sqrt{2}$ and, given a constant $\bar\kappa > 0$, set $s_n = n^{2\bar\kappa} e^{n\bar\beta\beta_c(\bar\varepsilon_n)}$ and $\bar b_n = \bar a_n \mathbb{E}_{\mathcal{C}} E_G(G_{n,i}\mathbb{1}_{\{G_{n,i}<s_n\}})$. Then, for all $\bar\beta \le \beta_c(\bar\varepsilon_n) - \sqrt{8\bar\kappa \log n/n}$, all $t \in [0,T]$, and any sequence $\bar\delta_n > n^{-\bar\kappa}$,*

$$\mathbb{P}_{\mathcal{C}}\left\{P_G\left(\left|\tfrac{1}{\bar b_n}\textstyle\sum_{i=1}^{\lfloor \bar a_n t\rfloor} G_{n,i} - \frac{\lfloor \bar a_n t\rfloor}{\bar a_n}\right| \ge \sqrt{t\bar\delta_n}\right) \le 2\bar\delta_n^{-1} n^{-\bar\kappa}\right\} \ge 1 - n^{-\bar\kappa + 2(c_\star+1)},$$

and $\mathbb{E}_{\mathcal{C}} E_G(G_{n,i}\mathbb{1}_{\{G_{n,i}<s_n\}}) = (1+o(1))\frac{n^{2c_\star-1}}{\bar\beta\sqrt{2\pi n}} e^{n\bar\beta^2/2}$.

Lemma 10.8. *Let $\bar a_n > 0$ be an increasing sequence such that $0 < \bar\varepsilon_n \equiv \frac{\log \bar a_n}{n \log 2} < 1$. Given $k > 2$ and $\{l_i, 1 \le i \le \lfloor \bar a_n t\rfloor\} \subseteq \mathcal{L}_n(k)$ let $(Y^k_{n,i}, 1 \le i \le \lfloor \bar a_n t\rfloor)$ be i.i.d. integer valued r.v.'s satisfying $P_Y(Y^k_{n,i} > m) \le e^{-m(n/k\varrho_{n,l_i}(0))(1-o(1))}(1+o(1))$ for all $m \ge \theta^\star_n$, where $\varrho_{n,l_i}(0)$ and $\theta^\star_n$ are as in (5.2)-(5.4). Let P_Y be the joint law of $(Y^k_{n,i}, 1 \le i \le \lfloor \bar a_n t\rfloor)$.*
(i) Given $\bar\kappa > 0$, let $\bar\beta$ and s_n be as in Lemma 10.7 and set $\bar b_n = \bar a_n \mathbb{E}_{\mathcal{C}} E_Y(Y^k_{n,i}\mathbb{1}_{\{Y^k_{n,i}<s_n\}})$. Then, for all $\bar\beta \le \beta_c(\bar\varepsilon_n) - \sqrt{8\bar\kappa \log n/n}$, all $t \in [0,T]$, and any sequence $\bar\delta_n > n^{-\bar\kappa}$,

$$\mathbb{P}_{\mathcal{C}}\left\{P_Y\left(\tfrac{1}{\bar b_n}\textstyle\sum_{i=1}^{\lfloor \bar a_n t\rfloor} Y^k_{n,i} - \frac{\lfloor \bar a_n t\rfloor}{\bar a_n} \ge \sqrt{t\bar\delta_n}\right) \le 2\bar\delta_n^{-1} n^{-\bar\kappa}\right\} \ge 1 - n^{-\bar\kappa + 2(c_\star+2)},$$

and $(1+o(1))\frac{k}{\bar\beta\sqrt{\pi n}} n^{2c_\star-1} e^{n\bar\beta^2/2} \le \mathbb{E}_{\mathcal{C}} E_Y(Y^k_{n,i}\mathbb{1}_{\{Y^k_{n,i}<s_n\}}) \le (1+o(1))\frac{k^2(k-1)}{2\bar\beta\sqrt{2\pi n}} n^{2c_\star-1} e^{n\bar\beta^2/2}$.
(ii) For all $\beta > 0$, all $t \in [0,T]$, and any $\bar\kappa', \bar\kappa'' > 0$, taking $\bar b_n = \bar a_n \mathbb{E}_{\mathcal{C}} E_Y(Y^k_{n,i})$,

$$\mathbb{P}_{\mathcal{C}}\left\{P_Y\left(\tfrac{1}{\bar b_n}\textstyle\sum_{i=1}^{\lfloor \bar a_n t\rfloor} Y^k_{n,i} \ge t n^{\bar\kappa' + \bar\kappa''}\right) \le n^{-\bar\kappa'}\right\} \ge 1 - n^{-\bar\kappa''},$$

and $(1+o(1))\frac{k}{\bar\beta\sqrt{\pi n}} n^{2c_\star-1} e^{n\bar\beta^2/2} \le \mathbb{E}_{\mathcal{C}} E_Y(Y^k_{n,i}) \le (1+o(1))\frac{k^2(k-1)}{2\bar\beta\sqrt{\pi n}} n^{2c_\star-1} e^{n\bar\beta^2/2}$.

Proof of Lemmata 10.7 and 10.8. We only indicate the main lines of the proofs. Consider first Lemma 10.7. Set $G'_{n,i} = G_{n,i}\mathbb{1}_{\{G_{n,i}<s_n\}}$. Then, for all $\epsilon > 0$,

$$P_G\left(\left|\tfrac{1}{\bar b_n}\textstyle\sum_{i=1}^{\lfloor \bar a_n t\rfloor} G_{n,i} - \sigma_n\right| \ge \epsilon\right) \le \epsilon^{-2}\bar\sigma_n + \hat\sigma_n, \tag{10.47}$$

where $\bar b_n \sigma_n = E_G \sum_{i=1}^{\lfloor \bar a_n t\rfloor} G'_{n,i}$, $\bar b_n^2 \bar\sigma_n = E_G \sum_{i=1}^{\lfloor \bar a_n t\rfloor} G'^2_{n,i}$, and $\hat\sigma_n = \sum_{i=1}^{\lfloor \bar a_n t\rfloor} P_G(G_{n,i} > s_n)$. Now $E_G G'_{n,i} = \frac{1}{p_{n,i}}\left(1 - (1-p_{n,i})^{s_n-1}\right)$, $E_G G'^2_{n,i} \le 2\frac{1-p_{n,i}}{p^2_{n,i}}\left(1 - (1-p_{n,i})^{s_n+1}\right)$, and $P_G(G_{n,i} > s_n) = (1-p_{n,i})^{s_n+1}$. Thus each of the σ_n's in (10.47) is a sum of a function of the variables $\varrho_{n,l_i}(0)$. Note that $\mathbb{P}_{\mathcal{C}}\left(|\sigma_n - \mathbb{E}_{\mathcal{C}}\sigma_n| \ge \epsilon'\right) \le \epsilon'^{-2}\mathbb{E}_{\mathcal{C}}\bar\sigma_n$ and, by first order Tchebychev inequalities, $\mathbb{P}_{\mathcal{C}}(\bar\sigma_n > \epsilon') \le \epsilon'^{-1}\mathbb{E}_{\mathcal{C}}\bar\sigma_n$ and $\mathbb{P}_{\mathcal{C}}(\hat\sigma_n > \epsilon') \le \epsilon'^{-1}\mathbb{E}_{\mathcal{C}}\hat\sigma_n$. Clearly, $\mathbb{E}_{\mathcal{C}}\sigma_n = \lfloor \bar a_n t\rfloor/\bar a_n$, while $\mathbb{E}_{\mathcal{C}}\bar\sigma_n \le n^{-2\bar\kappa+2c_\star+3/2}$ and $\mathbb{E}_{\mathcal{C}}\hat\sigma_n \le n^{-2(\bar\kappa-1)\beta_c(\bar\varepsilon)/\bar\beta+2c_\star}$ for all $\bar\beta \le \beta_c(\bar\varepsilon_n) - \sqrt{8\bar\kappa\log n/n}$. We do not reproduce the lenghty but simple calculations leading to these bounds. They resemble the high temperature moments estimates of the partition function of the REM: here the quantity

$$p^{-1}_{n,i} = 1 + (n-1)^{-1} e^{-\beta \max\{(H_n(y), H_n(x)) \,|\, \{x,y\} \in C^\star_{n,l_i}\}} \tag{10.48}$$

plays the role of the Boltzmann weights, the gist of the argument being that the truncation of the $G_{n,i}$'s at level s_n effectively results in a truncation of the $p^{-1}_{n,i}$'s at level s_n, and that s_n is, roughly speaking, of the order of the maximum of the $p^{-1}_{n,i}$'s. Based on this (10.47) is easily worked out.

To prove Lemma 10.8, (i), replace $G_{n,i}$ by $Y^k_{n,i}$ and $G'_{n,i}$ by $Y'_{n,i} = Y^k_{n,i}\mathbb{1}_{\{Y^k_{n,i}<s_n\}}$ in the above expressions of s_n, $\bar{s}_n$, and $\hat{s}_n$, and use that $\varrho_{n,l_i}(0) \geq e^{-\beta\max(H_n(y),H_n(x))}$ and $\varrho_{n,l_i}(0) \leq \sum_{\{x,y\}\in G(C^\star_{n,l_i})} e^{-\beta\max(H_n(y),H_n(x))}$ for any $\{x,y\} \in G(C^\star_{n,l_i})$. Again $\mathbb{E}_{\mathcal{C}}\sigma_n = \lfloor \bar{a}_n t\rfloor/\bar{a}_n$ while $\mathbb{E}_{\mathcal{C}}\bar{\sigma}_n \leq k^{-2}n^{-2\bar{\kappa}+2c_\star+5/2}$ and $\mathbb{E}_{\mathcal{C}}\hat{\sigma}_n \leq k^{2-\beta_c(\bar{\varepsilon})/\bar{\beta}}n^{-2(\bar{\kappa}-1)\beta_c(\bar{\varepsilon})/\bar{\beta}-1/2+2c_\star}$. Assertion (ii) of the lemma follows from two first order Tchebychev inequalities applied in a row. □

We are now equipped to estimate the sum (10.46). Consider first $S^2_{n,j,0}(t)$. By (a), there exists a subset $\{l_i, 1 \leq i \leq N_{n,j}(2)\} \subseteq \mathcal{L}_n(2)$ such that

$$S^2_{n,j,0}(t) = \tfrac{\ell^\circ_n}{b_n}\sum_{i=1}^{N_{n,j}(2)} \Lambda^\dagger_n(J^\dagger_n(i))\mathbb{1}_{\{J^\dagger_n(i)\in C^\star_{n,l_i}\}} \stackrel{d}{=} \tfrac{\ell^\circ_n}{b_n}\sum_{i=1}^{N_{n,j}(2)} G_{n,i}, \tag{10.49}$$

where $G_{n,i}$ is as in Lemma 10.7. The last equality follows from Proposition 5.1, (i), and holds in distribution. In the same way we have, by Proposition 5.1, (ii), that on $\Omega^\star \cap \Omega_0$,

$$S^k_{n,j,0}(t) \stackrel{d}{=} \tfrac{\ell^\circ_n}{b_n}\sum_{i=1}^{N_{n,j}(k)} Y^k_{n,i}, \quad 2 < k \leq k^\star_n, \tag{10.50}$$

where $Y^k_{n,i}$ is as in Lemma 10.8. On the one hand Lemma 10.7 combined with Proposition 10.1 yields an accurate estimate on the last sum in (10.49) for all $\beta \leq 2\beta_c(\varepsilon/2) - 4\sqrt{\bar{\kappa}\log n/n} - 5(c_\star+1)\log n/n$. On the other hand, the sum in the r.h.s. of (10.50) can be bounded from above using Lemma 10.8 and Proposition 10.1 . Note however that the temperature threshold in Lemma 10.8, (i), is a decreasing function of $\bar{a}_n$. Namely, for $\bar{a}_n = N_{n,j}(k)$, by (10.11)-(10.14), $\beta_c(\bar{\varepsilon}_n) - \sqrt{8\bar{\kappa}\log n/n} = 2\beta_c(\varepsilon/2) - 4\sqrt{\bar{\kappa}\log n/n} - \mathcal{O}(k\log n/n)$. We will thus use Lemma 10.8, (i), when e.g. $k \leq n^{1/4}$, and Lemma 10.8, (ii), else. Doing so we obtain that on $\Omega_0\cap\Omega_1\cap\Omega^\star\cap\Omega^{\text{BCL}}$, choosing

$$b_n = \ell^\circ_n\mathbb{E}_{\mathcal{C}}N_{n,j}(2)E_G(G_{n,i}\mathbb{1}_{\{G_{n,i}<s_n\}})\Big|_{t=1} = \tfrac{a_n}{(\beta/\sqrt{2})\sqrt{2\pi n}}e^{n(\beta/2)^2}(1+o(1)), \tag{10.51}$$

for all $0 < \beta \leq 2\beta_c(\varepsilon/2) - 4\sqrt{\bar{\kappa}\log n/n}(1+o(1))$, any $\bar{\kappa}, \bar{\kappa}', \bar{\kappa}'' > 0$, and any $\bar{\delta}_n > n^{-\bar{\kappa}}$, with $\mathbb{P}$- probability larger than $1 - k^\star_n n^{-\bar{\kappa}+2(c_\star+1)} - k^\star_n n^{-\bar{\kappa}''}$,

$$P^\dagger_{\pi^\circ_n}\left(\left|\sum_{k=2}^{k^\star_n} S^k_{n,j,0}(t) - t\right| \geq \sqrt{t\bar{\delta}_n}(1+\epsilon_{n,1}) + t\epsilon_{n,1}\right) \leq 2\bar{\delta}_n^{-1}k^\star_n n^{-\bar{\kappa}} + k^\star_n n^{-\bar{\kappa}'}, \tag{10.52}$$

for some constant $c_{j,1} > 0$.

Let us now establish that the contributions of the terms $S^k_{n,j,1}(t)$ and $S'^k_{n,j}(t)$ in (10.46) are subleading. Consider first $S^k_{n,j,1}(t)$, $k > 2$. Let $\{l_i, 1 \leq i \leq \Delta_{n,j}(k)\} \subseteq \mathcal{L}_n(k)$, $\Delta_{n,j}(k) \leq M_{n,j}(k) - N_{n,j}(k)$, be the labels of the distinct sets $C^\star_{n,l}, l \in \mathcal{L}_n(k)$, that $J^\dagger_n(i+1)$ visits as i runs through $I_{j,1}(k)$, and let $\{\alpha_{n,l_i}, 1 \leq i \leq \Delta_{n,j}(k)\}$ be the number of visits to each $C^\star_{n,l_i}$. Reasoning as in (10.50),

$$S^k_{n,j,1}(t) \stackrel{d}{=} \tfrac{\ell^\circ_n}{b_n}\sum_{i=1}^{\Delta_{n,j}(k)} \alpha_{n,l_i}Y^k_{n,i} \leq m_{n,j}\tfrac{\ell^\circ_n}{b_n}\sum_{i=1}^{M_{n,j}(k)-N_{n,j}(k)} Y^k_{n,i}, \tag{10.53}$$

where the last inequality is deterministic and follows from the fact that $0 < \alpha_{n,l_i} \leq m_{n,j}$ (see (10.9)) and $Y^k_{n,i} \geq 0$. In the same way, for $k > 2$, the sum $S'^k_{n,j}(t)$ obeys

$$S'^k_{n,j}(t) \stackrel{d}{=} \tfrac{\ell^\circ_n}{b_n}\sum_{i=1}^{\Delta'_{n,j}(k)} \alpha'_{n,l_i}Y^k_{n,i} \leq m_{n,j}\tfrac{\ell^\circ_n}{b_n}\sum_{i=1}^{M'_{n,j}(k)} Y^k_{n,i}, \tag{10.54}$$

for some $0 < \alpha'_{n,l_i} \leq m_{n,j}$ and $\Delta'_{n,j}(k) < M'_{n,j}(k)$. If $k = 2$ substitute $G_{n,i}$ for $Y^k_{n,i}$ in (10.53) and (10.54). Using again Lemma 10.7, Lemma 10.8, and Proposition 10.1, we obtain that under the assumption of Proposition 10.1, on $\Omega_0\cap\Omega_1\cap\Omega^\star\cap\Omega^{\text{BCL}}$, for b_n as in

(10.51), for all $0 < \beta \leq 2\beta_c(\varepsilon/2) - 4\sqrt{\bar\kappa \log n/n}(1+o(1))$, any $\bar\kappa, \bar\kappa', \bar\kappa'' > 0$, and any $\bar\delta_n > n^{-\bar\kappa}$, with $\mathbb{P}$- probability larger than $1 - 2k_n^\star n^{-\bar\kappa+2(c_\star+1)} - 2k_n^\star n^{-\bar\kappa''}$,

$$P^\dagger_{\pi_n^\circ}\left(\textstyle\sum_{k=2}^{k_n^\star}(S^k_{n,j,1}(t) + S'^k_{n,j}(t)) \geq (t+\sqrt{t\bar\delta_n})\epsilon_{n,2}\right) \leq 4\bar\delta_n^{-1}k_n^\star n^{-\bar\kappa} + 2_n^\star n^{-\bar\kappa'}, \quad (10.55)$$

where $\epsilon_{n,2} \leq c_{j,2}n^{-c_\star}(1+n^{-K(c_\star-1)+2}) + a_n 2^{-n+1} + c_{n,3}(n^{-2(c_\star-2)} + n^{-K(c_\star-1)+4})$ for some constants $c_{j,2}, c_{n,3} > 0$.

Combining (10.52) and (10.55) yields an estimate on $S^\dagger_{n,j}(t)$ for each $0 \leq j \leq \ell_n^\circ - 1$, and inserting in (10.45) we arrive at:

Lemma 10.9. *Let b_n be as in (10.51). Under the assumption of Proposition 10.1, on $\Omega_0\cap\Omega_1\cap\Omega^\star\cap\Omega^{BCL}$, the following holds: for all $0 < \beta \leq 2\beta_c(\varepsilon/2) - 4\sqrt{\bar\kappa\log n/n}(1+o(1))$, all $t \in [0,T]$, any $\bar\kappa, \bar\kappa', \bar\kappa'' > 0$, and any sequence $\bar\delta_n > n^{-\bar\kappa}$, with $\mathbb{P}$- probability larger than $1 - 4\ell_n^\circ k_n^\star n^{-\bar\kappa+2(c_\star+1)} - 4\ell_n^\circ k_n^\star n^{-\bar\kappa''}$,*

$$P^\dagger_{\pi_n^\circ}\left(\left|S^\dagger_n(t) - t\right| \geq \sqrt{t\bar\delta_n}(1+\epsilon_{n,0}) + t\epsilon_{n,0}\right) \leq 6\bar\delta_n^{-1}\ell_n^\circ k_n^\star n^{-\bar\kappa} + 4\ell_n^\circ k_n^\star n^{-\bar\kappa'}, \quad (10.56)$$

$\epsilon_{n,0} \leq c_{j,0}n^{-(c_\star-1)}(1+n^{-K(c_\star-1)+2})$ for some constant $c_{j,0} > 0$ and $K > 2$.

Choosing $\bar\delta_n = 1/n$, $\bar\kappa > 6 + 2(c_\star+1)$, and $\bar\kappa' = \bar\kappa'' > 6$ in Lemma 10.9, and taking $\bar c = 1$ in (10.17) yields the statement of Proposition 10.6. □

Proof of Theorem 3.5. Theorem 3.5 immediately follows from Proposition 10.6 since $S^\dagger_n(t)$ is a monotonous function whose limit is itself continuous. □

11. Appendix B: Speed of convergence to Perron projector.

All the matrix analysis results quoted in this section are well known and can be found in [29] (see in particular Subsection 8.5 p. 515 for nonnegative and primitive matrices).

11.1. **The case of nonnegative and primitive matrices.** Let $A = (a(i,j))_{1\leq i,j\leq N}$ be an $N\times N$ nonnegative and primitive matrix, that is to say, $A \geq 0$ and $A^k > 0$ for some $k \geq 1$. By Perron's theorem its eigenvalues satisfy

$$\lambda_0 > |\lambda_1| \geq |\lambda_2| \geq \cdots \geq |\lambda_{N-1}| \text{ and } \lambda_0 > 0, \quad (11.1)$$

and there exists a unique vector $u > 0$ such that $uA = \lambda_0 u$, and a unique vector $v > 0$ such that $Av = \lambda_0 v$, normalized such that

$$\textstyle\sum_{1\leq i\leq N} u(i) = 1, \sum_{1\leq i\leq N} u(i)v(i) = 1; \quad (11.2)$$

these are the left and right Perron eigenvectors of A. Futhermore, denoting by

$$L = (v(i)u(j))_{1\leq i,j\leq N} \quad (11.3)$$

the associated projector, we have $\lim_{m\to\infty}(\lambda_0^{-1}A)^m = L$. The lemma below provides an explicit bound on the speed of this convergence under an additionnal symmetry (or "reversibility") assumption. Write $A^m = (a^{(m)}(i,j))_{1\leq i,j\leq N}$.

Lemma 11.1. *Assume that there exists a vector $\pi_A > 0$ such that $\sum_{1\leq i\leq N}\pi_A(i) = 1$, and*

$$\pi_A(i)a(i,j) = \pi_A(j)a(j,i) \quad \textit{for all } 1 \leq i,j \leq N. \quad (11.4)$$

Then, for all $m \in \mathbb{N}$ and all $1 \leq i \leq N$, we have

$$\left|\textstyle\sum_{1\leq j\leq N} a^{(m)}(i,j) - \lambda_0^m v(i)\right| \leq |\lambda_1|^m/\sqrt{\pi_A(i)}. \quad (11.5)$$

Proof of Lemma 11.1. Let S be the diagonal matrix with diagonal entries $s(i,i) \equiv \sqrt{\pi_A(i)}$. The matrix $\widehat{A} \equiv SAS^{-1}$ has entries $\hat{a}(i,j) \equiv \sqrt{\pi_A(i)/\pi_A(j)}a(i,j)$ and so, by (11.4), is symmetric. From the identity $\widehat{A}^m = SA^mS^{-1}$ it follows that if A is a nonnegative and primitive matrix then so is $\widehat{A}$. Finally, since A and $\widehat{A}$ are similar they have the same eigenvalues, given by (11.1). For L given by (11.3) set $\widehat{L} \equiv SLS^{-1} = (\hat{v}(i)\hat{u}(j))_{1\le i,j\le N}$ where $\hat{u} \equiv uS^{-1}$, $\hat{v} \equiv Sv$. One checks that $\hat{u}\widehat{A} = \lambda_0\hat{u}$, $\widehat{A}\hat{v} = \lambda_0\hat{v}$ and thus, since $\widehat{A}$ is symmetric, $\hat{v}\widehat{A} = \lambda_0\hat{v}$, $\widehat{A}\hat{u} = \lambda_0\hat{u}$. But this implies that $\hat{u} = \hat{v}$ and so, $\widehat{L}$ is symmetric.

Set $R \equiv A - \lambda_0 L$ and $\widehat{R} \equiv SRS^{-1} = \widehat{A} - \lambda_0\widehat{L}$. It results from the above that $\widehat{R}$ is symmetric, and since R and $\widehat{R}$ are similar they have the same eigenvalues. The proof of Lemma 11.1 now hinges on the following key facts. Given a matrix B we denote by $\sigma(B)$ its spectrum and by $\rho(B) = \max\{|\lambda| : \lambda \in \sigma(B)\}$ its spectral radius. We have:

(i) for all $m \in \mathbb{N}$, $R^m = A^m - (\lambda_0 L)^m = A^m - \lambda_0^m L$, and

(ii) $\rho(A - \lambda_0 L) \le |\lambda_1| < \rho(A) = \lambda_0$.

For a proof see (b), (e), and (h) of Lemma 8.2.7 p. 498 of [29] together with the remark below Definition 8.5.0, p. 516.

To proceed observe that $R^m = S^{-1}\widehat{R}^m S$. Thus, denoting by δ_i the unit vector with components $\delta_i(j) = 1$ if $i = j$ and $\delta_i(j) = 0$ else, it follows from (i) that

$$a^{(m)}(i,j) - \lambda_0^m v(i)u(j) = (\delta_i, R^m\delta_j) = (\delta_i, S^{-1}\widehat{R}^m S\delta_j) = \sqrt{\frac{\pi_A(j)}{\pi_A(i)}}(\delta_i, \widehat{R}^m\delta_j). \quad (11.6)$$

Summing (11.6) over j we get, by Cauchy-Schwarz's inequality and (11.2), that

$$\left|\sum_{1\le j\le N} a^{(m)}(i,j) - \lambda_0^m v(i)\right| \le \sqrt{1/\pi_A(i)}\left[\sum_{1\le j\le N}(\delta_i, \widehat{R}^m\delta_j)^2\right]^{1/2}. \quad (11.7)$$

Since $\widehat{R}$ is symmetric, $\sum_{1\le j\le N}(\delta_i, \widehat{R}^m\delta_j)^2 = (\delta_i, \widehat{R}^{2m}\delta_i)$. Now,

$$(\delta_i, \widehat{R}^{2m}\delta_i) \le \sup_{\|x\|_2=1}(x, \widehat{R}^{2m}x) \le (\rho(\widehat{R}))^{2m} = (\rho(R))^{2m} \le |\lambda_1|^{2m}. \quad (11.8)$$

The middle inequality in (11.8) follows from the minimax characterization of the largest eigenvalue of $\widehat{R}^{2m}$; the ensuing equality states that having the same eigenvalues, $\widehat{R}$ and R have same spectral radius, and the last inequality is (ii). Combining (11.7) and (11.8) proves (11.5). The proof of the lemma is done. □

11.2. **Irreducible periodic matrices with period 2.** Assume that A is an $N \times N$ nonnegative and irreducible matrix, namely, $A \ge 0$ and $(I + A)^{N-1} > 0$. Assume further that there exists a permutation matrix B such that

$$B^TAB = \begin{pmatrix} 0 & A^{+-} \\ A^{-+} & 0 \end{pmatrix} \text{ and } B^TA^2B = \begin{pmatrix} A_2^{++} & 0 \\ 0 & A_2^{--} \end{pmatrix}, \quad (11.9)$$

where the square matrices $A_2^{++} \equiv A^{+-}A^{-+}$ and $A_2^{--} \equiv A^{-+}A^{+-}$ are primitive $N^+ \times N^+$, respectively $N^- \times N^-$ matrices for some $N^+ > 0$ and $N^- > 0$ such that $N^+ + N^- = N$. This implies that A is periodic with period two, that $A_2^{\pm}$ are aperiodic and, more generally, that powers of A can be analysed in terms of powers of primitive matrices. To simplify the presentation we assume from now on that B is the identity matrix.

Part of Perron's theorem generalizes to nonnegative and irreducible matrices: denoting by λ_0 its largest eigenvalue we know that $\rho(A) = \lambda_0 > 0$, that there exist a unique (up to a normalization) and positive left (respectively, right) Perron eigenvector associated to λ_0, and that λ_0 has multiplicity one. Thus the eigenspace associated to this eigenvector is one dimensional. However the largest eigenvalue, λ_0, is not the only eigenvalue of

maximal modulus, i.e. satisfying $\rho(A) = \lambda_0$. Indeed one readily derives from (11.9) that the spectrum of A is symmetric: writing $\lambda_0 \geq \lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_{N-1}$, we have

$$\lambda_i = -\lambda_{N-i-1}, \quad 0 \leq i \leq N-1, \tag{11.10}$$

and if $Aw = \lambda_i w$, $w = (w^+, w^-)$ where $w^\pm$ denote vectors in $\mathbb{R}^{N^\pm}$, then $A\bar{w} = -\lambda_{N-i-1}\bar{w}$ where $\bar{w} = (w^+, -w^-)$. Thus

$$\rho(A) = \lambda_0 = |\lambda_{N-1}| > |\lambda_i|, \quad 1 \leq i \leq N-2. \tag{11.11}$$

Using in addition that the matrices $A_2^\pm$ are primitive, one deduces that there exist unique left and right Perron eigenvectors, u and v,

$$uA = \lambda_0 u, \;\; u = (u^+, u^-) > 0, \tag{11.12}$$

$$Av = \lambda_0 v, \;\; v = (v^+, v^-) > 0, \tag{11.13}$$

normalized such that

$$\textstyle\sum_{1\leq i\leq N^+} u^+(i) = \sum_{1\leq i\leq N^-} u^-(i) = \frac{1}{2}, \tag{11.14}$$

$$\textstyle\sum_{1\leq i\leq N^+} u^+(i)v^+(i) = \sum_{1\leq i\leq N^-} u^-(i)v^-(i) = \frac{1}{2}, \tag{11.15}$$

and that, setting $\bar{u} = (u^+, -u^-)$ and $\bar{v} = (v^+, -v^-)$, $\bar{u}A = -\lambda_0\bar{u}$ and $A\bar{v} = -\lambda_0\bar{v}$.

With obvious notation define

$$L = (v(i)u(j))_{1\leq i,j\leq N}, \;\; \overline{L} = (\bar{v}(i)\bar{u}(j))_{1\leq i,j\leq N}. \tag{11.16}$$

The next lemma establishes that

$$\lim_{m\to\infty} (\lambda_0^{-1}A)^{2m} = (L + \overline{L}), \;\; \lim_{m\to\infty} (\lambda_0^{-1}A)^{2m+1} = (L - \overline{L}), \tag{11.17}$$

and gives bounds on the speed of convergence under the same symmetry assumption as in Lemma 11.1. Note that $L + \overline{L}$ has (i,j)-th entry $2v^\pm(i)u^\pm(j)$ if $1 \leq i,j \leq N^\pm$ and zero else, and that $L - \overline{L}$ has (i,j)-th entry $2v^\pm(i)u^\mp(j)$ if $1 \leq i \leq N^\pm$ and $1 \leq j \leq N^\mp$, and zero else.

Lemma 11.2. *Assume that there exists a vector $\pi_A > 0$ such that $\sum_{1\leq i\leq N} \pi_A(i) = 1$, and*

$$\pi_A(i)a(i,j) = \pi_A(j)a(j,i) \quad \textit{for all } 1 \leq i,j \leq N. \tag{11.18}$$

Then for all $m \in \mathbb{N}$ we have: for all $1 \leq i \leq N^\pm$,

$$\left|\textstyle\sum_{1\leq j\leq N^\pm} a^{(2m)}(i,j) - \lambda_0^{2m}v(i)\right| \leq |\lambda_1|^{2m}/\sqrt{\pi_A(i)}, \tag{11.19}$$

and for all $1 \leq i \leq N^\pm$,

$$\left|\textstyle\sum_{1\leq j\leq N^\mp} a^{(2m+1)}(i,j) - \lambda_0^{2m+1}v(i)\right| \leq |\lambda_1|^{2m+1}/\sqrt{\pi_A(i)}. \tag{11.20}$$

Proof of Lemma 11.2. One easily checks that for all $m \in \mathbb{N}$, $L^m = L$, $\overline{L}^m = \overline{L}$, $L\overline{L} = \overline{L}L = 0$, $LA = AL = 0$, and $\overline{L}A = A\overline{L} = 0$. Thus, setting $R = A - \lambda_0(L - \overline{L})$ we have, for all $m \in \mathbb{N}$,

(i) $R^{2m} = A^{2m} - (\lambda_0(L - \overline{L}))^{2m} = A^{2m} - \lambda_0^{2m}(L + \overline{L})$,

(ii) $R^{2m+1} = A^{2m+1} - (\lambda_0(L - \overline{L}))^{2m+1} = A^{2m+1} - \lambda_0^{2m+1}(L - \overline{L})$,

and, using (11.11),

(iii) $\rho(A - \lambda_0(L - \overline{L})) \leq \lambda_1 < \rho(A) = \lambda_0$.

Using this the proof of Lemma 11.2 is a simple rerun of the proof of Lemma 11.1. □

REFERENCES

[1] D. J. Aldous and M. Brown. Inequalities for rare events in time-reversible Markov chains. I. In *Stochastic inequalities (Seattle, WA, 1991)*, volume 22 of *IMS Lecture Notes Monogr. Ser.*, pages 1–16. Inst. Math. Statist., Hayward, CA, 1992.

[2] G. Ben Arous, A. Bovier, and J. Černý. Universality of the REM for dynamics of mean-field spin glasses. *Comm. Math. Phys.*, 282(3):663–695, 2008.

[3] G. Ben Arous, A. Bovier, and V. Gayrard. Aging in the random energy model. *Phys. Rev. Letts.*, 88(8):087201, 2002.

[4] G. Ben Arous, A. Bovier, and V. Gayrard. Glauber dynamics of the random energy model. I. Metastable motion on the extreme states. *Comm. Math. Phys.*, 235(3):379–425, 2003.

[5] G. Ben Arous, A. Bovier, and V. Gayrard. Glauber dynamics of the random energy model. II. Aging below the critical temperature. *Comm. Math. Phys.*, 236(1):1–54, 2003.

[6] G. Ben Arous and J. Černý. The arcsine law as a universal aging scheme for trap models. *Comm. Pure Appl. Math.*, 61(3):289–329, 2008.

[7] G. Ben Arous and V. Gayrard. Elementary potential theory on the hypercube. *Electron. J. Probab.*, 13:no. 59, 1726–1807, 2008.

[8] G. Ben Arous and O. Gün. Universality and extremal aging for dynamics of spin glasses on subexponential time scales. *Comm. Pure Appl. Math.*, 65(1):77–127, 2012.

[9] G. Bennett. Probability inequalities for the sum of independent random variables. *J. Amer. Statist. Assoc.*, 57(297):33–45, 1962.

[10] B. Bollobás, Y. Kohayakawa, and T. Luczak. On the evolution of random Boolean functions. In *Extremal problems for finite sets (Visegrád, 1991)*, volume 3 of *Bolyai Soc. Math. Stud.*, pages 137–156. János Bolyai Math. Soc., Budapest, 1994.

[11] J.-P. Bouchaud. Weak ergodicity breaking and aging in disordered systems. *J. Phys. I (France)*, 2:1705–1713, september 1992.

[12] J.-P. Bouchaud, L. Cugliandolo, J. Kurchan, and M. Mézard. Out of equilibrium dynamics in spin-glasses and other glassy systems. In A. P. Young, editor, *Spin glasses and random fields*. World Scientific, Singapore, 1998.

[13] J.-P. Bouchaud and D. S. Dean. Aging on Parisi's tree. *J. Phys I(France)*, 5:265, 1995.

[14] A. Bovier and V. Gayrard. Convergence of clock processes in random environments and ageing in the p-spin SK model. *Ann. Probab.*, 41(2):817–847, 2013.

[15] A. Bovier, V. Gayrard, and A. Švejda. Convergence to extremal processes in random environments and extremal ageing in SK models. *Probab. Theory Related Fields*, 157(1-2):251–283, 2013.

[16] J. Černý and V. Gayrard. Hitting time of large subsets of the hypercube. *Random Structures Algorithms*, 33(2):252–267, 2008.

[17] Y. S. Chow and H. Teicher. *Probability theory*. Springer Texts in Statistics. Springer-Verlag, New York, second edition, 1988. Independence, interchangeability, martingales.

[18] H. Cramér. *Mathematical Methods of Statistics*. Princeton Mathematical Series, vol. 9. Princeton University Press, Princeton, N. J., 1946.

[19] B. Derrida. Random-energy model: limit of a family of disordered models. *Phys. Rev. Lett.*, 45(2):79–82, 1980.

[20] B. Derrida. A generalization of the random energy model which includes correlations between energies. *J. Physique Lett.*, 46:401–407, 1985.

[21] P. Diaconis and D. Stroock. Geometric bounds for eigenvalues of Markov chains. *Ann. Appl. Probab.*, 1(1):36–61, 1991.

[22] R. Durrett and S. I. Resnick. Functional limit theorems for dependent variables. *Ann. Probab.*, 6(5):829–846, 1978.

[23] L. Fontes and P. Lima. Convergence of symmetric trap models in te hypercube. In V. Sidoravicius, editor, *New Trends in Mathematical Physics*, pages 285–297. Springer Netherlands, 2009.

[24] L. R. G. Fontes, M. Isopi, Y. Kohayakawa, and P. Picco. The spectral gap of the REM under Metropolis dynamics. *Ann. Appl. Probab.*, 8(3):917–943, 1998.

[25] V. Gayrard. Aging in reversible dynamics of disordered systems. II. Emergence of the arcsine law in the random hopping time dynamics of the REM. preprint, 2010. arXiv:1008.3849.

[26] V. Gayrard. Convergence of clock process in random environments and aging in Bouchaud's asymmetric trap model on the complete graph. *Electron. J. Probab.*, 17:no. 58, 33, 2012.

[27] R. J. Glauber. Time-dependent statistics of the Ising model. *J. Mathematical Phys.*, 4:294–307, 1963.
[28] M. Goldstein. Viscous liquids and the glass transition: A potential energy barrier picture. *The Journal of Chemical Physics*, 51(9):3728–3739, 1969.
[29] R. A. Horn and C. R. Johnson. *Matrix analysis*. Cambridge University Press, Cambridge, 1985.
[30] P. Mathieu and J.-C. Mourrat. Aging of asymmetric dynamics on the random energy model. arxiv:1212.5117, 2013.
[31] N. Metropolis, A. Rosenbluth, M. Rosenbluth, A. Teller, and E. Teller. Equations of state calculations by fast computing machines. *J. of Chemical Physics*, 21 (6):1087–1092, 1953.
[32] A. Sinclair and M. Jerrum. Approximate counting, uniform generation and rapidly mixing Markov chains. *Inform. and Comput.*, 82(1):93–133, 1989.
[33] E. Vincent. Ageing, rejuvenation and memory: the example of spin-glasses. In *Ageing and the glass transition*, volume 716 of *Lecture Notes in Phys.*, pages 7–60. Springer, Berlin, 2007.

V. GAYRARD, I2M, AIX-MARSEILLE UNIVERSITÉ, 39, RUE F. JOLIOT CURIE, 13453 MARSEILLE CEDEX 13, FRANCE

E-mail address: veronique.gayrard@math.cnrs.fr, veronique@gayrard.net